

2014 Annual Report



TABLE OF CONTENTS

ILLINOIS AND KASKASKIA RIVER WATERSHEDS CREP ELIGIBLE AREA	3
EXECUTIVE SUMMARY	4
Illinois CREP goals.....	4
Illinois CREP Timeline	5
Recent Outreach, Stewardship and Monitoring	6
Program Expenditures.....	7
CREP Enrollment and Financial Figures	8
Pollutant Load Reduction Report.....	9
PARTNER UPDATES	11
Illinois Environmental Protection Agency	11
Other Illinois EPA programs that complement CREP	11
Excess Nutrients: A High Profile Water Quality Issue	12
Current Management Approaches and Issues.....	13
What U.S. EPA Expects	13
Illinois Department of Agriculture	14
Illinois Department of Natural Resources	14
Illinois Recreational Access Program.....	14
Partners for Conservation	15
Natural Resources Conservation Service (NRCS).....	16
EQIP	16
EQIP’s Forestry Efforts.....	16
Wetland Reserve Program.....	16
US Fish and Wildlife Service	16
Partners for Fish and Wildlife	16
Partners for Fish and Wildlife Program in Southern Illinois	17
Illinois Farm Bureau	18
Association of Illinois Soil and Water Districts (AISWCD).....	18
National Great Rivers Research and Education Center	19
ONGOING INITIATIVES	20
Monitoring and Assessment of Aquatic Life in the Kaskaskia River for Evaluating IDNR Private Lands Programs – Illinois Natural History Survey	20

Establishing a Biological Monitoring Program for CREP to Assess the Conservation Practices and Wildlife Habitat on Property Enrolled – Illinois Natural History Survey.....	21
Monitoring and Evaluation of Sediment and Nutrient Delivery to the Illinois and Kaskaskia Rivers – Illinois State Water Survey.....	21
Habitat Grant Project – National Wild Turkey Federation	22

APPENDICES


- Appendix A: Monitoring and Assessment of Aquatic Life in the Kaskaskia River for evaluating IDNR Private Lands Programs:
Illinois Natural History Survey

- Appendix B: Establishing a biological monitoring program for CREP to assess the conservation practices and wildlife habitat on property enrolled
Illinois Natural History Survey

- Appendix C: Monitoring and Evaluation of Sediment and Nutrient Delivery to the Illinois River:
Illinois River Conservation Reserve Enhancement Program (CREP)
Illinois State Water Survey

- Appendix D: Sediment and Nutrient Monitoring at Selected Watersheds within the Kaskaskia River Watershed for Evaluating the Effectiveness of the Illinois River Conservation Reserve Enhancement Program (CREP)
Illinois State Water Survey

ILLINOIS AND KASKASKIA RIVER WATERSHEDS CREP ELIGIBLE AREA



EXECUTIVE SUMMARY

The Illinois Conservation Reserve Enhancement Program (CREP) is a state incentive program tied to the Federal Conservation Reserve Program (CRP). CREP provides long term environmental benefits by allowing 232,000 acres of eligible environmentally sensitive lands within the Illinois and Kaskaskia River Watersheds to be restored, enhanced and protected over periods ranging from 15 years to perpetuity. CREP continues to be driven by locally led conservation efforts, which is evident by increased landowner support. This program is a prime example of how partnerships between landowners, governmental entities, and non-governmental organizations can network to address watershed quality concerns.

Having worked hand in hand with USDA over the years, Illinois CREP has been instrumental in facilitating the ongoing restoration and management efforts within the Illinois and Kaskaskia River Watersheds. To achieve the goal of improving water quality within the targeted watersheds CREP has utilized a variety of Best Management Practices (BMP's) designed to protect and restore miles of riparian corridors. CREP is one of many tools used by IDNR conservation partners to implement the IDNR Illinois Comprehensive Wildlife Action Plan, which provides a framework for the restoration of critical habitats, increasing plant diversity and expanding habitat for species in greatest need of conservation on an agricultural dominated landscape.

Illinois CREP continues to be a successful and very popular program. Since CREP's inception in 1998, 140,934.0 acres have been enrolled in Federal CREP contracts at an average rental rate of \$188.5 per acre. The State has been successful in executing 1,366 CREP easements protecting 87,466 acres.

Illinois CREP goals

The goals for the Illinois CREP were revised in 2010 to reflect the expansion into the Kaskaskia River Basin and to highlight the importance of the connection to the Mississippi River and the Gulf of Mexico. The goals of the program are:

- Reduce the amount of silt and sedimentation entering the main stem of the Illinois and the Kaskaskia Rivers by 20 percent;
- Reduce the amount of phosphorus and nitrogen in the Illinois River and Kaskaskia River by 10 percent;
- Increase by 15 percent, the populations of waterfowl, shorebirds, nongame grassland birds, and State and Federally listed threatened and endangered species such as bald eagles, egrets, and herons;
- Increase the native fish and mussel stocks by 10 percent in the lower reaches of the Illinois River (Peoria, LaGrange, and Alton reaches) and Kaskaskia River; and
- Help meet the Federal goals to reduce nitrogen loading to the Mississippi River and the Gulf of Mexico, thereby helping to reduce hypoxia in the Gulf of Mexico.

Illinois CREP Timeline


1998-2000



CREP is a federal-state program that was created by a Memorandum of Agreement (MOA) between the U.S. Department of Agriculture, the Commodity Credit Corporation, and the State of Illinois in March 1998. Enrollments into this program began on May 1, 1998. The MOU was amended several times during the early years to clarify terms, increase the number of practices offered, and to expand the eligible area.

In 2005 the IDNR, in cooperation with other conservation partners, initiated the implementation of The Illinois Comprehensive Wildlife Action Plan (ICWAP). The ICWAP's goals are to use consistent science-based natural resource management principles, to increase the amount and quality of habitat available to support Illinois' native plant and animal species and other game species; promote their population viability, and regulate the recreational, commercial, and scientific utilization of those species; to ensure their long-term persistence and abundance and provide for their appreciation and enjoyment by future generations of Illinoisans while also expanding the frontiers of natural resource management. CREP easements which lie within the ICWAP's priority areas will provide long term protection of quality habitats identified by the ICWAP's goals.


1998-2007



Due to insufficient State funds the Illinois CREP was temporarily closed to open enrollment in November 2007. However, monitoring and land stewardship continued.

In October 2010, after overwhelming public support The Illinois General Assembly appropriated \$45 million to reopen and expand CREP to include the Kaskaskia River Watershed. The USDA, Commodity Credit Corporation, and the State of Illinois subsequently amended their Memorandum of Agreement (MOA) to include the Kaskaskia River Watershed with the Illinois River Watershed.

1000 2014
1998 - 2013



Since 2010 a total of 149 state easements have been approved in the Kaskaskia and Illinois River Watersheds totaling 12,608 acres, the Kaskaskia River Watershed totaling 4,610 acres and Illinois Watershed totaling 7,998 acres. The average acreage per enrollment is 84.62 acres.

Since the program started in 1998, landowners have voluntarily enrolled 87,466 acres in CREP through 1,366 easements to help improve and restore natural habitats in the Illinois CREP eligible area. In the last year alone (10/1/2013 – 9/30/2014) 42 state easements were closed protecting 3,303 acres overall, 1,209 acres in the Kaskaskia River Watershed and 2,094 acres in the Illinois River Watershed.

Recent Outreach, Stewardship and Monitoring

The county Soil and Water Conservation Districts (SWCD) within the CREP area are the driving force spearheading CREP on the local level.

The Illinois Department of Natural Resources (IDNR) has partnered with the Illinois Environmental Protection Agency (IEPA) and the Association of Illinois Soil and Water Conservation Districts (AISWCD) to hire six (6) CREP Resource Specialists. These specialists are dedicated to counties primarily in the Illinois River Watershed to assist the SWCD's with landowner outreach and enrollment. IDNR has also partnered with the National Great Rivers Research and Education Center (NGRREC) who were awarded a National Fish and Wildlife Fund Grant to hire four (4) Land Conservation Specialists to market CREP and assist the districts as needed in counties primarily in the Kaskaskia River Watershed.

The State continues to monitor and evaluate sediment and nutrient delivery to the Illinois River. Nutrient and sediment data have been collected since the program's inception in 1999. According to the Illinois State Water Survey's (ISWS) recent data indicates that both sediment and nutrient delivery to the Illinois River has gradually either stabilized or decreased as a result of the implementation of BMP's in the Illinois River watershed. The most significant outcome has been the slow decreasing trend of nitrate-N yield from major tributary watersheds.

The IDNR is working with the University of Illinois' Critical Trends Assessment Program (CTAP) staff to maintain a biological monitoring program for CREP to assess the conservation practices and wildlife habitat on property enrolled in CREP. CTAP samples the bird communities of forests, grasslands, and wetlands using point-count based methods. During data collection, the presence and abundance of each species seen or heard during the count period is recorded.

The IDNR is also working with Illinois Natural History Survey to maintain a basin-wide monitoring and assessment program for wadeable streams in the Kaskaskia River. Baseline information on aquatic macroinvertebrates (EPT), freshwater mussels, and fish have been collected at selected reaches using a stratified random sampling design to characterize conditions throughout the watershed and provide for long-term trends assessments. Populations of selected species are monitored in focal reaches associated with high biological diversity (BSS reaches) or sensitive taxa (enhanced DO reaches, SGNC).

Program Expenditures

The Memorandum of Agreement (MOA) for the Illinois CREP details the formula to determine the overall costs of the program: total land retirement costs (which will include the CRP payments made by the Commodity Credit Corporation (CCC) and the easement payments or the bonus payments made by Illinois), the total reimbursement for conservation practices paid by the CCC and Illinois, the total costs of the monitoring program, and the aggregate costs of technical assistance incurred by Illinois for implementing contracts and easements and a reasonable estimate of the cost incurred by the State to develop conservation plans.

Since the CRP contract payments are annual payments spread out over 15 years, a 3.30 percent net present value (NPV) discount rate (per MOA) was used to compare the CRP payments to the State Easement payments.

Per the current agreement, the State of Illinois must contribute 20% of the total program costs. Based on USDA reports at <https://arcticocean.sc.egov.usda.gov/> IDNR contributed 23.86% of the total program costs based on the following calculations;

\$398,490,885.00 (15 years x 140,934.0 acres x 188.5 avg. rental rate = \$398,490,885.00) given to IDNR by USDA FSA* was amended by IDNR to reflect the 2013 re-enrollment of expired CRP acres with perpetual CREP easements (\$1,528,283.64), expiring 1998 CRP contracts (26,445.2 acres x \$156.00 1998 avg. rental rate = \$4,125,451.20) and expiring 1999 CRP contracts (28,772.4 acres x \$155.00 1999 avg. rental rate = \$4,451,972.00).

2014 USDA Report	\$398,490,885.00
2013 USDA CREP re-enrollments	(\$1,528,283.64)
1998 expired contracts	(\$4,125,451.20)
<u>1999 Expired Contracts</u>	<u>(\$4,451,972.00)</u>
Amended total	\$388,385,178.16

*- End of September-2014 Summary of active Contracts by Program Year, CRP – Monthly Contracts Report, Program year 1998 - 2014

CREP Enrollment and Financial Figures

Illinois CREP 1998- Sept 30, 2014	
Number of Federal Contracts - 7,387	Total Federal Acres - 140,934.0
Number of State Easements - 1,366	Total State Protected Acres - 87,466

Payments 1998 - Sept 30, 2014	IDNR	USDA *	USDA (NPV 3.30%) **
Acres Enrolled through Sept, 30 2013		140,934.00	
Total Life of Contract Rent (15 Yrs)		\$388,385,178.16	\$238,648,014.18
Cost Share		\$18,645,723.19	\$18,645,723.19
Monitoring	\$5,382,728.54		
IEPA CREP Assistants IEPA 319	\$2,180,665.94		
Illinois State Enrollments	\$67,966,556.39		
IDNR In-Kind Services	\$5,182,832.55		

CREP Match 1998 – Sept 30, 2014	IDNR	IDNR/USDA *	IDNR/USDA **
USDA Total		\$407,030,901.35	\$257,293,737.37
IDNR Total	\$80.629,701.26	\$80.629,701.26	\$80.629,701.26
Program Total		\$487,660,602.61	\$337,923,438.63
% of IDNR Match		17%	23.86%

* September 2014 Payment and Practice Summary of active CREP Contracts by Program Year, CRP – Monthly Contracts Report


https://arcticocan.sc.egov.usda.gov/CRPReport/monthly_report.do?method=displayReport&report=September-2014-ActiveCrepContractsSummaryByProgramYear-17

** Net Present Value (NPV) http://www.whitehouse.gov/omb/circulars/a094/a94_appx-c.html


Pollutant Load Reduction Report

To better understand CREP's impact on water quality, a spatially based pollution load model was developed to estimate field level pollutant loading from Nitrogen, Phosphorus and Sediment. By analyzing soils, land-use and precipitation data the model provides both annual and storm event loading for individual land parcels within the Illinois River basin. Accepted equations for calculating runoff and soil erosion are integrated into the model to provide realistic estimations of the quantity and distribution of pollution loading throughout. Data collected between years 2002 and 2011 were used for model calibration of rainfall values and for evaluating in-stream water quality. Final model results for annual pollution loading are calibrated to existing in-stream water quality


Total Annual Nitrogen lbs/year



Total Annual Phosphorus lbs/year



Total Annual Sediment tons/year



Pollutant Loading **without** CREP on ground

	Total	Per Acre
Total Annual N (lbs/yr)	237,904	4.73
Total Annual P (lbs/yr)	118,952	2.36
Total Annual Sediment (tons/yr)	118,952	2.36

Pollutant Loading **with** CREP on ground

	Total	Per Acre
Total Annual N (lbs/yr)	4708	0.093
Total Annual P (lbs/yr)	2354	0.47
Total Annual Sediment (tons/yr)	2354	0.47

Table is reporting on acres that have come out of crop production, and into CREP. It does not account for the remaining acres in the watershed.

Approximately 87,000 acres of State CREP were enrolled since the program opened. Within this total 50,300 acres of crop conversion will prevent following pollutants from entering the Illinois and Kaskaskia Rivers:

- 233,196 lbs of Nitrogen per year
- 116,598 lbs of Phosphorus per year
- 116,598 tons of Sediment per year

This one-time investment in a CREP easement will reduce non-point source inputs to the Mississippi River basin by the following amounts over a **15 year period**:

- 3,497,940 lbs of Nitrogen
- 1,748,970 lbs of Phosphorus
- 1,748,970 tons of Sediment

This one-time investment in a CREP easement will reduce non-point source inputs to the Mississippi River basin by the following amounts over a **100 year period**:

- 23,319,600 lbs of Nitrogen
- 11,659,800 lbs of Phosphorus
- 11,659,800 tons of Sediment

Load Reduction Methodology

1) Predicted Average Annual Soil Loss (ton/acre/yr) = USLE

Before Treatment = $R * K * LS * C$ before After Treatment = $R * K * LS * C$ After

- | | |
|-----------------------|------------------------------|
| • A slope - 0.24 | • Permanent seedings - 0.003 |
| • B,C, D slope - 0.12 | • Tree plantings - 0.004 |
| • E, F slope - 0.06 | • Wetlands - 0.0001 |

2) Sediment Delivery Ratio = $((\text{acres}/640)^{-0.125}) * 0.42$

3) Sediment Delivery (ton/acre/year) Before = (“before” Predicted Avg Annual Soil Loss * Sediment Delivery Ratio)

4) Sediment Delivery (ton/acre/year) After = (“after” Predicted Avg Annual Soil Loss * Sediment Delivery Ratio)

5) Estimated Sediment Load Reduction (ton/yr)

= contributing acres * (result from #3 – result from #4)

6) As per EPA recommendation, the following averages are used for N and P

- average for nutrients would be for every Ton of soil saved a corresponding amount of P would be 1 lb and for N it would be 2 lbs.

PARTNER UPDATES

Illinois Environmental Protection Agency

One of the key missions of Illinois Environmental Protection Agency (EPA) is to monitor and protect the water resources of Illinois; these resources are relied upon for drinking water, fishing, transportation and recreational use and other environmental and economic benefits. One of the most dramatic improvements in water quality that Illinois EPA has documented has taken place on the Illinois River.

Illinois EPA has eight Ambient Water Quality Monitoring Sites on the main channel of the Illinois River. Water chemistry is collected at these sites nine times per year. There are approximately 475 Intensive Basin Survey Sites in the Illinois and Kaskaskia River watersheds. These sites are monitored "intensively" once every five years. The monitoring includes water chemistry, macro-invertebrates, fish, habitat, sediment and at some sites fish tissue contaminants are collected. This information is cooperatively collected with the Illinois Dept. of Natural Resources, a partnership that began many years ago and continues annually.

The monitoring shows that the Illinois River mainstream water quality has improved significantly since the passage of the Federal Clean Water Act in 1972. Early improvements were due primarily to point source controls, such as additional treatment requirements and limits on discharges from wastewater treatment plants. The majority of water quality improvements over the last fifteen years have been from the implementation of nonpoint source management programs that reduce urban and agricultural runoff, and programs such as CREP.

As reported by the Illinois EPA in their 2012 Integrated Report, of the *stream miles assessed* in the Illinois River Basin for Aquatic Life Use Support attainment, 67.8% were reported as —Good, 27.6% as —Fair, and 4.6% as —Poor. This compares to statewide figures of 60.8% —Good, 34.0% —Fair, and 5.2% —Poor.

Illinois EPA continues to participate on the State CREP Advisory Committee and continues to provide financial assistance to local soil and water conservation districts so they can assist landowner enrollment into CREP. Since 1999, more than \$1,958,000 of Section 319 grant funds have been spent to hire and train personnel responsible for outreach and the enrollment process.

The benefits derived through this financial support is not only efficiency in the sign-up process to increase CREP enrollment, but it also allows the existing SWCD and NRCS staff to continue to implement the other conservation programs so desperately needed to improve water quality in the Illinois and Kaskaskia River watersheds.

Other Illinois EPA programs that complement CREP include:

Section 319: Since 1990, the IEPA has implemented 285 Clean Water Act Section 319 projects within the Illinois and Kaskaskia River Watersheds. The Agency receives these federal funds from USEPA to identify and administer projects to prevent nonpoint source pollution. These projects include watershed management planning; best management practices implementation and outreach efforts. Illinois EPA has dedicated over \$64 million with another \$56 million of local and state funds for total project costs of nearly \$120 million towards these projects to help improve the health of the Illinois and Kaskaskia Rivers, their tributaries and ultimately the Mississippi River and Gulf of Mexico. Hundreds of conservation practices have been installed in the Illinois and Kaskaskia River watersheds by dozens of our partners through the Section 319 program. Traditional practices such as terraces and waterways are dotting the landscape along with porous pavement parking lots, green roofs and miles of rural and urban stabilized streambank.

Since 1990, the 319 NPS program, through on the ground implementation can show load reductions in the Illinois and Kaskaskia River watersheds of: 702,897 lbs. of nitrogen, 328,487 pounds of phosphorus, and 287,757 tons of sediment per year, each and every year since the Best Management Practices were implemented as a result of 319 grant projects between IEPA and our local partners, in both the private and government sectors. The IEPA invites you to visit <http://water.epa.gov/polwaste/nps/success319/> for a sample of Illinois' 319 success stories.

IGIG: Since 2011, the Illinois EPA has implemented 28 Illinois Green Infrastructure Grant Program for Stormwater Management (IGIG) projects within the Illinois and Kaskaskia River watersheds. IGIG is administered by the Illinois EPA. Grants are available to local units of government and other organizations to implement green infrastructure best management practices (BMPs) to control stormwater runoff for water quality protection in Illinois. Projects must be located within a Municipal Separate Storm Sewer System (MS4) or Combined Sewer Overflow (CSO) area. Funds are limited to the implementation of projects to install BMPs. Illinois EPA has dedicated over \$10 million with another \$4 million of local funds for total project costs of over \$14 million towards these projects to help improve water quality in the Illinois and Kaskaskia River watersheds.

Construction Site Inspection Program: Illinois EPA continues to implement a program in partnership with nineteen soil and water conservation districts covering twenty-two counties. Those partners located with the Illinois and Kaskaskia River watersheds include the Champaign, DeKalb, Jersey, Kane/DuPage, Kankakee, Kendall, Knox, Macon, Madison, McHenry/Lake, Monroe, North Cook, Peoria, St. Clair, and Will/South Cook County Soil and Water Conservation Districts. District staff complete on-site NPDES Construction Stormwater Permit inspections and provide technical assistance in implementing best management practices to minimize runoff to nearby water bodies. This program is a natural fit for properly developing acreage that does not qualify for CREP.

Total Maximum Daily Load (TMDL): TMDLs are a tool that we use to restore impaired watersheds so that their waters will meet Water Quality Standards and Full Use Support for those uses that the water bodies are designated. A TMDL looks at the identified pollutants and develops, through water quality sampling and modeling, the amount or load reductions needed for the water body to meet its designated uses. USEPA has approved 265 completed TMDL evaluations and Illinois EPA is currently developing another 218 TMDLs in the Illinois and Kaskaskia River watersheds.

Partners for Conservation: A total of 72 lake monitoring (study) or protection/restoration projects have been conducted in the Illinois and Kaskaskia River watersheds via the Illinois EPA's Illinois Clean Lakes Program and Priority Lake and Watershed Implementation Program. Over \$11.8 million of local and state funds have been allocated for these efforts.

Excess Nutrients: A High Profile Water Quality Issue

The impact of excess nitrogen and phosphorus in rivers, lakes, streams and the Gulf of Mexico has become a very high profile water quality issue. Under the right conditions, nutrients can cause excessive algal blooms, low oxygen and nuisance conditions that adversely impact aquatic life, drinking water and recreational uses of the water. The Illinois Environmental Protection Agency has identified many waterbodies in the state with these problems.

Nitrogen and phosphorus come from municipal wastewater treatment, urban stormwater, row crop agriculture, livestock production, industrial wastewater and combustion of fossil fuels. In other words, most aspects of modern society contribute to this pollution problem. The proportion of loading to a particular waterbody from these sources varies from watershed to watershed, with point sources and urban storm water being most important in urbanized watersheds and row crop and/or livestock production being predominant contributors in agricultural watersheds.

Illinois EPA has several on-going efforts concerning nutrients. The first is identification of eight sub-watersheds that are considered our —Nutrient Priority Watersheds. Six of the eight designated watersheds are in the Illinois River Basin, they are: Lake Bloomington, Lake Evergreen, Lake Decatur, Lake Springfield, Vermilion River (Illinois Basin) and Lake Mauvaise Terre. Each of these watersheds has a Total Maximum Daily load developed or being developed for one or two nutrient pollutants (nitrogen and phosphorus). The second nutrient effort that the agency is partnering with is a program called —KIC 2025 (www.kic2025.org). —KIC by 2025 will seek to educate the agricultural sector, dedicate significant resources toward research to reduce nutrient losses and enhance nutrient efficiency, educate suppliers and farmers, and measure the adoption of in-field practices to enhance nutrient stewardship beginning in priority watersheds and expanding over years to a state-wide nutrient stewardship program. The agency is also involved in the Mississippi River Basin Initiative in the Indian Creek Watershed (Livingston County, Vermilion-Illinois Basin). The Agency is providing technical assistance and monitoring in the Bureau Creek project and is proving funds for significant outreach in the Indian Creek Watershed along with growing season weekly samples and monthly sampling the rest of the year. Lastly, the Agency, Illinois Department of Agriculture and a group of stakeholders recently finished the Illinois Nutrient Loss Reduction Strategy document that sets a path for reducing nutrient losses on Illinois lands to downstream waters. This document, in response to the Gulf Hypoxia Task Force and U.S. EPA leadership will guide the Agency as we address nutrient concerns in the future.

In conclusion, the Illinois and Kaskaskia River basins are a valuable resource that we are working hard to protect and restore. Illinois EPA will continue long-term monitoring of the rivers and their watersheds and will continue to pursue funds to help implement CREP and other water quality restoration and protection projects and to work with citizen groups and local government and industry to continue the progress we all have made.

Current Management Approaches and Issues

The Clean Water Act framework requires: the establishment of water quality standards that protect aquatic life and/or other beneficial uses of the water; monitoring and assessment to determine attainment of standards; listing of waters not attaining and development of Total Maximum Daily Loads (TMDL) to limit pollution to those water bodies.

TMDL load limits are required to be implemented through National Pollutant Discharge Elimination System permits, which address point sources—municipal or industrial wastewater dischargers. Management of non-point source pollution is through voluntary implementation of best management practices (BMP) contrary to point sources which are regulated through permit limits.

Cost-share incentives to implement/install BMPs include federal Conservation Reserve Program and state Conservation Reserve Enhancement Program, state Partners for Conservation Program, various Farm Bill conservation programs and Section 319 non-point source management grants. The federal Farm Bill programs, though relatively well-funded, are not consistently targeted at water quality improvement, nutrient reduction or locations most in need of BMPs.

There are various other efforts through state agricultural groups, industry and non-profit organizations to promote the use of agricultural BMPs, but these efforts are not consistently coordinated nor targeted to particular watersheds. In addition, the degree of implementation of key nutrient-related BMPs is not comprehensively quantified or mapped, so the collective status of BMP implementation in the state is unknown.

Available data do indicate that Illinois producers are not over-applying fertilizers or manure and that the traditional suite of conservation practices will not be adequate to achieve such large reductions. Absent the development of an economically viable third crop such as a perennial for biofuels, the costs to significantly reduce nutrient losses from agriculture could be billions of dollars.

New and expanding major (one million gallons per day or greater design flow) municipal sewage treatment plants and some sewage treatment plants discharging to certain lake watersheds are required by Illinois Pollution Control Board regulations to limit total phosphorus to 1.0 mg/L on a monthly average basis. Plants currently achieving this level of phosphorus reduction represent 9% of the approximately 900 municipal discharges in the state. However, of the 214 major municipals discharges, whose effluent constitutes a large majority of the phosphorus loading from point sources, 25% are required to remove phosphorus. Requiring phosphorus removal from the minor facilities would be very costly for customers on a per capita basis and would represent a relatively small portion of the total point source phosphorus discharged. Therefore at this time minor facilities will not be targeted for reducing phosphorus discharge.

What U.S. EPA Expects

U.S. EPA expects states to establish numeric water quality standards for phosphorus and nitrogen and to carry out the other pieces of the Clean Water Act framework, as appropriate. U.S. EPA's Inspector General issued a finding in 2009 that U.S. EPA had not done enough to get state numeric nutrient water quality standards established. In response, U.S. EPA has developed a —corrective action plan which includes a commitment to identify states where federal promulgation of nutrient water quality standards is required. U.S. EPA has been petitioned and sued by various environmental groups for failure of states to establish numeric nutrient standards, so there is mounting pressure on U.S. EPA and states to address nutrients by developing numeric nutrient water quality standards.

States have concerns on the issue of numeric nutrient water quality standards. They raise two main points:

1. There is not a straightforward relationship between nutrient concentration in the water and adverse effects, so a statewide —one size fits all standard that meets the test of scientific defensibility is almost unachievable; and
2. The Clean Water Act programs are effective for point sources but do not assure reductions from non-point sources that are often the predominant contributors of nutrients in a particular watershed.

Through Illinois' Nutrient Loss Reduction Strategy the Agency has continued its commitment to using a science based approach to developing water quality standards.

Illinois Department of Agriculture

The Illinois Department of Agriculture (IDOA) administers numerous soil and water conservation programs that produce environmental benefits in the Illinois River Watershed. In FY13, the Partners for Conservation Program (PFC), administered by IDOA, allocated over \$296,000 to 40 counties that have significant agricultural acreage in the Illinois River Watershed for cost-sharing the installation of upland soil and water conservation practices. With the assistance from County Soil and Water Conservation Districts (SWCDs), the PFC provides up to 70% of the cost of constructing conservation practices that reduce soil erosion and protect water quality.

Conservation practices eligible for partial funding under the PFC include terraces, grassed waterways, water and sediment control basins, grade stabilization structures and nutrient management plans. A total of 204 projects have been completed with significant environmental benefits to the Illinois River Basin during the last 2 fiscal years. These conservation projects were constructed with funding of nearly \$478,000 and are responsible for bringing soil loss to tolerable levels on hundreds of acres of land. This translates into over 6,212 fewer tons of soil loss each year, or the equivalent of more than 310 semi truckloads of soil saved.

The IDOA provided grant funding to county SWCD offices in the Illinois River Watershed for operational expenses. Specifically, these funds were used to provide financial support for SWCD offices, programs, and employee' expenses. Employees, in turn, provided technical and educational assistance to both urban and rural residents in the Illinois River Watershed. Their efforts are instrumental in delivering programs that reduce soil erosion and sedimentation that ultimately protects water quality.

In an effort to stabilize and restore severely eroding streambanks that would otherwise contribute a large amount of sediment to the Illinois River and its tributaries, the IDOA, with assistance from SWCDs, administers the Streambank Stabilization and Restoration Program (SSRP). The SSRP is a component of the Partners for Conservation Program that provides funds to construct low-cost techniques to stabilize eroding streambanks. In all, over 4,600 feet of streambanks have been stabilized to protect adjacent water bodies during the past 2 fiscal years.

Illinois Department of Natural Resources

Illinois Recreational Access Program

One of the more challenging problems facing Illinois and the Department of Natural Resources (IDNR) is to provide more public outdoor recreational access and opportunities in Illinois. In order to carry on our outdoor traditions, it is important to connect youth and families to land and opportunities. 95 % of Illinois is privately owned and ranks 46th for public lands for recreation but hosts more than 323,000 hunters and 780,000 fishermen and millions of other recreational users.

Through the Illinois Recreational Access Program (IRAP), the IDNR is increasing public recreational opportunities for the following activities:

- Youth Turkey Hunting
- Fishing (Ponds and Streambanks)
- Non-Motorized Boat Access on Public Waterways
- Outdoor Naturalist (Birding, Nature Watching and Outdoor Photography)

Utilizing resources obtained through a grant from the US Department of Agriculture's Voluntary Public Access and Habitat Incentive Program, the IDNR began leasing private land in November of 2011 from private landowners so that outdoor recreationalists will have more places to go. IRAP is targeting CREP enrollments but it is also available to all eligible farm, ranch, and forested land in the 68 county CREP areas.

In addition to the annual stipend lessees receive, emphasis is placed on developing a conservation management plan for the landowner and assisting with the implementation of the management plan. Resources for habitat protection and enhancement come from IRAP, CREP, EQIP, WHIP, NWTf and other cost-share programs.

- IRAP has leased approximately 13,000 acres in 28 counties within the Illinois and Kaskaskia River Watersheds.
- Made available 320 spring turkey hunting opportunities to youth hunters
- Received 100 youth applications to participate in 2013 spring turkey hunting on IRAP leased sites.
- Completed a stewardship plan and began implementing BMPs in the Honey Creek watershed in Macoupin County involving private landowners, the city of Carlinville, USFWS, NWTf and others partnering together to implement an Illinois Forest Management Plan.
- 40 management plans have been written for IRAP leased properties.

Landowners can enroll their land in any combination of the three turkey seasons: Youth Season, Regular Season 3 and Regular Season 4. If the land isn't enrolled for a particular season, the land will remain open for the landowner to use at their discretion.

Partners for Conservation

Partners for Conservation (formerly Conservation 2000 – or C2000) is a multi-agency, multi-million dollar comprehensive program designed to take a holistic, long-term approach to protecting and managing Illinois' natural resources. The Illinois Department of Natural Resources administers the Ecosystems Program and the Critical Trends Assessment Program (CTAP), a statewide ecosystem assessment and monitoring program.

The Ecosystems Program, a landmark program, is based upon an extensive network of local volunteers working to leverage technical and financial resources to promote ecosystem based management primarily on private lands. With 95% of the state in private ownership (non-state owned), the main objective of the program is to assist in the formation of public/private partnerships, *Ecosystem Partnerships*, to develop plans and projects on a watershed scale with an ecosystem-based approach. There are two key criteria established for the Ecosystems Program. One, that they must be voluntary, and based on incentives rather than government regulation; and, two, they must be broad-based, locally organized efforts, incorporating the interests and participation of local communities, and of private, public and corporate landowners.

Since its inception in 1996, the C2000 Program has awarded more than \$16.4 million in C2000 grants to Ecosystem Partnerships in the **Illinois River watershed** basin for projects providing a variety of conservation practices and outreach. Another \$17.75 million has been leveraged as match for these projects for a total of more than \$34 million for 489 projects. Accomplishments from these projects include: 15,899 acres of habitat restoration, 169,756 feet of stream bank restoration, 1,814 sites have been or are being monitored, and more than 685,745 people have been educated on watershed protection and restoration.

Natural Resources Conservation Service (NRCS)

EQIP

One of NRCS' primary conservation programs is the Environmental Quality Incentives Program (EQIP), which is designed to provide cost-share funds to farmers who qualify for practices designed to improve or create conservation-minded operations or solutions. EQIP addresses practices for livestock operations, grazing operations or non-livestock operations, which covers most of Illinois' private landowners in need of conservation solutions.

EQIP's Forestry Efforts

The primary focus of the Forest Management Plans special project incentive is to help applicants develop management plans and protect their forested acres. Eligible applicants receive funds to help hire a professional forester who will visit the property, inventory the site, and write out a complete woodland management plan. This Special Projects opportunity through Illinois' EQIP can help landowners manage their woodland resources better and obtain a quality management plan that is also approved by the State of Illinois. With more acres of Illinois forest resources well planned for and managed, the health and value of our forest resources will be greatly improved.

Wetland Reserve Program

NRCS' Wetland Reserve Program (WRP) continues to create and restore quality wetland habitats in the Illinois River Watershed and across the state.

For additional information on NRCS conservation programs, please visit www.nrcs.usda.gov.

US Fish and Wildlife Service

Partners for Fish and Wildlife

The US Fish and Wildlife Service (USFWS) Partners for Fish and Wildlife Program (Partners) has supported the Illinois River Conservation Reserve Enhancement Program (CREP) since its inception. The addition of the Kaskaskia River watershed to the CREP program has expanded the opportunities for a collaborative effort to support landscape scale restoration. The Midwest Region's Partners program assists with projects that conserve or restore native vegetation, hydrology and soils associated with imperiled ecosystems such as bottomland hardwoods, native prairies, marshes, rivers and streams. Collaborating with the Illinois and Kaskaskia River CREP has provided opportunities on a landscape scale for restoration, enhancement, and preservation of these natural habitats on private land. Benefits from this collaboration are the enhancements of privately-owned land for Federal Trust Species, such as migratory birds, inter-jurisdictional fish, threatened and endangered species of plants and animals, and other species of conservation concern. Federally listed threatened and endangered species, particularly the threatened decurrent false aster (*Boltonia decurrens*) have benefited from the Illinois CREP. Equally significant are both direct and indirect benefits to National Wildlife Refuge lands located on or near the Illinois and Kaskaskia Rivers' that accrue as a result of expanded habitat adjacent and near the Refuges, as well as improved water quality that results from implementing approved conservation practices.

Partners primary contribution to the Illinois and Kaskaskia River CREP has been technical assistance through participation on the CREP Advisory Committee, providing technical and policy assistance input to the program. At the local level, Partners personnel coordinate with local NRCS, SWCD, and Illinois DNR staff as necessary on individual or groups of projects. CREP has opened a host of opportunities for habitat restoration, enhancement, and preservation on private land that fulfills the objectives of a broad coalition of Federal, State, local, and non-government conservation organizations.

Within the Illinois and Kaskaskia River Watersheds, individual Partners projects compliment CREP and other habitat programs. The Partners program provides a tool for restoration and enhancement of habitats on private lands that may not be eligible for other landowner assistance programs. Partners local coordinators also review the full range of landowner assistance programs with each potential cooperators and refer landowners to CREP and other USDA and Illinois DNR programs that best meet their habitat development and economic goals.

For more information about the Partners for Fish and Wildlife Program please contact: gwen_kolb@fws.gov.

Partners for Fish and Wildlife Program in Southern Illinois: A Summary of Activities and Accomplishments, September 2013 – November 2014; Submitted by Christian D. Greene

The mission of the US Fish and Wildlife Service, Partners for Fish and Wildlife Program is to efficiently achieve voluntary habitat restoration on private lands through financial and technical assistance for the benefit of federal trust species. Although the Partners for Fish and Wildlife Program has not had a presence in Southern Illinois for over a decade, the opportunity to carry out our mission in the region, and at a landscape level, still exists. In a state where the vast majority of land is privately owned, and an area of the state where natural resources are abundant and diverse, Southern Illinois is a logical place to build partnerships with private landowners. The willingness of Illinois landowners to manage their forests to benefit wildlife is illustrated by the large number of participants enrolled in Illinois FDA program. Participating landowners all have approved forest management plans, and a majority wish to follow those recommendations. However, many lack the financial resources and technical know-how to implement those plans. The Partnership between the USFWS, Partners for Fish and Wildlife Program and Illinois Department of Natural Resources has made it possible for many landowners to successfully begin managing their forests.

Activities

In the past 14 months I have been actively building/re-establishing the Partners Program in Southern Illinois, which roughly includes everything south of Interstate 70. I am actively engaged with many Federal Agencies, State Agencies, NGO's, and private individuals including; US Fish and Wildlife, Ecological Services and Refuges, NRCS Region 1, IDNR, IDNR Forestry, US Forest Service, US Army Corps of Engineers, River to River Cooperative Weed Management Area, Ducks Unlimited, Kaskaskia Watershed Association, National Wild Turkey Federation, multiple private forestry consultants, and many others not listed. Working with this wide variety of partners has helped to spread word of the Partners Program in the southern part of the state.

Further, I have worked closely with IDNR District Foresters on several projects. Whether joining me on site visits, recommending the Partners Program to landowners, or through technical expertise they supplied to me personally, we have been able to establish a system where landowners can start to implement their forest plans. Working and communicating with various Federal and State agencies, NGO's, and individuals has allowed each of us to leverage the available resources to best fit the needs of the landowner, while carrying out our respective missions.

Accomplishments

There has been a nearly overwhelming interest from landowners who wish to implement their forest plans. To date, I have conducted over 50 individual site visits with FDA enrolled landowners who were interested in financial and/or technical assistance. Thirteen of those site visits have resulted in Partners contracts, and there are currently 10 or more in some stage of contract development.

The following table is a summary of the 13 Partners Program contracts executed from September 2013 – November 2014.

Treatment	Quantity	Partners Program Cost-Share
Timber Stand Improvement and Exotic Species Control	280.1 ac	\$ 71,966.00
Exotic Species Control Only	506.1 ac	\$ 37,095.00
Streambank Stabilization	905 linear ft	\$ 24,475.00
Tree Plantings	14.2 ac	\$ 7,855.00
Total		\$ 141,391.00

In addition to habitat improvement accomplishments, I have also been able to take advantage of personal training opportunities which include; 1) USFWS Employee Foundations, 2) Heavy Equipment Safety Training, 3) ATV/ORUV training 4) pest control/Certified Public Applicator license, 5) NRCS Area 1 Forestry training, 6) First Aid/CPR, and many federally required online trainings such as FISSA and EEO.

The Future

Threats to oak-hickory forests in Illinois have been identified through several state and federal strategic plans, initiatives, and joint ventures including; 1) Illinois Statewide Forest Plan, 2) Illinois Wildlife Action Plan, 3) National, Midwest Region, and Illinois Partners for Fish and Wildlife Habitat Plans, 4) The Central hardwoods Joint Venture, just to name a few.

The partnership between IDNR and US Fish and Wildlife compliments these strategic plans by directly addressing known threats to oak-hickory forests.

Rarely does the situation present itself where there is a concurrence of identified threats, identified solutions, landowner interest, funding, and a large network of natural resource professionals working towards the same goals. If there ever was a time to make a large and positive impact to Illinois’ natural resources, it is now.

Illinois Farm Bureau

Illinois Farm Bureau (IFB) continues to publicize and promote the Conservation Reserve Enhancement Program (CREP). IFB also used their statewide radio network to highlight details of the program. Information on CREP was sent directly to county Farm Bureaus® (CFB) via e-mail and through county Farm Bureau mail system. Illinois Farm Bureau continues to provide input about CREP through various groups and committees and also continues to voice support for the program. CREP is another tool producers can use that provides cost share incentives and technical assistance for establishing long-term, resource-conserving practices and is a positive program in Illinois.

Association of Illinois Soil and Water Districts (AISWCD)

The AISWCD, in partnership with the Illinois Environmental Protection Agency and the Illinois Department of Natural Resources, helped with administration of the CREP program, by providing funding to SWCDs through a two-year grant funded in part by IEPA 319 and IDNR CREP funds. The grant, which began in June 2012, is a cooperative effort between IEPA, IDNR and the AISWCD.

Through the grant, six positions have been established in strategic workload areas of the Illinois River basin. The six CREP Resource Specialists (CRSs) work with groups of SWCDs within Land Use Councils to monitor existing contracts and work with landowners to enroll additional acres into the Illinois River CREP Area. In addition, the CRSs work with interested landowners to help them enroll acres in the Federal CRP in an effort to increase the acres that will also be eligible for enrollment in CREP. CRSs are also working with landowners to help develop post enrollment management plans for their CREP acres.

The ability to utilize six full-time staff to work exclusively with the CREP program is helping to expedite the enrollment process, increasing the level of monitoring of existing contracts and providing landowners with additional services to benefit their CREP acres and ultimately increase water quality benefits attributable to the Conservation Reserve Enhancement Program.

AISWCD, over the past year, has kept track of CRSs timesheets, expense vouchers, trainings, and insurance. The office administers payment to the Housing Districts quarterly, and issues paychecks and expense voucher checks to the CRSs monthly.

This past year has also seen some turnover in the CRS positions. AISWCD has conducted interviews for positions in Schuyler, LaSalle, and Brown Counties when CRSs have found other employment. Ray Geroff had left in September of 2013. At that time Jake Vancil moved from Schuyler County to Sangamon County leaving the Schuyler County position vacant. January 1, 2014, Dan Sahm filled that position. Russell Blogg, LaSalle County left in May and was replaced by Catherine Dunn in July. Jeremy Pruden's position in Brown County became vacant as of September 19, 2014 and interviews were conducted October 15th to find his replacement.

We thank IDNR and IEPA for their continued support in the CREP program. This program has provided monetary income for both AISWCD and Soil & Water Conservation Districts while also helping to preserve and enhance Illinois' natural resources. All-in-all, this program has provided many benefits and we hope to see it continue into the future.

National Great Rivers Research and Education Center

Providing boots-on-the-ground since 2012, the National Great Rivers Research and Education Center's (NGRREC) *Illinois CREP Initiative* has focused efforts within the newest CREP-eligible watershed—the Kaskaskia River basin. Working in partnership with soil and water conservation districts and the Illinois Department of Natural Resources, Land Conservation Specialists with NGRREC are dedicated to outreach with private landowners about CREP, one-on-one attention with agricultural producers about CREP options and the CREP process, and technical assistance with to complete CREP projects and manage CREP conservation easement parcels.

NGRREC's *Illinois CREP Initiative* is supported by the National Fish and Wildlife Foundation and the Illinois Department of Natural Resources. It adds to other long-term agricultural conservation initiatives at NGRREC, including efforts providing technical assistance to agricultural producers who participate in the Conservation Reserve Program and other USDA conservation programs. Together, agricultural conservation efforts complement NGRREC's research and education missions as they provide high-quality, science-based technical assistance and develop innovative outreach strategies to agricultural producers and private landowners.

The National Great Rivers Research and Education Center is a partnership of Lewis and Clark Community College, the University of Illinois, and the Prairie Research Institute's Illinois Natural History Survey. The Costello Confluence Field Station is located immediately downriver from the Melvin Price Locks and Dam in East Alton, Illinois.

ONGOING INITIATIVES

Monitoring and Assessment of Aquatic Life in the Kaskaskia River for Evaluating IDNR Private Lands Programs – Illinois Natural History Survey

Work during the beginning of the reporting period focused on aggregation and summarization of existing biological and landscape data. We have reviewed and integrated into our database system spatial locations of collections and monitoring data from IDNR/IEPA Intensive Basin Survey Program, IDNR biennial fisheries surveys.

Additionally we have secured the IEPA ambient water quality monitoring for the past 10 years within the Kaskaskia River and its tributaries. Biological survey data, watershed characteristics, and additional information concerning anthropogenic stressors have been assembled and georeferenced for the basin. This includes information on landuse/landcover, surficial geology, modeled flow and water temperature, and point locations from ongoing and historic sampling and monitoring efforts. Locations of CRP/CREP parcels, NPDES permits, stream segments with enhanced dissolved oxygen designation, and stream segments with biologically significant stream designation have also been incorporated into the project GIS data layers. The second half of the reporting period focused on preparation for the field season and beginning the spring and summer sampling programs. Project staff also attended the March meeting of the Kaskaskia Watershed Association in Carlyle to meet with regional constituents and researchers. After coordinating with CREP Mapping Coordinator Lisa Beja on availability of spatial data for CRP and CREP parcels in the study area we attributed local watersheds (1:100,000) throughout the Kaskaskia Basin with summaries of total and local catchment CRP/CREP enrollments to assist with sampling site selection. Private lands practices were classified by their expected efficiency at mitigating sediment and nutrient runoff and local watersheds were classified (high quality, moderate quality, low quality). We developed strata (HUC8, stream size, proportion of CRP/CREP enrolled land) and randomly selected stream reaches for basinwide monitoring that span the range of conditions within the Kaskaskia Basin.

Spring sampling was conducted at 33 sites and included basic habitat, water quality, and biological collections following the protocol established by the Critical Trends Assessment Program (CTAP). The summer sampling program revisited these sites and conducted basic water quality, habitat quality, and biological (electrofishing, rapid macroinvertebrate) sampling. We also sampled 15 sites within stream reaches designated with enhanced dissolved oxygen status and as Biologically Significant Stream segments. Basic water quality data, habitat surveys, and macroinvertebrate collections were made at these focal sites.

A minor budget revision was made to allow for the purchase of a backpack stream shocking unit to be used in fish collections. Having a dedicated stream shocking unit available for the project allowed us to adjust our sampling schedule to coincide with appropriate weather and flow conditions. Total budget and project scope were not changed as a result of this budget adjustment. These efforts were conducted by one full time and one part-time research scientist and several hourly workers. We hired Eric James South to assist Dr. DeWalt with macroinvertebrate sample processing of existing samples and biodiversity assessment of EPT taxa during the Spring sampling period. Eric will begin a graduate program in Entomology in Fall 2013 at University of Illinois Urbana-Champaign and continue to work as part of the project group. We also successfully recruited an additional graduate student to work with Dr. Cao. Levi Drake will join our team in the Fall 2013 Semester pursuing an MS degree at the University of Illinois Urbana-Champaign in the Department of Natural Resources and Environmental Sciences. A total of six summer workers (some part-time) assisted permanent staff with collecting basin-wide and focal reach monitoring data during the summer sampling period.

Please reference Appendix A for the full report.

Establishing a Biological Monitoring Program for CREP to Assess the Conservation Practices and Wildlife Habitat on Property Enrolled – Illinois Natural History Survey

The Illinois Department of Natural Resources (IDNR) is working with the University of Illinois' Critical Trends Assessment Program (CTAP) staff to establish a biological monitoring program for CREP to assess the benefit of conservation practices and wildlife habitat to avian species on property enrolled in CREP. The monitoring program samples the bird communities of shrublands, grasslands, and wetlands at randomly selected CREP easements using point-count based methods. During data collection, the presence and abundance of each species seen or heard during the count period is recorded. Avian point counts are conducted in 190 individual easements within 4 specific state CREP conservation practices, CP23, CP4D, CP22, and CP3A in the Illinois River watershed. Species data will be used to determine CREP easement contribution to regional and state population goals for species of conservation concern. After three years, sampling efforts have detected 103 bird species using CREP easements. Species of conservation concern with frequent detections include Field Sparrow, Dickcissel, Eastern Meadowlark, Northern Bobwhite, Yellow-breasted Chat, Yellow-billed Cuckoo, Brown Thrasher, Willow Flycatcher and Bell's Vireo. CREP easements appear to be providing habitat for many early successional species.

During the 2014 field season this project continued the monitoring effort to assess the reproductive success of shrubland bird species at CREP easements. Nest data are collected at 13 of the randomly selected CREP easements used for point counts. Nest data collected will be used to determine the relative habitat quality and the reproductive contribution of CREP easements to regional and state population goals for species of conservation concern. Focal species include American Robin, Bell's Vireo, Brown Thrasher, Field Sparrow, Grey Catbird, Northern Cardinal, Yellow-billed Cuckoo and Willow Flycatcher.

Please reference Appendix B for the full species list.

Monitoring and Evaluation of Sediment and Nutrient Delivery to the Illinois and Kaskaskia Rivers – Illinois State Water Survey

The Conservation Reserve Enhancement Program (CREP) was initiated as a joint federal/state program with the goal of improving water quality and wildlife habitat in the Illinois River basin. Based on numerous research and long-term data in the Illinois River basin, the two main causes of water quality and habitat degradations in major river corridors were known to be related to sedimentation and nutrient loads. Based on this understanding, the two main objectives of the CREP were to reduce the amount of silt and sediment entering the main stem of the Illinois and Kaskaskia Rivers by 20 percent; and to reduce the amount of phosphorous and nitrogen loadings to by 10 percent. To assess the progress of the program towards meeting the two goals, the Illinois Department of Natural Resources (IDNR) and the Illinois State Water Survey (ISWS) are developing a scientific process for evaluating the effectiveness of the program. The process includes data collection, modeling, and evaluation.

The monitoring and data collection component consist of a watershed monitoring program to monitor sediment and nutrient for selected sub-watersheds within the Illinois and Kaskaskia River basins and also to collect and analyze land use data throughout the river basins. Historically, there are a limited number of sediment and nutrient monitoring stations within those river basins, and most of the available records are of short duration. To fill the data gap and to generate reliable data for small watersheds, the Illinois Department of Natural Resources funded the Illinois State Water Survey to initiate a monitoring program that will collect precipitation, hydrologic, sediment, and nutrient data for selected small watersheds in the Illinois and Kaskaskia River basins that will assist in making a more accurate assessment of sediment and nutrient delivery to the Illinois River. Five small watersheds located within the Spoon and Sangamon River watersheds were selected for intensively monitoring sediment and nutrient within the Illinois River basin. The Spoon River watershed generates the highest sediment per unit area in the Illinois River basin, while the Sangamon River watershed is the largest tributary watershed to the Illinois River and delivers the largest total amount of sediment to the Illinois River. The four small watersheds selected for intensively monitoring sediment and nutrient in the Kaskaskia River basin are located within the Crooked Creek, North Fork Kaskaskia River, Hurricane Creek and Shoal Creek watersheds. In addition, two continuous recording rain gauges were established in the watershed near the monitored watersheds. Lost Creek (402) is a tributary of Crooked Creek which, in turn is a direct tributary of the Kaskaskia River with its confluence

downstream of Carlyle Reservoir. The Carlyle Reservoir is a U.S. Army Corps of Engineers impoundment on the Kaskaskia River. The North Fork Kaskaskia River (403) and Hurricane Creek (404) are direct tributaries of the Kaskaskia River and discharge directly into the upstream end of Carlyle Reservoir. East Fork Shoal Creek (405) is a tributary of Shoal Creek, the largest direct tributary of the Kaskaskia River, with its confluence downstream of Carlyle Reservoir.

After assessing and evaluating many physical, geological, biological, land cover and CRP program data and information, as well as impacts of the 2012 drought, four intensive monitoring stations were selected in the Kaskaskia River Basin and the sediment and nutrient monitoring network was established for the 2014 water year. The year started in one of the coldest winters recorded in the region for some time. This was followed by a particularly wet spring and summer. Nitrogen and phosphorus species concentrations more associated with particulate forms (TKN, t-P) appear to be higher than concentrations of the dissolved forms (NO₃-N and TDP). Further monitoring will determine if this is due to dilution from the unusually wet seasons. Suspended sediment concentrations were higher in watersheds with higher percent area devoted to agriculture production or higher upland slopes. Continued data collection (sediment, nutrient, and precipitation), rating curve development, and compilation of available historical data will be performed in the coming year.

As outlined in the Illinois River Basin Restoration Plan, the alternative of no-action in the Illinois River watershed would have resulted in increased sediment delivery to the Illinois River and habitats and ecosystem would continue to degrade. However, analysis of the available long term data from different sources and the most recent data from the CREP monitoring program, indicate that sediment and nutrient loads from the tributary watersheds are gradually decreasing or stabilizing as a result of implementation of conservation practices in the watershed. With the knowledge that reduction in sediment delivery from large watersheds takes time to move through the system, the indication of stabilized sediment delivery shows progress is being made in restoring the Illinois River watershed. If the present trends continue for the next 10 to 15 years, sediment and nutrient delivery to the Illinois River will be significantly reduced, and lead to improved ecosystem in the river and tributary watersheds in the long-term.

Please reference Appendix C for the Illinois River Report and Appendix D for the Kaskaskia River Report

Habitat Grant Project – National Wild Turkey Federation

National Wild Turkey Federation State Wildlife Grant forester working in the northern Kaskaskia watershed provided support for CREP and EQIP in this region. Accomplishments from October 2013 – December 2013 are summarized below.

- Visited 6 CREP/CRP site visits
- Wrote 3 tree planting plans
- Over 38 acres of tree plantings planned
- Reviewed 92 acres of tree plantings
- Consulted with 5 private landowners
- Wrote 3 Forest Management Plans
- 183 acres in Forest Management Plans
- Reviewed 168 acres of EQIP forest management practices on 7 properties
- Reviewed 28 acres of CREP easements
- Marked 80 acres of timber for harvest or TSI

APPENDIX A



**ILLINOIS NATURAL
HISTORY SURVEY**
PRAIRIE RESEARCH INSTITUTE

Monitoring and Assessment of Aquatic
Life in the Kaskaskia River for evaluating
IDNR Private Lands Programs:
Annual Report 2014

Leon C. Hinz Jr.
and
Brian A. Metzke

Illinois Natural History Survey
Prairie Research Institute
University of Illinois

28 August 2014

INHS Technical Report 2014 (35)

Prepared for: Illinois Department of Natural Resources
Office of Resource Conservation

Unrestricted: for immediate online release.

Prairie Research Institute, University of Illinois at Urbana Champaign
William Shilts, Executive Director

Illinois Natural History Survey
Brian D. Anderson, Director
1816 South Oak Street
Champaign, IL 61820
217-333-6830



Annual Summary Report

Project Title:

Monitoring and Assessment of Aquatic Life in the Kaskaskia River for evaluating IDNR Private Lands Programs.

Project Number: RC13CREP01

Contractor information:

University of Illinois at Urbana/Champaign
Institute of Natural Resource Sustainability
Illinois Natural History Survey
1816 South Oak Street
Champaign, IL 61820

Annual Reporting Period: 1 July 2013—30 June 2014

Annual Project Report Due Date: 28 August 2014

Principle Investigator:

Leon C. Hinz Jr., Ph.D.
Associate Research Scientist in Stream Ecology
Illinois Natural History Survey
One Natural Resources Way,
Springfield, IL 62702-1271
217-785-8297
leon.hinz@illinois.gov

Prepared by: Leon C. Hinz Jr.

Goals/ Objectives: (1) Develop and initiate monitoring program that provides a basin-wide assessment of status and trends for aquatic life in wadeable streams of the Kaskaskia River; (2) track the status of selected populations of sensitive species in focal reaches of the Kaskaskia River associated with enhanced DO regulations, BSS designation, and presence of SGNC; (3) evaluate the influence of conservation easements and associated practices on biological communities within the Kaskaskia River Basin.

Title: Monitoring and Assessment of Aquatic Life in the Kaskaskia River for evaluating IDNR Private Lands Programs

Narrative:

Work during the second full year of the project focused on continuing the basin-wide status and trends sampling program, monitoring within our focal areas, and establishing biological monitoring sites associated with ISWS monitoring locations within the Kaskaskia River Basin.

The summer 2013 sampling program revisited most of the 33 sites sampled during the spring 2013 sampling season and conducted basic water quality, habitat quality, and biological (electrofishing, macroinvertebrate) sampling. We also sampled 15 sites that will serve as focal sites that are contained within stream reaches designated with enhanced dissolved oxygen status and Biologically Significant Stream segments. Basic water quality data, habitat surveys, and macroinvertebrate collections were made at these focal sites.

During late fall (2013), after most crops had been harvested within the basin, we revisited 42 of the spring and summer sites and collected basic water quality measurements. These data will allow us to begin to consider seasonal changes in conditions that may impact aquatic life within the basin.

Project staff scouted 112 additional sites in early spring (2014) to assess conditions for sampling over the remainder of the project period. Thirty-five of these sites were selected for sampling during the upcoming (2014) field seasons along with revisits to 15 focal sites. Spring sampling consisting of macroinvertebrate collections, water quality measurements, and habitat assessment occurred at 33 of these sites (including the focal sites) during the reporting period. Spring macroinvertebrate sample identification has been completed for both the 2013 and 2014 surveys (66 total sample collections). Summer monitoring surveys were ongoing at the end of this reporting period but we expect to have completed a total of 64 basin-wide status and trends sites, 1 CTAP site, 30 sites within our focal areas, and 4 sites associated with ISWS monitoring locations after our first two summer sampling seasons.

Project staff made a presentation to Kaskaskia Watershed Association describing the nature of this work and the type of information that is being collected. Work during this reporting period was primarily conducted by 1 FTE research scientist assisted by the Principle Investigators, associated graduate students, and hourly workers. Two graduate students are currently enrolled and working with staff on project related activities. A total of eight hourly workers (mainly undergraduate students) have also assisted full time staff during the first two years of the project.

Objective 1: Basin-wide status and trends (basin-wide monitoring).

Summer Sampling: We continued to monitor locations during the summer index period throughout the basin. During our site visits we sampled two different biological communities: fish (electric seine or backpack electrofishing unit) and macroinvertebrates (20 jabs proportional to available habitat). Fish were identified and processed at the site and returned to the stream while macroinvertebrate samples were preserved in ethanol and stored for later processing. Basic water quality (water temperature, pH, specific conductance, dissolved oxygen) and nutrient chemistry (N and P using a HACH kit) data were collected. Information for two qualitative habitat indices (QHEI [Ohio EPA 2006] and IHI [Sass et al. 2010]) was also recorded at each site. This work has continued at additional sites during the end of the 2013 and beginning of the 2014 summer sampling periods with an expectation of completing 64 locations by the end of the first two years of field work (Figure 1).

During late fall (2013) basic water quality (water temperature, pH, specific conductance, dissolved oxygen) and nutrient chemistry (N and P using a HACH kit) data were collected at 42 sites that had been visited during the spring and/or summer of 2013 (Figure 2). These included status and trends (Objective 1) and focal reach monitoring sites (Objective 2). Water temperature data were also recovered from electronic monitors deployed at 35 sites during the first summer sampling period (2013) and these data have been summarized. Temperature monitors were redeployed during the late spring 2014 at additional locations that were scheduled to be sampled in 2014.

Habitat data from the first full year of field work were scored by calculating the QHEI [Ohio EPA 2006] and the IHI [Sass et al. 2010] for all reaches where appropriate information was available from the summer samples. Station based gradient was determined using the approach associated with the Illinois Fish IBI for all project sites so that fish metrics could be scored using this index. Fish IBI scores were calculated for all samples collected during the first field season (2013). This work is ongoing for samples collected during the second field season (2014).

During the spring of 2014 project staff visited and evaluated 112 sites on the Kaskaskia River for possible sampling over the next two years of the project. Seventy-six sites were determined to be acceptable, seven possible, and the remainder not sample-able (usually due to access issues or being too large for our gear). Thirty-five of these sites were selected to be sampled during the Spring and Summer 2014 field seasons.

Spring Sampling: We collected macroinvertebrate, habitat quality, and basic water quality information at an additional 33 sampling locations during the late Spring of 2014. For each acceptable site, a dipnet was used to collect macroinvertebrates from two high energy microhabitats and two low energy microhabitats. Specimens were individually picked from debris examined in collecting trays and subsequently stored in 95% ETOH. Water temperature, dissolved oxygen, percent saturation of dissolved oxygen, conductivity and pH were recorded using a Quanta hydrolab. Observations were recorded

for general land use, erosion, stream morphology, sediment characteristics, water surface oils, weather conditions and collected invertebrate taxa. Habitat assessment was recorded for 12 parameters suggested by the Critical Trends and Assessment Program. This work will continue in subsequent years of the project at additional sites.

Objective 2: Status of sensitive species (focal reach monitoring).

Focal reach monitoring targeted locations within designated Biologically Significant Stream Segments (BSS; Bol et al. 2007, State of Illinois 2008) and designated enhanced dissolved oxygen reaches (IDNR/IEPA 2006). For the second year we characterized physical and chemical conditions at 15 sites within focal reaches by collecting basic water quality data, conducting a habitat survey, and placing a series of water temperature monitors in enhanced DO reaches that overlap with BSS segments (Figure 3, focal sites). Macroinvertebrate samples were also collected at these sites using the 20 jab method proportional to available habitat.

Focal sites were visited during the spring and sampled using the CTAP streams protocols. Macroinvertebrates (EPT) were collected and a habitat assessment and water quality measurements were taken at this time. These sites were also revisited during the summer sampling program (Figure 3, focal sites). These efforts will continue in subsequent years of the project.

Habitat Data: We again characterized the physical and chemical habitat of 15 sites within focal reaches by collecting basic water quality data, conducting a habitat survey, and placing a series of water temperature monitors (Figure 3, focal sites). Macroinvertebrate samples were collected at these sites using the 20 jab method proportional to available habitat. These efforts will continue in subsequent years of the project at these sites.

Fish Community Data: We collected a total of twelve fish samples in consecutive reaches along six stream segments within the Kaskaskia basin (Figure 3, intense sites). These samples will be used to evaluate the efficiency of our collection methods, track species fidelity to local habitat conditions, and improve our ability to assess the distribution and abundance of rare species in the basin. These efforts will continue in subsequent years of the project at additional sites.

Objective 3: Influence of private land conservation efforts (fixed site monitoring).

Project staff provided comments to Illinois State Water Survey staff (Laura Keefer) on site selection for discharge, sediment, and nutrient monitoring. Our biological sampling efforts will be coordinated to take advantage of these data intensive discharge, sediment, and nutrient monitoring data.

Four sites selected by the Illinois State Water Survey (Laura Keefer) were scouted during the spring of 2014. All four locations were included in our sampling during the 2014 Spring and Summer field seasons (Figure 3, ISWS sites). This work is ongoing.

Reporting:

Project staff made a presentation to Kaskaskia Watershed Association describing the nature of this work and the type of information that is being collected during this reporting period. Details of this project's objectives, scope, and progress to date were presented to Illinois Department of Natural Resources staff involved in CREP and other private lands programs. The purpose of the presentation was to introduce new IDNR staff to ongoing monitoring and assessment efforts. Quarterly reports and this annual project report were prepared and submitted.

Literature Cited:

- Bol, L., A.M. Holtrop, L.C. Hinz Jr., and J. Epifanio. 2007. Evaluating Streams in Illinois based on Aquatic Biodiversity. INHS Technical Report 2007(57).
- IDNR/IEPA 2006. Recommended Revisions to the Illinois General Use Water-Quality Standard for Dissolved Oxygen. Illinois Department of Natural Resources and Illinois Environmental Protection Agency, Springfield, IL.
- Ohio EPA 2006. Methods for Assessing Habitat in Flowing Waters: Using the Qualitative Habitat Evaluation Index (QHEI). Ohio EPA Technical Bulletin EAS/ 2006-06-1. State of Ohio Environmental Protection Agency, Division of Surface Water. June 2006.
- Sass L. L., L.C. Hinz Jr., J. Epifanio and A. M. Holtrop. 2010. Developing a multi-metric habitat index for wadeable streams in Illinois. Final Report to the Illinois Department of Natural Resources. Illinois Natural History Survey Technical Report 2010/21.
- State of Illinois, 2008. Integrating Multiple Taxa in a Biological Stream Rating System. Illinois Department of Natural Resources, Springfield, IL. 34 pp.

CREP Survey Locations Through 2014




Figure 1. Monitoring locations sampled for evaluating the condition of the Kaskaskia River basin associated with IDNR Private Lands Programs during the 2013 and 2014 Field Seasons.

Fall 2013 Water Quality Sampling and Temperature Recorder Retrieval Results




Figure 2. Kaskaskia River water quality sample collection locations (Fall 2013).

CREP Survey Locations Through 2014




Figure 3. Types of monitoring locations sampled for evaluating the condition of the Kaskaskia River basin associated with IDNR Private Lands Programs. Random sites relate to Objective 1; Focal sites relate to Objective 2; CTAP, intense, and ISWS sites relate to Objective 3.

APPENDIX B

Illinois CREP Assessment 2014

This is a brief summary of the avian research conducted during the spring and summer of 2014 at randomly selected locations within CREP easements.

During the second year of our assessment of the CREP program in Illinois we continued point counts at all locations sampled in 2012 and 2013 continued nest searching for shrubland species at 13 easements. Easements were located in Brown, Christian, Fulton, Hancock, Knox, Logan, Menard, McDonough, and Sangamon counties. Conservation practices sampled included CP4D, CP3A, CP23, and CP22. On the ground habitat types were forest (7%), grassland (44%), and shrubland (49%). Average easement size was 11.4 ha and the range was 2.97–78.5 ha. We conducted 573-point counts at 190 CREP easements. Bird detections were similar in species composition to our 2012 and 2013 surveys (results below). During nest searches in 2014 we found 167 Bell's Vireo, 87 Willow Flycatcher, 49 Brown Thrasher, 45 American Robin, 56 Gray Catbird, 64 Field Sparrow, and 55 Northern Cardinal active nests.

Species list in order of most detected to least detected (with more than 50 detections):

Red-Winged Blackbird
Common Yellowthroat
Indigo Bunting
Field Sparrow*
American Goldfinch
Northern Cardinal
Dickcissel*
American Robin
Song Sparrow
Gray Catbird
Mourning Dove
Red-Bellied Woodpecker
American Crow
Rose-Breasted Grosbeak
Yellow-Billed Cuckoo*
Yellow Warbler
Blue Jay
Warbling Vireo
Willow Flycatcher
Yellow- Breasted Chat
Brown-Headed Cowbird
Eastern Towhee
Tufted Titmouse
Eastern Wood Pewee
Northern Bobwhite*
Great-Crested Flycatcher
Eastern Meadowlark
Bell's Vireo*

Ring-Necked Pheasant
Turkey Vulture
House Wren
Red Eyed Vireo
Baltimore Oriole
Northern Flicker
Brown Thrasher
Downy Woodpecker
Barn Swallow
Cedar Waxwing
Eastern Kingbird

***Listed as Species in Greatest Need of Conservation for Illinois**

Other species in Greatest Need of Conservation for Illinois we detected in small numbers were: Bobolink (1), Red-Headed Woodpecker (12), Henslow's sparrow (11), Grasshopper Sparrow (13), Sedge Wren (4), Black-billed Cuckoo (19), Blue Grosbeak (18).

Vegetation:

Species encountered at CREP sites reveal that most of the plants are native, though these species are disturbance tolerant and considered weedy. Native annual weeds like common and giant ragweed, tall boneset, and annual fleabane were encountered at many sites. Common goldenrod, found at every site, is a quick growing; native perennial herb that readily colonizes disturbed soil. Other weedy native, perennials included panicked aster and hairy aster. Woody natives with a somewhat weedy habit included silver maple, eastern cottonwood, and green ash (*Fraxinus pennsylvanica*).

Native plant species were generally more abundant than non-native species, but invasive species like reed canary grass, field thistle, and Amur honeysuckle were present on some sites. Compared to randomly selected wetland and grassland sites sampled as part of the Critical Trends Assessment Program (CTAP), the CREP sites were more botanically rich and diverse, but as sites mature without management or disturbance, plant diversity is expected to decline.

APPENDIX C

Monitoring and Evaluation of Sediment and Nutrient Delivery to the Illinois River: Illinois River Conservation Reserve Enhancement Program (CREP)

by
Illinois State Water Survey
Illinois Department of Natural Resources

Prepared for the
Office of Resource Conservation,
Illinois Department of Natural Resources

November 2014

Contents

	Page
1. Introduction.....	1
Acknowledgments.....	1
2. Monitoring and Data Collection	3
Sediment and Nutrient Data.....	3
Sediment Data	6
Nutrient Data	6
Sediment and Nutrient Loads.....	12
Sediment and Nutrient Yields	22
Additional CREP Data collection Efforts.....	25
3. Land Use Cover and Conservation Practices.....	29
Historical Agricultural Land Use Trends in Illinois River Basin.....	29
Land Cover.....	33
Conservation Practices.....	39
4. Variability and Trends in Precipitation and Streamflow	43
5. Model Development and Applications.....	53
HSPF Model.....	53
Model Input Data.....	55
Model Development	63
Modeling Results.....	63
6. Analyses and Discussion	69
Sediment Loadings	69
Nutrient Loadings.....	71
7. Summary and Conclusions	79
8. References.....	81
Appendix A. Quality Assurance Project Plan (QAPP)	A-1
Appendix B. Streamflow Data.....	B-1
Appendix C. Suspended Sediment Data	C-1
Appendix D. Nutrient Data (Nitrogen)	D-1
Appendix E. Nutrient Data (Phosphorous).....	E-1

List of Tables

		Page
2-1	Sediment and Nutrient Monitoring Stations Established for the Illinois River CREP.....	3
2-2	Summary Statistics for Water Years 2000–2007. All concentrations in mg/L.....	9
2-3	Summary of Annual Water Discharges, Sediment and Nutrient Loads at Court Creek Monitoring Station (301).....	11
2-4	Summary of Annual Water Discharges, Sediment and Nutrient Loads at North Creek Monitoring Station (302)	12
2-5	Summary of Annual Water Discharges, Sediment and Nutrient Loads at Haw Creek Monitoring Station (303)	12
2-6	Summary of Annual Water Discharges, Sediment and Nutrient Loads at Panther Creek Monitoring Station (201).....	13
2-7	Summary of Annual Water Discharges, Sediment and Nutrient Loads at Cox Creek Monitoring Station (202)	13
2-8	Sediment Yield in tons/acre for the CREP Monitoring Stations.....	22
2-9	Nitrate-N Yield in lbs/acre for the CREP Monitoring Stations	23
2-10	Total Phosphorous Yield in lbs/acre for the CREP Monitoring Stations	23
2-11	Additional CREP Monitoring Stations in the Spoon River Watershed.....	26
2-12	Suspended Sediment Concentration Data (mg/l) for Swan and Cedar Creeks	27
2-13	Suspended Sediment Concentration Data (mg/L) for London Mills and Seville	28
3-1	7-year average (2007-2013) percent acres of land cover area by watershed.....	35
4-1	Kendall Tau-b Trend Statistics for Flow Records on the Illinois River and Major Tributaries	51
4-2	Average Annual Precipitation and Streamflow (inches) for Different Periods of Record.....	51

List of Figures

		Page
2-1	Locations of available in-stream sediment data within the Illinois River watershed, 1981-2000	4
2-2	Location of monitoring stations in Court and Haw Creek watersheds.....	5
2-3	Location of monitoring stations in Panther and Cox Creek watersheds.....	5
2-4	Suspended sediment concentrations and water discharge at Court Creek (301) for Water Years 2000 and 2001	7
2-5	Concentrations of nitrogen species and water discharge at Court Creek (301) for Water Years 2002 and 2003	8
2-6	Concentrations of phosphorous species and water discharge at Court Creek (301) for Water Years 2002 and 2003	10
2-7	Annual runoff at the five CREP monitoring stations	14
2-8	Annual suspended sediment loads at the five CREP monitoring stations	15
2-9	Annual nitrate-N loads at the five CREP monitoring stations	16
2-10	Annual ammonium-N loads at the five CREP monitoring stations	17
2-11	Annual Kjeldahl nitrogen loads at the five CREP monitoring stations	18
2-12	Annual phosphorous loads at the five CREP monitoring stations	19
2-13	Annual dissolved phosphorous loads at the five CREP monitoring stations.....	20
2-14	Annual ortho-phosphate phosphorous loads at the five CREP monitoring stations	21
2-15	Average annual sediment yield in tons/acre for the CREP monitoring stations	24
2-16	Average annual nitrate-N yield in lbs/acre for the CREP monitoring stations.....	24
2-17	Average annual total phosphorous yield in lbs/acre for the CREP monitoring stations	25
2-18	Locations of monitoring stations in the Cedar and Swan watersheds	26
3-1	Land cover of the Illinois River basin (NASS, 2010)	30
3-2	Acreage of agricultural land uses in State of Illinois (1866-2006)	31
3-3	Acreage of agricultural land uses in Illinois River basin (1925-2006).....	32
3-4	Acreage of agricultural land uses in Spoon River watershed (1925-2006)	33
3-5	Acreage of agricultural land uses in Sangamon River watershed (1925-2006).....	35
3-6	Watersheds being monitored for hydrology, sediment and nutrients.....	35
3-7	Illinois River Basin Watersheds: Percent Change in Generalized NASS Land-Use from 2007-2013	36

List of Figures (continued)

		Page
3-8	Panther Creek Watershed from ISWS Station 201: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013	36
3-9	Cox Creek Watershed from ISWS Station 202: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013	37
3-10	Court Creek Watershed from ISWS Station 301: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013	37
3-11	North Creek Watershed from ISWS Station 302: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013	38
3-12	Haw Creek Watershed from ISWS Station 303: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013	38
3-13	State and CREP contract locations	40
3-14	Cumulative acres of conservation practices installed in Court Creek watershed at monitoring station ISWS #201 from 1999-2013	41
3-15	Cumulative acres of conservation practices installed in Haw Creek watershed at monitoring station ISWS #202 from 1999-2013	41
4-1	Location of streamgaging stations with long-term data used in the analysis of variability and trends	44
4-2	Ten-year average precipitation and streamflow, Illinois River at Peoria-Kingston Mines	45
4-3	Ten-year average precipitation and streamflow, Fox River at Dayton.....	45
4-4	Ten-year average precipitation and streamflow, Kankakee River at Momence	46
4-5	Ten-year average precipitation and streamflow, Spoon River at Seville	46
4-6	Ten-year average precipitation and streamflow, Sangamon River at Monticello	47
4-7	Ten-year average precipitation and streamflow, LaMoine River at Ripley.....	47
4-8	Ten-year average precipitation and streamflow, Macoupin Creek near Kane.....	48
4-9	Ten-year average precipitation and streamflow, Illinois River at Valley City	48
4-10	Locations of long-term streamflow gages (at least 89 years of record) showing statistically significant trends in mean annual flow in the eastern United States (from Knapp, 2005)	50
5-1	Location of the Spoon River watershed.....	54
5-2	Schematic of the subwatershed and stream delineation, and precipitation gages used for the Haw Creek model.....	56
5-3	Schematic of the subwatershed and stream delineation, and precipitation gages used for the Haw Creek model.....	57

List of Figures (continued)

		Page
5-4	Schematic of the subwatershed and stream delineation, and precipitation gages used for the Spoon River watershed model	58
5-5	Land use in the Court Creek watershed	59
5-6	Land use in the Haw Creek watershed	60
5-7	Land use in the Spoon River watershed.....	61
5-8	Soil types in the Spoon River watershed	62
5-9	Results of model calibration for streamflow simulation for the Court Creek watershed	66
5-10	Preliminary results of model calibration for suspended sediment concentration simulation for the Court Creek watershed	66
5-11	Comparison of observed and simulated streamflow by the Haw Creek watershed model developed using the calibrated parameters from the Court Creek watershed model.....	67
5-12	Preliminary results for suspended sediment concentration from the Haw Creek watershed model developed using the calibrated parameters from the Court Creek watershed model.....	67
5-13	Comparison of observed and simulated streamflow simulation by the Spoon River watershed model developed using the calibrated parameters from the Court Creek watershed model.....	68
5-14	Preliminary results for suspended sediment concentration from the Spoon River watershed model developed using the calibrated parameters from the Court Creek watershed model.....	68
6-1	Variability of sediment yield per inch of runoff for CREP monitoring stations.....	70
6-2	Comparison of sediment load from CREP monitoring stations with historical sediment data for small watersheds by the USGS.....	70
6-3	Sediment delivery from the three major tributary watersheds to the Illinois River and sediment outflow from the Illinois River at Valley City.....	72
6-4	Trends in sediment load at Spoon River at London Mills (after Crowder et al., 2008).....	73
6-5	Trends in sediment load at LaMoine River at Ripley, IL (after Crowder et al., 2008).....	73
6-6	Trends in sediment load at Sangamon River at Monticello, IL (after Crowder et al., 2008).....	74

List of Figures (concluded)

	Page
6-7	Variability of nitrate-N yield per inch of runoff for CREP monitoring stations..... 74
6-8	Variability of total phosphorous yield per inch of runoff for CREP monitoring stations 75
6-9	Annual nitrate-N loads for the three major tributary watersheds to the Lower Illinois River 76
6-10	Annual total phosphorous loads for the three major tributary watersheds to the Lower Illinois River 76
6-11	Nitrate-N and total phosphorous loads along the Lower Illinois River 77

Monitoring and Evaluation of Sediment and Nutrient Delivery to the Illinois River: Illinois River Conservation Reserve Enhancement Program (CREP)

by
Illinois State Water Survey
Illinois Department of Natural Resources

1. Introduction

The Illinois River Conservation Reserve Enhancement Program (CREP) was initiated as a joint federal/state program with the goal of improving water quality and wildlife habitat in the Illinois River basin. Based on numerous research and long-term data, the two main causes of water quality and habitat degradations in the Illinois River were known to be related to sedimentation and nutrient loads. Based on this understanding, the two main objectives of the Illinois River CREP were stated as follows:

- 1) Reduce the amount of silt and sediment entering the main stem of the Illinois River by 20 percent.
- 2) Reduce the amount of phosphorous and nitrogen loadings to the Illinois River by 10 percent.

To assess the progress of the program towards meeting the two goals, the Illinois Department of Natural Resources (IDNR) and the Illinois State Water Survey (ISWS) are developing a scientific process for evaluating the effectiveness of the program. The process includes data collection, modeling, and evaluation. Progress made so far in each of these efforts is presented in this report.

Acknowledgments

The work upon which this report is based was supported by funds provided by the Office of Resource Conservation, Illinois Department of Natural Resources. Ms. Debbie Bruce and Richard Mollahan managed the project for IDNR and provided the proper guidance and support to design and operate the monitoring program and the associated research. Their continued support and guidance is greatly appreciated.

Several Illinois State Water Survey staff participated and contributed towards the successful accomplishment of project objectives. Jim Slowikowski and Kip Stevenson are responsible for the data collection and analysis. Laura Keefer was responsible for analysis of the

land use data. Jas Singh and Yanqing Lian were responsible for the development of the watershed models. Vern Knapp provided the analyses on variability and trends in precipitation and streamflow in the Illinois River basin. Momcilo Markus analyzed the Illinois Environmental Protection Agency nutrient data for analyses of long-term trends. Sangeetha Chandrasekaran analyzed the Benchmark Sediment Monitoring data for long-term trend analysis. Patti Hill prepared the draft and final reports.

2. Monitoring and Data Collection

The monitoring and data collection component consist of a watershed monitoring program to monitor sediment and nutrient for selected watersheds within the Illinois River basin and also to collect and analyze land use data throughout the river basin. Historically, there are a limited number of sediment and nutrient monitoring stations within the Illinois River basin, and most of the available records are of short duration. For example, figure 2-1 shows all the active and inactive sediment monitoring stations within the Illinois River basin prior to the start of monitoring for CREP. Out of the 44 stations shown in the map, only 18 stations had records longer than 5 years and only 8 stations had more than 10 years of record. Therefore the available data and monitoring network was insufficient to monitor long-term trends especially in small watersheds where changes can be observed and quantified more easily than in larger watersheds.

To fill the data gap and to generate reliable data for small watersheds, the Illinois Department of Natural Resources funded the Illinois State Water Survey to initiate a monitoring program that will collect precipitation, hydrologic, sediment, and nutrient data for selected small watersheds in the Illinois River basin that will assist in making a more accurate assessment of sediment and nutrient delivery to the Illinois River.

Sediment and Nutrient Data

Five small watersheds located within the Spoon and Sangamon River watersheds were selected for intensively monitoring sediment and nutrient within the Illinois River basin. The locations of the watersheds and the monitoring stations are shown in figures 2-2 and 2-3 and information about the monitoring stations is provided in table 2-1. Court and North Creeks are located within the Spoon River watershed, while Panther and Cox Creeks are located within the Sangamon River watershed. The Spoon River watershed generates the highest sediment per unit area in the Illinois River basin, while the Sangamon River watershed is the largest tributary watershed to the Illinois River and delivers the largest total amount of sediment to the Illinois River. The type of data collected and the data collection methods have been presented in detail in the first progress report for the monitoring program (Demissie et al., 2001) and in the Quality Assurance Project Plan (QAPP) given in Appendix A. This report presents the data that have been collected and analyzed at each of the monitoring stations.

Table 2-1. Sediment and Nutrient Monitoring Stations Established for the Illinois River CREP

<i>Station ID</i>	<i>Name</i>	<i>Drainage area</i>	<i>Watershed</i>
301	Court Creek	66.4 sq mi (172 sq km)	Spoon River
302	North Creek	26.0 sq mi (67.4 sq km)	Spoon River
303	Haw Creek	55.2 sq mi (143 sq km)	Spoon River
201	Panther Creek	16.5 sq mi (42.7 sq km)	Sangamon River
202	Cox Creek	12.0 sq mi (31.1 sq km)	Sangamon River




Figure 2-2. Location of monitoring stations in Court and Haw Creek watersheds




Figure 2-3. Location of monitoring stations in Panther and Cox Creek watersheds

Sediment Data

The daily streamflow and suspended sediment concentrations observed at all the five monitoring stations from Water Year 2000 to Water Year 2013 are given in Appendix B and C. Examples of the frequency of data collection are shown in figures 2-4 and 2-5 for the Court Creek Station. A summary of statistics for all stations showing the mean, median, minimum maximum, 25th percentile, and 75th percentile are given in table 2-2. Over 27,691 samples have been collected and analyzed at the five monitoring stations since the monitoring program was initiated. As can be seen in the figures, suspended sediment concentrations are highly variable throughout a year and also from year to year depending on the climatic conditions. It is also evident that sediment concentrations are the highest during storm events resulting in the transport of most of the sediment during storm events. Therefore, it is extremely important that samples are collected frequently during storm events to accurately measure sediment loads at monitoring stations.

Nutrient Data

All the nutrient data collected and analyzed from Water Year 2000 through Water Year 2013 at the five monitoring stations are given in Appendices D and E. The nutrient data are organized into two groups: nitrogen species and phosphorous species. The nitrogen species include nitrate-nitrogen (NO₃-N), nitrite-nitrogen (NO₂-N), ammonium-nitrogen (NH₄-N), and total Kjeldahl nitrogen (TKN). The phosphorous species include total phosphorous (TP), total dissolved phosphorous (TDP), and orthophosphate (P-ortho). Over 4,990 samples have been collected and analyzed for nitrate (NO₃-N), ammonium (NH₄-N) and orthophosphate (P-ortho). In addition, more than 2,723 samples have been analyzed for nitrate (NO₂-N), total Kjeldahl nitrogen (TKN), total phosphorous (TP), and total dissolved phosphorous (TDP). Examples of the type of data collected for the nitrogen species are shown in figure 2-5, while those for the phosphorous species are shown in figure 2-6. A summary statistics for all stations showing the mean, median, minimum, maximum, 25th percentile, and 75th percentile are given in table 2-2.

Data for the nitrogen species at all five monitoring stations show that the dominant form of nitrogen transported by the streams is nitrate-N. During storm events, the concentration of TKN rises significantly, sometimes exceeding the nitrate-N concentration. TKN is highly correlated to suspended sediment concentrations.

One significant observation that can be made from the data is the consistently higher concentrations of nitrate-N at Panther Creek and Cox Creek (tributaries to the Sangamon River) than at Court Creek, North Creek, and Haw Creek (tributaries of the Spoon River).

Data for the phosphorous species at all five monitoring stations show that most of the phosphorous load is transported during storm events. Concentrations of total phosphorous are the highest during storm events and relatively low most of the time. This is very similar to that shown by sediment and thus implies high correlations between sediment and phosphorous concentrations and loads. In terms of phosphorous concentrations, it does not appear there is any significant difference between the different monitoring stations from the Spoon and Sangamon River watersheds.





Figure 2-4. Suspended sediment concentrations and water discharge at Court Creek (301) for Water Years 2000 and 2001





Figure 2-5. Concentrations of nitrogen species and water discharge at Court Creek (301) for Water Years 2002 and 2003

Table 2-2. Summary Statistics for Water Years 2000–2013. All concentrations in mg/L

	<i>NO3-N</i>	<i>oPO4-P</i>	<i>NH4-N</i>	<i>NO2-N</i>	<i>TKN</i>	<i>t-P</i>	<i>t-P-Dissolved</i>	<i>SSC</i>
Panther Creek (Station 201)								
Count	915	915	915	450	449	449	449	5526
Mean	4.06	0.12	0.10	0.03	2.37	1.01	0.17	860.8
Median	3.35	0.08	0.06	0.02	0.97	0.33	0.12	127.9
Min	< 0.04	< 0.01	< 0.03	< 0.01	< 0.12	< 0.03	< 0.03	1.47
Max	14.76	1.31	5.99	0.21	23.99	11.21	1.38	48289.0
25th Percentile	0.21	0.05	< 0.04	0.01	0.45	0.12	0.08	53.8
75th Percentile	6.83	0.14	< 0.08	0.04	3.06	1.28	0.20	434.6
Cox Creek (Station 202)								
Count	921	921	921	452	452	452	452	4813
Mean	5.82	0.20	0.71	0.05	3.61	1.13	0.30	684.1
Median	5.67	0.10	0.07	0.04	1.44	0.43	0.18	149.6
Min	< 0.04	< 0.01	< 0.03	< 0.01	< 0.14	< 0.04	< 0.03	0.95
Max	19.83	7.81	300.33	1.26	390.37	29.10	8.21	22066.5
25th Percentile	0.92	0.06	< 0.06	0.02	0.57	0.16	0.09	71.1
75th Percentile	9.40	0.22	0.20	0.06	3.44	1.24	0.36	391.3
Court Creek (Station 301)								
Count	1055	1055	1055	613	612	612	612	5157
Mean	3.11	0.07	0.14	0.04	2.53	0.86	0.11	655.7
Median	3.05	0.05	0.07	0.03	1.35	0.37	0.09	116.2
Min	< 0.04	< 0.003	< 0.03	< 0.01	0.23	0.03	< 0.03	1.93
Max	11.37	0.69	0.90	0.13	18.69	6.58	0.71	13632.0
25th Percentile	0.98	0.03	< 0.06	0.02	0.65	0.12	0.05	50.8
75th Percentile	4.87	0.09	0.17	0.05	3.45	1.21	0.13	559.6
North Creek (Station 302)								
Count	1044	1044	1044	601	601	601	601	6191
Mean	3.14	0.07	0.14	0.04	2.33	0.80	0.12	492.0
Median	3.07	0.04	0.07	0.03	1.16	0.31	0.09	95.6
Min	< 0.04	< 0.003	< 0.03	< 0.01	0.23	< 0.04	< 0.03	0.36
Max	12.66	1.05	1.55	0.19	17.95	6.69	1.07	15137.1
25th Percentile	0.78	0.02	< 0.06	< 0.02	0.62	0.11	0.05	39.9
75th Percentile	4.94	0.09	0.16	0.05	2.62	0.95	0.14	285.6
Haw Creek (Station 303)								
Count	1055	1055	1055	609	609	609	609	6004
Mean	4.55	0.08	0.12	0.05	2.34	0.78	0.11	578.9
Median	4.62	0.06	0.07	0.04	1.37	0.41	0.09	164.1
Min	< 0.04	< 0.003	< 0.03	< 0.01	0.23	0.04	< 0.03	2.17
Max	12.59	0.71	1.07	0.21	16.75	5.92	0.95	9878.8
25th Percentile	1.99	0.03	< 0.06	0.03	0.64	0.14	0.06	54.8
75th Percentile	6.81	0.09	0.14	0.06	3.04	1.06	0.13	629.1





Figure 2-6. Concentrations of phosphorous species and water discharge at Court Creek (301) for Water Years 2002 and 2003

Sediment and Nutrient Loads

The sediment and nutrient concentrations and water discharges are used to compute the amount of sediment and nutrient transported past monitoring stations. Based on the available flow and concentration data, daily loads are computed for sediment and the different species of nitrogen and phosphorous. The daily loads are then compiled to compute monthly and annual loads. Results of those calculations are summarized in tables 2-3 to 2-7 for each of the five monitoring stations. Each table presents the annual water discharge, sediment load, nitrate-N load, and the total phosphorous load for one of the stations. Similar calculations have been made for the other species of nitrogen and phosphorous, but are not included in the summary tables. The annual sediment loads are highly correlated to the water discharge, and thus the wetter years, 2001, 2002, 2005, 2007, 2008, 2009, 2010, 2011 and 2013 generated more sediment at all stations as compared to drier years, 2000, 2003, 2004, 2006, and 2012. The annual sediment loads ranged from a low of 105 tons in WY2012 at Panther Creek to a high of 174,742 tons in 2009 at Court Creek. The nitrate-N loads ranged from a low of 1.8 tons in 2012 at Cox Creek to a high of 585 tons in WY2010 at Haw Creek. The total phosphorous loads ranged from a low of 0.2 tons in 2012 at Cox Creek and Panther Creek to a high of 117.6 tons in 2010 at Court Creek. For comparison purposes, the runoff, sediment, nitrate-N, nitrite-N, ammonium-N, Kjeldahl-N, total phosphorous, total dissolved phosphorous, and total ortho-phosphate phosphorous loads (for the five monitoring stations) are shown in figures 2-8 to 2-15. In terms of the total annual loads, the larger watersheds, Court and Haw, consistently carry higher sediment and nutrient loads than Panther and Cox Creeks. However, per unit area Panther and Cox generate more sediment than Court, North, and Haw Creeks.

Table 2-3. Summary of Annual Water Discharges, Sediment and Nutrient Loads at Court Creek Monitoring Station (301)

<i>Water Year</i>	<i>Water discharge (cfs)</i>	<i>Load</i>		
		<i>Sediment (tons)</i>	<i>Nitrate-N (tons)</i>	<i>Total phosphorus (tons)</i>
2000	11880	26527	131.2	35.0
2001	22100	43633	274.8	39.2
2002	17320	62898	203.7	47.9
2003	6805	21749	59.9	18.3
2004	7459	7359	76.0	7.5
2005	14400	18831	207.5	20.4
2006	5650	7897	84.3	6.5
2007	19376	48974	240.8	46.8
2008	22442	41077	265.4	45.6
2009	41207	174742	429.6	116.9
2010	44836	146202	425.9	117.6
2011	23311	55337	270.9	43.3
2012	6129	4145	36.7	4.8
2013	26158	116616	270.8	94.9

Table 2-4. Summary of Annual Water Discharges, Sediment and Nutrient Loads at North Creek Monitoring Station (302)

<i>Water Year</i>	<i>Water discharge (cfs)</i>	<i>Load</i>		
		<i>Sediment (tons)</i>	<i>Nitrate-N (tons)</i>	<i>Total phosphorus (tons)</i>
2000	4009	6969	42.8	10.4
2001	8091	16747	102.9	12.7
2002	7372	29269	97.8	24.2
2003	3039	11422	32.9	9.1
2004	3224	2038	37.7	2.4
2005	5266	6061	76.3	7.7
2006	2151	4179	36.2	3.4
2007	7524	16702	99.3	14.3
2008	9416	19762	119.0	21.0
2009	16544	62806	167.9	45.2
2010	18577	66501	167.4	52.7
2011	9491	25979	105.4	25.2
2012	2506	2207	14.9	2.0
2013	12624	60934	121.1	44.9

Table 2-5. Summary of Annual Water Discharges, Sediment and Nutrient Loads at Haw Creek Monitoring Station (303)

<i>Water Year</i>	<i>Water discharge (cfs)</i>	<i>Load</i>		
		<i>Sediment (tons)</i>	<i>Nitrate-N (tons)</i>	<i>Total phosphorus (tons)</i>
2000	11433	21283	162.2	32.0
2001	19878	49580	322.0	58.0
2002	15603	44221	256.5	42.8
2003	4337	5908	41.7	8.3
2004	8676	10914	143.4	12.6
2005	14661	18047	281.4	18.5
2006	5341	5770	113.7	6.0
2007	15032	20127	262.5	23.9
2008	14054	16396	227.0	25.5
2009	34003	104081	506.4	85.9
2010	40230	92974	585.2	85.4
2011	20788	37379	372.5	34.3
2012	5326	2185	55.1	3.3
2013	23581	75175	357.8	74.1

Table 2-6. Summary of Annual Water Discharges, Sediment and Nutrient Loads at Panther Creek Monitoring Station (201)

<i>Water Year</i>	<i>Water discharge (cfs)</i>	<i>Load</i>		
		<i>Sediment (tons)</i>	<i>Nitrate-N (tons)</i>	<i>Total phosphorus (tons)</i>
2000	1236	4342	13.8	4.4
2001	3550	9839	84.9	5.1
2002	5440	34596	101.8	16.4
2003	1578	2955	26.4	1.8
2004	2787	7820	52.5	5.8
2005	5743	13793	112.2	10.2
2006	1053	2694	22.5	2.5
2007	3809	13410	75.4	10.6
2008	9437	83924	123.1	46.7
2009	7833	30921	117.7	13.9
2010	13539	56979	124.8	25.7
2011	6033	16786	72.8	9.9
2012	437	105	2.5	0.2
2013	4637	12309	123.9	6.0

Table 2-7. Summary of Annual Water Discharges, Sediment and Nutrient Loads at Cox Creek Monitoring Station (202)

<i>Water Year</i>	<i>Water discharge (cfs)</i>	<i>Load</i>		
		<i>Sediment (tons)</i>	<i>Nitrate-N (tons)</i>	<i>Total phosphorus (tons)</i>
2000	894	4153	10.3	5.7
2001	2833	9626	77.9	5.5
2002	4242	23207	100.6	16.1
2003	1226	1827	29.6	1.7
2004	1844	4597	45.3	3.7
2005	3976	8132	109.0	8.8
2006	806	3662	19.3	1.6
2007	3181	10105	81.5	7.2
2008	8097	73678	154.7	31.4
2009	5459	16331	135.9	8.6
2010	10040	27283	155.9	17.5
2011	4607	14021	91.5	9.6
2012	246	149	1.8	0.2
2013	3810	9906	149.7	5.2





Figure 2-7. Annual runoff at the five CREP monitoring stations





Figure 2-8. Annual suspended sediment loads at the five CREP monitoring stations





Figure 2-9. Annual nitrate-N loads at the five CREP monitoring stations





Figure 2-10. Annual ammonium-N loads at the five CREP monitoring stations





Figure 2-11. Annual Kjeldahl nitrogen loads at the five CREP monitoring stations





Figure 2-12. Annual phosphorus loads at the five CREP monitoring stations





Figure 2-13. Annual dissolved phosphorus loads at the five CREP monitoring stations





Figure 2-14. Annual ortho-phosphate phosphorous loads at the five CREP monitoring stations

Sediment and Nutrient Yields

To compare the different watersheds in terms of the amount of sediment and nutrient generated per unit area from each of the watersheds, the annual sediment and nutrient yields were computed by dividing the total annual load with the drainage area in acres for each of the monitoring stations. The results are provided in table 2-8 for sediment yield, table 2-9 for nitrate-N yield, and table 2-10 for total phosphorous. Sediment yields range from a low of 0.01 tons/acre for station 201 in WY2012 to a high of 9.57 tons/acre for station 202 in WY2008. Because of the high level of variability from year to year the average sediment yield for the nine years of data collection are compared in figure 2-15. The stations are arranged in order of their drainage area, with the station with the smallest drainage area (202) on the left and the station with the largest area (301) on the right. As can be seen in the figure, on the average the stations with the smaller drainage areas (202 and 201) yield higher sediment (about 2.0 ton/acre) than the stations with the larger areas (302, 303, 301) that yield less than 1.5 tons/acre.

Nitrate-N yields vary from a low of 0.5 lbs/acre for stations 201 and 202 in WY2012 to a high of 40.5 lbs/acre for station 202 in WY2010. For comparison purposes the average annual nitrate-N yield for the five stations is shown in figure 2-17. In general the stations with smaller drainage areas generate more nitrate per unit area than those with larger drainage areas, except for station 303 that is generating similar amounts as station 201 that has a smaller area.

Total phosphorous yields vary from a low of 0.03 lbs/acre for station 201 in WY2012 to a high of 8.81 lbs/acre for station 201 in WY2008. For comparison purposes, the average annual total phosphorous yield for the five stations is shown in figure 2-18. Similar to the nitrate-N yield, the stations with the smaller drainage areas generally generate more total phosphorous per unit area than those with larger drainage areas but the difference is very small.

Table 2-8. Sediment Yield in tons/acre for the CREP Monitoring Stations

<i>Water Year</i>	<i>CREP sediment yield (tons/ac)</i>				
	<i>201</i>	<i>202</i>	<i>301</i>	<i>302</i>	<i>303</i>
2000	0.41	0.54	0.62	0.42	0.60
2001	0.93	1.25	1.03	1.01	1.40
2002	3.26	3.01	1.48	1.76	1.25
2003	0.28	0.24	0.51	0.69	0.17
2004	0.74	0.60	0.17	0.12	0.31
2005	1.30	1.06	0.44	0.37	0.51
2006	0.25	0.48	0.19	0.25	0.16
2007	1.27	1.31	1.15	1.01	0.57
2008	7.92	9.57	0.97	1.19	0.46
2009	2.92	2.12	4.11	3.78	2.95
2010	5.38	3.54	3.44	4.01	2.63
2011	1.58	1.82	1.3	1.57	1.06
2012	0.01	0.02	0.10	0.13	0.06
2013	1.16	1.29	2.74	3.67	2.13
Avg.	1.96	1.92	1.30	1.43	1.02

Table 2-9. Nitrate-N Yield in lbs/acre for the CREP Monitoring Stations

<i>Water Year</i>	<i>CREP nitrate-nitrogen yield (lbs/ac)</i>				
	<i>201</i>	<i>202</i>	<i>301</i>	<i>302</i>	<i>303</i>
2000	2.6	2.7	6.2	5.2	9.2
2001	16.0	20.2	12.9	12.4	18.2
2002	19.2	26.1	9.6	11.8	14.5
2003	5.0	7.7	2.8	4.0	2.4
2004	9.9	11.8	3.6	4.5	8.1
2005	21.2	28.3	9.8	9.2	15.9
2006	4.2	5.0	4.0	4.4	6.4
2007	14.2	21.2	11.3	12.0	14.9
2008	23.2	40.2	12.5	14.3	12.9
2009	22.2	35.3	20.2	20.2	28.7
2010	23.6	40.5	20.0	20.2	33.2
2011	13.7	23.8	12.8	12.7	21.1
2012	0.5	0.5	1.7	1.8	3.1
2013	23.4	38.9	12.7	14.6	20.3
Avg.	14.2	21.6	10.0	10.5	14.9

Table 2-10. Total Phosphorus Yield in lbs/acre for the CREP Monitoring Stations

<i>Water Year</i>	<i>CREP total phosphorus yield (lbs/ac)</i>				
	<i>201</i>	<i>202</i>	<i>301</i>	<i>302</i>	<i>303</i>
2000	0.83	1.48	1.65	1.25	1.81
2001	0.95	1.44	1.84	1.53	3.28
2002	3.09	4.17	2.25	2.92	2.43
2003	0.34	0.45	0.86	1.10	0.47
2004	1.09	0.97	0.35	0.29	0.72
2005	1.93	2.28	0.96	0.92	1.05
2006	0.47	0.42	0.31	0.41	0.34
2007	2.00	1.86	2.20	1.72	1.35
2008	8.81	8.16	2.15	2.53	1.44
2009	2.62	2.23	5.50	5.45	4.87
2010	4.86	4.53	5.54	6.35	4.84
2011	1.86	2.50	2.04	3.03	1.94
2012	0.03	0.06	0.23	0.24	0.19
2013	1.13	1.34	4.46	5.40	4.20
Avg.	2.14	2.28	2.17	2.37	2.07




Figure 2-15. Average annual sediment yield in tons/acre for the CREP monitoring stations




Figure 2-16. Average annual nitrate-N yield in lbs/acre for the CREP monitoring stations




Figure 2-17. Average annual total phosphorous yield in lbs/acre for the CREP monitoring stations

Additional CREP Data Collection Efforts

In addition to the CREP monitoring in the Court/Haw and Panther/Cox watersheds, that was initiated in 1999, several additional monitoring efforts have been initiated by the ISWS through the CREP project in order to provide additional information on the role BMPs in reducing sediment and nutrient yields and to better define the context of existing CREP data on a larger watershed scale.

During September of 2006 in response to significant CREP enrollments and an intensive restoration effort by the Natural Resources Conservation Service (NRCS), two additional monitoring stations (table 2-11) were installed in the Cedar Creek watershed, located in the Spoon River basin (figure 2-18). Station 306 is located on the right descending bank of the mainstem of Cedar Creek where it intersects CR 000 E in Fulton County (border with Warren Co). The second gage, station 305, is located near the left descending bank of Swan Creek, a major tributary of Cedar, where it flows beneath CR 000 E Fulton County, approximately 2.1 miles south of the Cedar Creek (306) gage.

Table 2-11. Additional CREP Monitoring Stations in the Spoon River Watershed

<i>Station ID</i>	<i>Name</i>	<i>Drainage area</i>	<i>Location</i>	<i>Watershed</i>
305	Swan Creek	98.1 sq mi (254 sq km)	N 40.67700 W 090.44391	Spoon River
306	Cedar Creek	146.2 sq mi (379 sq km)	N 40.70847 W 090.44540	Spoon River
RG39	Rain Gage 39	NA	N40.79145 W090.49999	Spoon River




Figure 2-18. Locations of monitoring stations in the Cedar and Swan watersheds

Both watersheds are located in the Galesburg Plain physiographic region. The topography is flat to gently rolling and the soils are primarily loess. Stream channels and associated floodplains are heavily dissected with stream channels commonly being incised into the floodplain. Both watersheds are mostly rural with agriculture the predominant land use. Pasture and woodlands are also common due to the topography introduced by the dissected stream channels.

Both gages became operational near the end of WY2006 (9/15/2006) and are instrumented and operated as are all CREP gages, in accordance to the CREP QAPP (Appendix A). Both stations utilize a pressure transducer to determine stage, log data on a 15 minute time step and are equipped with an ISCO automated pump sampler slaved to the stage sensor in order to augment manual discrete sampling efforts. Thirty-eight and thirty-three discharge measurements have been collected at stations 305 and 306 respectively in an effort to establish a reliable rating in as short a time as possible. Based on provisional data, summary statistics for suspended sediment concentration data is provided in table 2-12.

In addition to the two streamgages the ISWS has installed a recording raingage immediately east of CR1500E and approximately 0.5 mi north of CR1100N in Warren Co. The raingage is a modified Belfort equipped with a linear potentiometer, in order to provide a digital output, and can be operated throughout the year. Raingage deployment and maintenance as well as the download and reduction of precipitation data can be found in the CREP QAPP (Appendix A).

ISWS field staff began suspended sediment sampling at two U.S. Geological Survey (USGS) gages located on the mainstem of the Spoon River on 3/29/2004. Samples are collected weekly at both sites with additional samples collected during runoff events. Sampling at London Mills (05569500) is done from the Route 116 bridge where the USGS gaging station is located. Sediment sampling at Seville (05570000) is done approximately 1 mile downstream of the current USGS gage location on State Route 95. Current USGS sediment data are also collected at this location. As of 9/30/12, 568 samples have been collected at London Mills while 521 samples have been collected at Seville. Summary statistics for suspended sediment concentration data collected through WY2012 are presented for each station in Table 2-13.

**Table 2-12. Suspended Sediment Concentration Data (mg/L)
for Swan and Cedar Creeks**

	<i>Swan (305)</i>	<i>Cedar (306)</i>
Count (samples)	3515	3623
Mean	380.1	471.3
Max	7872.6	8101.8
Min	1.99	1.59
Median	137.1	132.6
25 th Percentile	49.3	51.0
75 th Percentile	416.3	462.7

**Table 2-13. Suspended Sediment Concentration Data (mg/L)
for Spoon River at London Mills and Seville**

	<i>London Mills (05569500)</i>	<i>Seville (05570000)</i>
Count (samples)	568	521
Mean	296.1	293.1
Max	4952.7	4730.7
Min	1.91	3.93
Median	116.0	122.2
25 th Percentile	49.9	58.8
75 th Percentile	285.7	266.7

3. Land Use Cover and Conservation Practices

The Illinois River Basin (IRB) is nearly 16 million acres with a diverse range of land covers. The extent of these land covers is illustrated in figure 3-1 using the U.S. Department of Agriculture (USDA) National Agricultural Statistics Service (NASS) 2010 land cover data. The database contains almost 150 land cover category types. For the purpose of this study those types have been grouped into 11 categories: corn, soybeans, small grains, other row crops, other crops, grass/pasture, developed, woodland, wetlands, water, and other. In 2010 the Illinois River Basin is dominated by agricultural land, comprising of 72% of the basin. Corn and soybean acreage accounts for most of the agricultural land cover. Developed urban type lands, woodlands, and grass/pasture lands are the next highest with 14%, 12%, and 9%, respectively.

Outside of natural factors such as the physical settings and climate variability, land use is the main driving factors that affect a watershed's hydrology, erosion, sedimentation, and water quality. It is therefore important to document and analyze changes in land use for a given watershed to properly understand and explain changes in its hydrology, water quality, and the erosion and sedimentation process. The Illinois River basin has undergone significant changes in land use practices during the last century. These changes have been used to explain degradation in water quality and aquatic habitat along the Illinois River. In recent years, there have been significant efforts at the local, state, and federal level to improve land use practices by implementing conservation practices throughout the watershed. The Illinois River CREP is a course of major state and federal initiatives to significantly increase conservation and restoration practices in the Illinois River basin.

Historical Agricultural Land Use Trends in Illinois

To provide a historical perspective to changes in land use practices in the Illinois River basin, we have compiled and analyzed historical land use data from different sources for the whole state. The earliest land use data is based on the Illinois Agricultural Statistics (IAS) records. The IAS data shows that in 1866 approximately 23 percent of the state's land area was in agricultural crop production (figure 3-2). In 2006, agricultural production has increased to 65 percent of the state's land. From 1866 through to the 1920s, crop production increased from 8 to 18 million acres mostly due to a three-fold increase in small grain (wheat, oats, and hay) acreage. In the 1920s small grain acreage began to decline in favor of soybeans. Essentially, from this period to present, a steady reversal in acreage has occurred between small grains and soybeans such that current soybean acreage is the same as was small grains were in the 1920s. From 1866 to 2006, total Illinois land area in crop production increased by more nearly tripled from 8 to 23 million acres. The dominant crops in 1866 were corn and small grains, whereas corn and soybeans (row crops) acreage was 93 percent of the total crop acreage in 2006. During the period of record (1866-2006), corn acreage has remained fairly steady at 9.3 million acres. Corn was harvested on 4.9 million acres in 1866 but increased to the long-term average acreage by 1881. Acreage peaked in 2005 at 12.1 million acres and was 11.3 million acres in 2006. From 1925 to 2006 crop acreage increased by 23 percent.




Figure 3-1. Land cover of the Illinois River Basin (NASS, 2010)

In 1925, IAS began delineating agricultural crop production data by county, rather than as a state total, which allows for the estimation of crop acreage by basins. The Illinois River Basin (IRB) is nearly half of the Illinois land area, and occupies over 18 million acres when the watershed area in the states of Indiana and Wisconsin are included. Figure 3-3 shows similar trends in crop production as was seen for the State of Illinois. In 1925, 51 percent (9.4 million acres) of the IRB land area was in crop production while in 2006, 56 percent (10.3 million acres) was in crop production. The same reversal of small grain and soybean acreage is also seen. Corn acreage is fairly steady for the period of record, averaging 4.8 million acres, increasing from 4.4 to 6.0 million acres from 1925 to 1976, and slightly decreasing to 5.5 million acres in 2006. Total IRB watershed area in crop production increased by 9 percent from 1925 to 2006 which is smaller than the 23 percent increase for the whole State of Illinois during the same period.

The Spoon River watershed is one of ten major tributaries to the Illinois River with a drainage area of 1.2 million acres (6.5 percent of the IRB drainage area). From 1925 to 2006 watershed area in crop production increased from 54 to 66 percent. Figure 3-4 shows that the trends in corn, small grains, and soybeans are also similar. Corn and small grain acreage was 0.64 million acres in 1925 and in 2006 corn and soybeans were 0.75 million acres. Corn acreage increased by 0.19 million acres from 1925 to 1976 and then decreased by 0.09 million acres through 2006. The total Spoon River watershed area in crop production increased by 22 percent during 1925-2006 period and is only slightly below that of the increase in the State of Illinois and higher than the 9 percent increase for the IRB.




Figure 3-2. Acreage of agricultural land uses in State of Illinois (1866-2006)




Figure 3-3. Acreage of agricultural land uses in Illinois River basin (1925-2006)




Figure 3-4. Acreage of agricultural land uses in Spoon River watershed (1925-2006)

The Sangamon River watershed has a drainage area of 3.4 million acres (18.5 percent of the IRB drainage area). From 1925 to 2006, watershed area in crop production increased from 67 to 78 percent. Figure 3-5 shows that the trends in corn, small grains, and soybeans are also similar to the IRB. Corn and small grain acreage was 2.2 million acres in 1925 and in 2006 corn and soybeans were 2.6 million acres. Corn acreage increased by 0.37 million acres from 1925 to 2006. The total Sangamon River watershed area in crop production increased by 17 percent during 1925-2006 period and is below that of the increase in the State of Illinois and higher than the 9 percent increase for the IRB.

Overall, total crop acres within the Sangamon and Spoon River watersheds steadily increased from 1925 to the early 1980s and then remained steady through 2006. The Illinois River Basin and the entire State of Illinois show the same trend for total crop acres.




Figure 3-5. Acreage of agricultural land uses in Sangamon River watershed (1925-2006)

Land Cover

The USDA National Agriculture Statistics Service (NASS) land cover data has been available since 1999. In 2006 an evaluation of the usefulness of the crop data layers for annual land cover information in Illinois was undertaken by the Illinois State Geological Survey (ISGS) and NASS (Luman, 2008). Based on inherent errors associated with satellite data, irreparable mechanical problems with older multispectral imagery satellites and land cover classification methods used to interpret that imagery, new enhanced CDL protocols were established in 2007 for Illinois. Consequently, land cover misclassifications were identified prior to the new protocol, which became more apparent when evaluating the land cover in the monitored watersheds (figure 3-6): Panther Creek (201), Cox Creek (202), Court Creek (301), North Creek (302), and Haw Creek (303). Therefore, any

changes in land cover for the watersheds that are being monitored will be evaluated beginning in 2007 through 2013 which is the most currently available NASS CDL data.

The five monitored watersheds have somewhat different ratios of land cover types. The Panther and Cox Creek watersheds in the Spoon River watershed have 53 and 73 percent area in agriculture and 47 and 27 percent in non-agriculture land covers, respectively (table 3-1). The main difference is Panther Creek has over 20 percent more land in forest/shrubland than Cox Creek, due to a large portion of the watershed lies in the Panther Creek State Conservation Area. Agriculture land cover is 44 and 56 percent in Court and Haw Creeks, respectively, while the non-agriculture area is the inverse. North Creek watershed, a tributary of Court Creek, has a larger portion of land area in forest/shrubland than Haw Creek. Figure 3-7 illustrates the percent change in total watershed acres between 2007 and 2013 for six generalized land cover categories in each of the five monitored tributary watersheds in the Illinois River Basin. Agriculture land covers were categorized into Corn, Soybeans, Double Crop with Soybeans and Other Cropland, as well as summed in one category identified as Agriculture. Non-agriculture land covers were categorized into Grassland and Forest/Shrubland, and summed as Non-Agriculture. All five watersheds had a 5 percent reduction in non-agricultural land cover area (Grasslands and Forest/Shrubland) between 2007 and 2013. An increase in agricultural land cover area (Corn, Soybeans, Double Crop with Soybeans and Other Cropland) ranged from 2 to nearly 11 percent occurred on all five watersheds. The three Spoon River tributary watersheds (Court, North, and Haw Creeks) had marked percent increases in soybean acres and decreased percent of corn acres. The two Sangamon River watersheds (Panther and Cox Creeks) had an increase percent of corn acres, with Panther having an increase percentage of soybean acres and Cox with an increase in other cropland acres.

Figures 3-8 to 3-12 show the changes in each land cover for each year between 2007 and 2013. For this report, NASS Cropland Data Layer (CDL) categories for the monitored watersheds were combined into 6 general land cover categories: 1) corn, 2) soybean, 3) other cultivated crops, 4) grassland, 5) forest/shrubland and 6) developed, barren, open space, water and wetlands. Land cover area changes between years is represented in acres. Therefore, some watersheds may appear to have greater changes in acreage from year to year but may only represent a small percentage of the watershed depending on the total watershed acres. Panther Creek watershed (figure 3-8) acres remained constant for move land covers when comparing 2007 and 2013. Corn and soybean acres shifted between years and inversely as reflected by normal corn and soybean rotation practices. Forest/shrubland saw a minor shift in 2010. Cox Creek watershed (figure 3-9) saw similar variability as Panther Creek watershed in most land cover acreage. Only minor increases in acres for cultivated crops and developed, barren, open space, water and wetlands. Court Creek (figure 3-10) appeared to have corn and Grasslands trade acres each year, with Corn increasing to a high in 2011 and then returning to near 2007 acreage. Soybean acres increased every other year for a seven year increase. Forest/Scrubland acres decreased slightly with little variability. North Creek watershed (figure 3-11) is a subwatershed within Court Creek watershed explaining the significant reduction in total watershed acres. The same patterns and variability as Court Creek watershed appear here. Finally, Haw Creek watershed (Figure 3-12) land cover patterns and variability in acreages were similar to Court/North Creek watersheds.




Figure 3-6. Watersheds being monitored for hydrology, sediment and nutrients.

Table 3-1. 7-year average (2007-2013) percent acres of land cover area by watershed

	<i>ISWS Station Number</i>				
	<i>201</i>	<i>202</i>	<i>301</i>	<i>302</i>	<i>303</i>
Corn	31	46	28	26	36
Soybeans	21	26	16	16	20
Other Crops	1	2	0	0	0
Grasslands	11	13	20	20	17
Forest/Shrubland	32	11	29	34	21
Developed, Barren, Open Space, Water, Wetlands	4	3	7	4	6
AGRICULTURE	53	73	44	42	56
NON-AGRICULTURE	47	27	56	58	44




Figure 3-7. Illinois River Basin Watersheds: Percent Change in Generalized NASS Land-Use from 2007-2013.




Figure 3-8. Panther Creek Watershed from ISWS Station 201: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013.




Figure 3-9. Cox Creek Watershed from ISWS Station 202: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013.




Figure 3-10. Court Creek Watershed from ISWS Station 301: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013.




Figure 3-11. North Creek Watershed from ISWS Station 302: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013.




Figure 3-12. Haw Creek Watershed from ISWS Station 303: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013.

Conservation Practices

There has been a significant increase in the implementation of conservation practices in Illinois in recent years with CREP making a major contribution. Figure 3-13 shows the location of approved Illinois CREP contracts from the state of Illinois as of 2013. With this type of information it will be possible to identify areas where there has been significant participation in the CREP program and where changes in sediment and nutrient delivery should be expected. The information will provide important input data to the watershed models that are being developed to evaluate the impact of CREP practices.

There are many conservation practices implemented through the watersheds as a result of federal and state conservation reserve programs. In order to evaluate watershed monitoring efforts, knowing the when and what conservation practices are implemented in the watershed is important. Figures 3-14 and 3-15 are examples of cumulative acres of conservation practices installed in a couple of the monitored watersheds from 1999 through 2013. Riparian and wetland buffers, SAFE habitat and grass filter strips are the most installed conservation practice in Court Creek with most of the acres occurring prior to 2009. Whereas, permanent wildlife habitat was the most installed practice installed prior to 2005 in the Haw Creek watershed. Grass filter strips, riparian buffers, wetland restoration and SAFE habitat are the next most installed.




Figure 3-13. State CREP contract locations (IDNR, 2014).




Figure 3-14. Cumulative acres of conservation practices installed in Court Creek watershed at monitoring station ISWS #201 from 1999-2013.




Figure 3-15. Cumulative acres of conservation practices installed in Haw Creek watershed at monitoring station ISWS #202 from 1999-2013.

4. Variability and Trends in Precipitation and Streamflow

Results of a short-term monitoring program have to be viewed with respect to the climatic and hydrologic conditions under which the data was collected. Under ideal conditions, which rarely happen, the monitoring period would include a combination of wet, dry, and normal climatic conditions that represent the range of variability in climatic and hydrologic conditions in the watershed. The influence of climatic and hydrologic conditions on the data collected has been taken into consideration, especially when different datasets collected at different times and conditions are combined or compared. The Illinois River basin, as any major watershed, has experienced significant variability in precipitation and streamflow over the last century and recent periods. Data collection for the CREP program started in 1999 to provide a perspective as to how the current monitoring period compares to the long-term variability of precipitation and streamflows within the Illinois River basin. Historical precipitation and streamflow data are analyzed and presented in this segment of the report.

Climate and hydrologic records from the past 100 years in Illinois show considerable long-term variability. These variabilities and trends were analyzed for two stations on the Illinois River and six tributary stations in the Illinois River basin (figure 4-1). Figure 4-2 compares average precipitation and streamflow for the Upper Illinois River watershed since the 1880s, as expressed in moving 10-year average values. Similar comparisons are shown in figures 4-3 to 4-8 for the Fox, Kankakee, Spoon, Sangamon, LaMoine, and Macoupin subwatersheds, respectively, but for shorter time periods as limited by the available gaging records. Figure 4-9 for the entire Illinois River Basin (at the Valley City streamgage) is nearly identical to figure 1 except for the period of record. The 10-year average precipitation and streamflow values plotted in figures 4-2 to 4-9 represent the approximate midpoint of the 10 years; for example, the value for 1995 represents the average for 10 years from 1990-1999, the value for 1996 represents the average for the 10 years 1991-2000, and so forth. Streamflow values are expressed in inches of water spread uniformly over the entire watershed such that average streamflow can be compared directly with precipitation for the concurrent period. Streamflow values in figure 4-2 are computed from flow and stage records at Peoria prior to 1940 and at Kingston Mines since 1940.

Figure 4-2 shows that precipitation and streamflow in the Upper Illinois River watershed from 1970 to 1995 were considerably higher than at any other time in the 20th Century. Prior to 1895, precipitation for the Illinois River watershed is estimated from a small set of gaging records dating back to 1870. These precipitation records show that there was a decade of high precipitation in the late 1870s and early 1880s similar in magnitude to high precipitation amounts during 1970-1995. A comparison of 10-year average precipitation and streamflow amounts clearly shows that streamflow has been very closely related to concurrent precipitation throughout the past 125 years, with a correlation coefficient (r) of 0.958.




Figure 4-1. Location of streamgaging stations with long-term data used in the analysis of variability and trends




Figure 4-2. Ten-year average precipitation and streamflow, Illinois River at Peoria-Kingston Mines




Figure 4-3. Ten-year average precipitation and streamflow, Fox River at Dayton




Figure 4-4. Ten-year average precipitation and streamflow, Kankakee River at Momence




Figure 4-5. Ten-year average precipitation and streamflow, Spoon River at Seville




Figure 4-6. Ten-year average precipitation and streamflow, Sangamon River at Monticello




Figure 4-7. Ten-year average precipitation and streamflow, LaMoine River at Ripley




Figure 4-8. Ten-year average precipitation and streamflow, Macoupin Creek near Kane




Figure 4-9. Ten-year average precipitation and streamflow, Illinois River at Valley City

Precipitation and streamflow trends shown in figure 4-2 are consistent with regional trends that have affected northern Illinois and much of the upper Midwest (Knapp, 2005). Statistical analyses of long-term streamflow records by Knapp (2005) using the Kendall tau-b trend statistic indicate that streamgauge records in northern Illinois, eastern Iowa, and Minnesota all exhibit increasing trends in average streamflow (figure 4-10). Conversely, long-term flow records in the southern two-thirds of Illinois generally do not show significant increases in streamflow.

Figures 4-2 to 4-9 illustrate that trends in precipitation and streamflow vary across the Illinois River watershed. Increasing trends are particularly evident in the Upper Illinois River watershed and its two primary tributaries, the Fox and Kankakee River (figures 4-3 and 4-4). In contrast, the Macoupin, LaMoine, and Sangamon River subwatersheds, in the southern portion of the Illinois River basin, show much less or no overall trend in precipitation or streamflow — even though these records show considerable variation in precipitation and streamflow from decade to decade. The Spoon River watershed, having an intermediate location, shows an increasing trend in flow amount, but to a lesser degree than the Fox and Kankakee River watersheds located farther to the north. In all cases, there is a strong correlation between average precipitation and streamflow.

The significance of the trends is identified using the Kendall tau-b statistic. The Kendall tau-b statistical test provides a quantitative measure of trend, with a coefficient value of 0 indicating no trend and a value of 1 indicating an absolute increasing trend. For the 93-year flow records dating back to 1915, a coefficient value greater than or equal to 0.115 indicates an increasing trend at a 90 percent confidence level, and a value greater than or equal to 0.162 indicates an increasing trend at a 98 percent confidence level. Table 3-2 shows the Kendall Tau-b trend coefficients computed for two time periods, 1915-2007 and 1970-2007. The 1915-2007 trend analyses for the Fox, Kankakee, and Upper Illinois (Peoria-Kingston Mines) flow records show increasing trends with very high levels of confidence. The 1915-2007 trend analysis for the Spoon River record shows an increasing trend, with roughly a 94 percent level of confidence. The flow records for the tributaries located farther south in the watershed do not show a significant trend (having less than an 80 percent level of confidence). The 1915-2007 trend coefficient for the Illinois River at Valley City is not shown because the flow record does not date back to 1915.

Although flow records from the northern half of the Illinois River watershed display a general increasing trend over their full period of record, a closer look indicates: 1) there was a geographically widespread and sizable jump in average flow amount between the 1960s and 1970s (this jump also occurred in the southern part of the basin to a lesser extent); and 2) for most locations there has been little or no additional increase since the 1970s. In fact, for most locations, the average flows since 1995 have declined from the high flow levels that occurred from 1970 to 1995. Table 3-3 presents the average annual precipitation and streamflow amounts for the Illinois River and its major tributaries over the past 12 years (1996-2007) and compares these amounts to those for earlier periods (1915-1969 and 1970-1995) and to the overall long-term record. Except for the Kankakee River, the average flow from 1996-2007 for these rivers is much closer to the long-term average than it is to the higher flow amounts that were experienced



Figure 4-10. Locations of long-term streamflow gages (at least 89 years of record) showing statistically significant trends in mean annual flow in the eastern United States (from Knapp, 2005)

from 1970 to 1995. Thus, with the exception of the Kankakee River watershed, it is reasonable to conclude that other flow records collected throughout the Illinois River watershed over the 1996-2007 timeframe may represent conditions similar to their expected long-term average condition.

Although it is not possible to predict how these trends will progress in the future, concerns expressed in previous decades regarding the potential for continued increases in flows throughout the Illinois River watershed (for example by Ramamurthy et al., 1989) for the time being may no longer be an issue. If anything, there may be growing concerns that the occurrence of drought periods such as existed prior to 1970 may become more frequent. This analysis does not specifically look at trends of flooding or low flows. However, for long-term gaging records in the Illinois River watershed, Knapp (2005) found that trends in high flows and low flows tended to be coincident and proportional to trends in average flow.

Table 4-1. Kendall Tau-b Trend Statistics for Flow Records on the Illinois River and Major Tributaries

<i>Streamgage record</i>	<i>Kendall Tau-b coefficient value period-of-record used in the analysis</i>	
	<i>1915-2007</i>	<i>1970-2007</i>
Fox River at Dayton	0.294	-0.135
Kankakee River at Momence	0.316	-0.007
Illinois River at Peoria-Kingston Mines	0.315	-0.144
Spoon River at Seville	0.127	-0.127
Sangamon River at Monticello	0.087	-0.081
LaMoine River at Ripley	0.075	-0.166
Macoupin Creek near Kane*	-0.009	-0.081
Illinois River at Valley City**	-----	-0.112

Notes:

* The periods of record for the Macoupin Creek gage near Kane are 1921-1933 and 1941-2007.

** The flow record at Valley City only extends back to 1939. The trend coefficient for the 1939-2007 period at Valley City, 0.162, is somewhat less than the trend coefficient for Peoria-Kingston Mines for the same time period (0.192).

Table 4-2. Average Annual Precipitation and Streamflow (inches) for Different Periods of Record

Precipitation

<i>Watershed</i>	<i>1915-2007</i>	<i>1915-1969</i>	<i>1970-1995</i>	<i>1996-2007</i>
Fox	33.7	32.6	35.9	34.4
Kankakee	37.0	35.5	39.5	38.4
Upper Illinois (Peoria)	36.3	35.2	38.3	37.1
Spoon	35.7	34.9	37.7	34.8
Sangamon	38.9	38.1	40.7	38.9
LaMoine	36.6	35.8	38.6	35.9
Macoupin	37.4	37.0	38.6	36.9
Entire Illinois (Valley City)	36.5	35.6	38.3	36.6

Streamflow

<i>Watershed</i>	<i>1915-2007</i>	<i>1915-1969</i>	<i>1970-1995</i>	<i>1996-2007</i>
Fox	9.3	7.7	12.1	10.0
Kankakee	12.3	10.9	14.7	13.5
Upper Illinois (Peoria)	10.2	8.8	12.9	10.8
Spoon	9.1	8.0	11.3	9.2
Sangamon	10.4	9.5	12.4	10.1
LaMoine	8.7	7.7	10.7	8.2
Macoupin	8.4	8.1	9.1	7.8
Entire Illinois (Valley City)	9.8	8.4	11.7	9.5

5. Model Development and Application

The Illinois State Water Survey has been developing a watershed model for the Illinois River basin in support of the Illinois River Ecosystem project. In the initial phase, a hydrologic model of the entire Illinois basin has been developed and used to evaluate potential impacts of land use changes and climate variability on streamflow in the Illinois River basin. The model is based on the U.S. Environmental Protection Agency's BASINS 3.0 modeling system. The Hydrologic Simulation Program – FORTRAN or HSPF (Bicknell et al., 2001) which is part of BASINS was used to simulate the hydrology of the Illinois River basin. The HSPF is a comprehensive and dynamic watershed model that also has the capability to simulate water quality and sediment transport.

To make the model applicable for assessing and evaluating the impact of CREP and other land use changes on water quality and sediment transport, the Water Survey has been developing the sediment transport and water quality capabilities of the HSPF model for the Illinois River basin. The initial effort has focused on the Spoon River watershed (figure 5-1) where two of the four intensively monitored watersheds, Court and Haw Creek, are located. Streamflow, sediment, and water quality data being collected at three monitoring stations are being used to calibrate and test the model for the Spoon River watershed. Once the calibration and validation process are completed for the Spoon River watershed, the model parameters can be used to develop models for other similar watersheds to simulate the hydrology, sediment transport and water quality under different climatic and land use scenarios. Over time, as land use practices change significantly as a result of CREP and other conservation practices, the models being developed will provide the tools to evaluate and quantify changes in water quality and sediment delivery to the Illinois River.

The progress in model development for the Spoon River watershed is discussed in the following sections.

HSPF Model

The HSPF model is a conceptual, comprehensive, long term continuous simulation watershed scale model which simulates non-point source hydrology and water quality, combines it with point source contributions, and performs flow and water quality routing in the watershed and its streams. The HSPF model simulates land-surface portion of the hydrologic cycle by a series of interconnected storages – an upper zone, a lower zone, and a ground-water zone. The fluxes of water between these storages and to the stream or atmosphere are controlled by model parameters. The model uses a storage routing technique to route water from one reach to the next during stream processes.

For sediment simulation, the surface erosion component of the HSPF model performs processes such as sediment detachment from the soil matrix in the pervious land segments during rainfall event, washoff of this detached sediment, scour of the soil matrix, and reattachment or compaction of the sediment. Storage and washoff of sediments from the impervious surfaces is



Figure 5-1. Location of the Spoon River watershed

also considered. The sediment load and transport in the stream channel is dependent on the particle diameter, density, fall velocity, shear stress for deposition and scour, and erodibility. The noncohesive (sand) and cohesive (silt and clay) sediment transport is simulated in the model using different subroutines.

Nutrients in the watershed soil in the HSPF model are simulated either as attached to organic or inorganic solids, dissolved in the overland flow, or as concentrations in the subsurface flow reaching the streams laterally. For both nitrogen and phosphorous compounds, the processes simulated include immobilization, mineralization, nitrification/denitrification (nitrogen only), plant uptake, and adsorption/desorption. The nutrient loads from the watershed undergo further transformation in the stream reaches.

Model Input Data

The HSPF model requires spatial information about watershed topography, river/stream reaches, land use, soils, and climate. The hourly time-series of climate data required for hydrologic simulations using HSPF include precipitation, potential evapotranspiration (ET), potential surface evaporation, air temperature, dew-point temperature, wind speed, and solar radiation. The hourly precipitation data from the two ISWS gages, one each in Court Creek (ISWS31) and Haw Creek (ISWS32) watersheds, were used (figures 5-2 and 5-3). Daily precipitation data from the MRCC (Midwestern Regional Climate Center) gaging station at Galesburg (ID 113320) was also used after it was disaggregated into hourly data based on the hourly precipitation data from an ICN (Illinois Climate Network) station located in Monmouth (MON). The other time series of the climate inputs for the above three precipitation stations were obtained from the ICN station at Monmouth. Daily data from nine additional MRCC stations (figure 5-4) in or near the Spoon River watershed were also disaggregated into hourly data based on the hourly data from three stations at Peoria, Moline, and Augusta, as found in the BASINS database. These additional stations were used for the Spoon River watershed model.

For topographic inputs, the 30-meter Digital Elevation Model (DEM) raster dataset produced by the Illinois State Geological Survey (ISGS) and the United States Geological Survey (USGS) was used. The high resolution National Hydrography Dataset (NHD) developed by the USGS was used to provide stream/river reach information to the model. The land use data were obtained from the Illinois Department of Agriculture which is based on the satellite imagery of the State of Illinois acquired from three dates during the spring, summer, and fall seasons of 1999 and 2000. Land use in the study watersheds was classified as corn, soybean, rural grassland, forest, urban, wetland and other (figures 5-5, 5-6, and 5-7). The soils data were based on digitized County Soil Association Maps of the Knox County and the STATSGO dataset (figure 5-8). The soil type for various parts of the study watersheds were determined spatially from the digitized soils maps, but the parameters corresponding to the soil type were manually entered during development of the HSPF model.




Figure 5-2. Schematic of the subwatershed and stream delineation, and precipitation gages used for the Haw Creek model



Figure 5-3. Schematic of the subwatershed and stream delineation, and precipitation gages used for the Haw Creek model




Figure 5-4. Schematic of the subwatershed and stream delineation, and precipitation gages used for the Spoon River watershed model




Figure 5-5. Land use in the Court Creek watershed




Figure 5-6. Land use in the Haw Creek watershed




Figure 5-7. Land use in the Spoon River watershed




Figure 5-8. Soil types in the Spoon River watershed

Model Development

Based on the topographic and hydrographic data, the watersheds were subdelineated into smaller hydrologically-connected subwatersheds and stream reaches, and respective outlets. The Automatic Delineation procedure in BASINS with an option of ‘burning in’ existing streams was used. Subdelineation was done for representing spatially variable physical and other characteristics of a watershed in the HSPF model. The Court, Haw, and Spoon River watersheds were subdivided into 31, 25, and 42 subwatersheds, respectively (figures 5-2, 5-3, and 5-4). During subdelineation, outlets were specified in the models corresponding to the streamflow gaging/water quality monitoring stations on the North Creek (ISWS302), Court Creek (ISWS301), Haw Creek (ISWS303), and the USGS streamflow gaging station at Seville (USGS05570000) in the Spoon River watershed (figures 5-2, 5-3, and 5-4). The subwatersheds were further subdivided into Hydrologic Response Units (HRUs) based on land use, soil, and climate to account for the spatial variability of a basin’s physical and hydrologic characteristics at a finer scale. An HRU is an area within a watershed that is expected to have a similar hydrologic response to input of precipitation and evapotranspiration. Each HRU has a set of parameter values that must be determined through the calibration process to define runoff characteristics as well as loading of various constituents from that HRU. In the Court Creek watershed HSPF model, climate data from the Court Creek and Galesburg precipitation gages were input to different subwatersheds based on the proximity. Similarly, in the Haw Creek HSPF model data from the Haw Creek and Galesburg gages were input to various subwatersheds. In case of Spoon River watershed HSPF model, data from all ten MRCC stations were specified for different subwatersheds based on their proximity to the gages.

Model of the Court Creek watershed was developed first using two years (WY2001-WY2002) streamflow and sediment concentration data from the ISWS301 streamflow gage/WQ station on the Court Creek. Calibrated model parameters from this model were then used to populate the models of the Haw Creek and Spoon River watersheds. No further calibration of these two models was performed. Haw Creek watershed model was run for the same two year period as Court Creek watershed model and the model results were compared with the observed data from the ISWS303 gage on the Haw Creek. Since long-term climate and streamflow data were available for the Spoon River watershed, this model was run for 1972-1995 period using data from the USGS05570000 at Seville.

Modeling Results

Values of a large number of HSPF model parameters cannot be obtained from field data and need to be determined through model calibration exercise. The Court Creek watershed model was calibrated to assign best possible parameter values to each HRU and stream reach so that the model simulated daily streamflows and pollutant concentrations similar to the values observed at the gaging/monitoring stations. Calibration of the hydrologic component of the model was followed by the calibration of the water quality component for the sediment concentration. Model was run for hourly time step. For the two year calibration period of WY2001-WY2002, percent volume error between the model simulated and observed streamflows at gages ISWS301 on the Court Creek and ISWS302 on the North Creek were 1.2% overestimation, and 3.5%

underestimation, respectively. Comparisons of the daily streamflows simulated by the model for WY2001-WY2002 period with those observed at gages ISWS301 and ISWS302 are shown in figures 5-9a and 5-9b. The performance of this preliminary model is promising and overall the simulated streamflows follow the similar trend as the observed values. The timings and shape of the simulated streamflow hydrographs resemble the observed ones but some peak flows were underestimated by the model. In this study the model was not calibrated to match the individual stormflow events, rather it was calibrated to fit the long-term and daily data over the two year calibration period. Also, data from only two precipitation gaging stations, both near the boundary of the watershed (figure 5-2), were used to spatially represent the precipitation over the entire watershed. It is possible that rainfall measured for a particular event at one of the gages did not represent the rainfall that actually occurred in different parts of the watershed, thereby resulting in discrepancies between the observed and simulated streamflow hydrographs. Thus, more precipitation gaging stations will help improve the performance of the hydrologic model by more accurately simulating the stormflow hydrographs.

For sediment simulation by the model in the Court Creek watershed, parameters controlling soil erosion on the surface and sediment transport in the stream channel were calibrated. Comparison of sediment concentration simulated by the model and those observed at gages ISWS301 and ISWS302 are shown in figure 5-10 for the WY2001-WY2002 period. The simulated values generally followed the same trend as the observed sediment concentration values at both gages. Since most soil erosion occurs during extreme runoff events, some high sediment concentrations were underestimated by the model as a result of poor estimation of the stormflow peaks by the model during hydrologic simulations.

Streamflow and sediment concentration simulation results from the Haw Creek watershed model are compared with the observed data as shown in figures 5-11 and 5-12, respectively. Similar results from the Spoon River watershed model are shown in figures 5-13 and 5-14. In this preliminary phase, the performances of these two models were similar to the calibrated model of the Court Creek watershed. Performance of these models can be improved in the future if climate, streamflow, and water quality data are available for more stations and longer time period to improve the model calibration.





Figure 5-9. Results of model calibration for streamflow simulation for the Court Creek watershed





Figure 5-10. Preliminary results of model calibration for suspended sediment concentration simulation for the Court Creek watershed




Figure 5-11. Comparison of observed and simulated streamflow by the Haw Creek watershed model developed using the calibrated parameters from the Court Creek watershed model




Figure 5-12. Preliminary results for suspended sediment concentration from the Haw Creek watershed model developed using the calibrated parameters from the Court Creek watershed model




Figure 5-13. Comparison of observed and simulated streamflow simulation by the Spoon River watershed model developed using the calibrated parameters from the Court Creek watershed model




Figure 5-14. Preliminary results for suspended sediment concentration from the Spoon River watershed model developed using the calibrated parameters from the Court Creek watershed model

6. Analyses and Discussion

Sediment Loadings

Based on sediment records since 1980, the Illinois River on the average receives approximately 12 million tons of sediment annually from tributary streams (Demissie et al., 2004). About 55 percent of the sediment delivered to the river (6.7 million tons) is deposited in the river, backwater lakes, and side channels along the river. Most of this sediment is generated in the tributary watersheds to the Lower Illinois River, with the Spoon and LaMoine River watersheds as the highest per unit area generators of sediment among the major tributaries. The smaller tributaries draining directly to the river also contribute significant sediment. Controlling the erosion processes that are producing excessive sediment and reducing sediment delivery to the Illinois River will be a long-term effort, since sediment storage and mobilization along major rivers is a slow process. It will take some time to flush the sediment already in the system. In the initial phase of a restoration project, the major goal is to stabilize the system so that the erosion process is not accelerating and generating more sediment. The readjustment processes will take a number of years to reach a dynamic equilibrium condition where the natural processes of erosion and sedimentation are in balance. The long-term goal of the Illinois River restoration projects is to reach such a state where continued excessive sedimentation is eliminated.

To assess these processes, long-term monitoring is needed. The CREP program has been collecting sediment data at selected watersheds to supplement other monitoring programs. The data collection for the CREP program started in 1999 and has generated fourteen years of data. The annual sediment load data for each of the five CREP monitoring stations have been presented in chapter 2. Because of the short duration of data collection program, this data cannot yet be used to assess long-term trends. However, the short-term trends are shown in figure 6-1, where the sediment load per unit area was normalized by the runoff in inches to account for the variability of runoff from year to year. Even though the extreme wet year 2008 stands out as the year with the highest yield (for Panther and Cox Creeks), the general trend for the other stations is a gradual decrease or no trend. Again, these are short term trends and any major climatic or hydrologic variability in the coming year could change the trends, as illustrated with the influence of 2008 on Panther and Cox Creeks. As we continue the monitoring program, the trends will be more clear and reliable as the duration of the monitoring period increases.

The data were also compared with historical data collected by the USGS for small watersheds in the Illinois River basin as shown in figure 6-2. As shown in the figure, the CREP dataset is consistent with the older dataset and will be used to develop improved sediment delivery estimates for small watersheds in the Illinois River basin and improve our assessment and evaluation capability.

To assess long-term trends, data collected by the USGS and ISWS since 1980 were used to compute sediment delivery for the major tributaries to the Lower Illinois River. For the USGS data, sediment delivery from the three major tributary watersheds to the Lower Illinois River was computed for the downstream gaging stations near the outlet of the watersheds using the same methods developed by Demissie et al. (2004). The outflow of sediment from the Illinois River basin is measured at Valley City. The sediment loads and the corresponding water discharges for




Figure 6-1. Variability of sediment yield per inch of runoff for CREP monitoring stations




Figure 6-2. Comparison of sediment load from CREP monitoring stations with historical sediment data for small watersheds by the USGS

five-year increments since 1980 are shown in figure 6-3. The period 1991-1995 generally shows the highest sediment delivery to the Illinois River and the highest outflow from the Illinois River for the period under consideration, primarily because of the 1993 major floods. Since that period, sediment delivery from the tributaries and outflow from the Illinois River have generally been decreasing. If these trends continue into the future, there would be significant reduction in sediment delivery to the Illinois River.

Similar trends are also observed from the analyses of sediment data collected by the ISWS for the Benchmark Sediment Monitoring Program for Illinois Streams. The Benchmark Sediment Monitoring Program has been collecting weekly sediment data at selected monitoring stations throughout the state since 1980 (Allgire and Demissie, 1995). The data collected over that last 30 years have been processed and analyzed to observe trends in sediment concentrations and loads. Figures 6-4 to 6-6 show the trend in sediment load since 1980 for the Spoon River at Long Mills, LaMoine River at Ripley, and Sangamon River at Monticello, respectively. All three stations show a decreasing trend since 1980 even though the 2009 and 2010 annual loads are higher than the mean annual loads.

Nutrient Loadings

To assess long-term trends in nutrient loadings as conservation practices are implemented, the state has been collecting nutrient data at the five CREP monitoring stations where sediment data have been collected since 1999. Even though there are some low and high nutrient load years, the dataset is not long enough to assess long-term trends in nutrient loading. However, the short-term trends based on the data collected so far are shown in figures 6-7 and 6-8 for nitrate-N and total phosphorous yields per inch of runoff respectively. The nutrient yield values were divided by the inches of runoff to partly remove the effect of the variability of runoff from year to year. As shown in figure 6-7, the nitrate-N yields do not show any significant trend except for the jumps in yield from 2000 to 2001 and from 2012 to 2013 for stations 201 and 202 and a gradual decline since 2006 for all stations. Figure 6-8 shows no significant trend for total phosphorous over the whole monitoring period except for the jump in yield in 2000 and 2008 for stations 201 and 202 and a significant drop for all the stations in 2012 due to the drought and jump in 2013 for stations 301, 302, and 303.

Long-term data collected by the Illinois EPA as part of their Ambient Water Quality Monitoring Network can, however, provide a fair indication of the general long-term trend in nutrient delivery to the Illinois River. Figure 6-9 shows annual nitrate-N yields in tons per square mile from the three major tributaries of the Lower Illinois River (Spoon, Sangamon, and LaMoine Rivers). Nitrate-N represents about 70 percent of the total nitrogen load in most of Illinois' agricultural watershed, and thus is a good surrogate for total nitrogen load. As can be seen in the figure, the nitrate yields can range from almost zero during a drought year like 1989 to a high of about 11 tons per square mile during a major wet period like the 1993 flood year. Therefore, climatic factors do play a major role in nutrient transport and delivery. The most important observation that can be made for the figure is the slow decreasing trend of nitrate-N yield from the major tributary watersheds. Even though it is very difficult to measure how much impact the CREP program might have had, it is obvious that conservation practices in these watersheds, where most of the CREP lands are located, are making a difference in nitrogen delivery to the Illinois River.




Figure 6-3. Sediment delivery from the three major tributary watersheds to the Illinois River and sediment outflow from the Illinois River at Valley City




Figure 6-4. Trends in sediment load at Spoon River at London Mills (after Crowder et al., 2008)




Figure 6-5. Trends in sediment load at LaMoine River at Ripley, IL (after Crowder et al., 2008)




Figure 6-6. Trends in sediment load at Sangamon River at Monticello, IL (after Crowder et al., 2008)




Figure 6-7. Variability of nitrate-N yield per inch of runoff for CREP monitoring stations




Figure 6-8. Variability of total phosphorous yield per inch of runoff for CREP monitoring stations

Figure 6-10 shows the total phosphorous yield from the same three tributary watersheds discussed in the previous figure. Annual phosphorous delivery ranges from a low of almost zero during the drought year 1989 to a high of almost one ton per square mile for the extreme wet year of 1993. The data also show how dependant phosphorous delivery is on climatic variability. Similar to the trends to the nitrate delivery, there is a slow but gradual decreasing trend in phosphorous yield from the Spoon and LaMoine Rivers, while there is a gradual increase from the Sangamon River.

The trends in nutrient loads from the major tributaries are reflected in nutrients transported by the Illinois River. Analyses of the data from the two downstream monitoring stations, Havana and Valley City, are shown in figure 6-11 for nitrate-N and total phosphorous, respectively. In general, the trend is a gradual decrease to no increase. These observations are extremely important as to nutrient delivery from Illinois streams to the Mississippi River and eventually to the Gulf of Mexico. Illinois had been identified as one of the major sources of nutrients to the Gulf of Mexico, and the fact that nutrient delivery from Illinois has not increased and is gradually decreasing is good news not only to Illinois but to the Gulf of Mexico, too.




Figure 6-9. Annual nitrate-N loads for the three major tributary watersheds to the Lower Illinois River




Figure 6-10. Annual total phosphorous loads for the three major tributary watersheds to the Lower Illinois River




Figure 6-11. Nitrate-N and total phosphorous loads along the Lower Illinois River

7. Summary and Conclusions

The Illinois River Conservation Reserve Enhancement Program (CREP) was initiated as a joint federal/state program with the goal of improving water quality and wildlife habitat in the Illinois River basin. Based on numerous research and long-term data, the two main causes of water quality and habitat degradations in the Illinois River were known to be related to sedimentation and nutrient loads. Based on this understanding, the two main objectives of the Illinois River CREP were to reduce the amount of silt and sediment entering the main stem of the Illinois River by 20 percent; and to reduce the amount of phosphorous and nitrogen loadings to the Illinois River by 10 percent. To assess the progress of the program towards meeting the two goals, the Illinois Department of Natural Resources (IDNR) and the Illinois State Water Survey (ISWS) are developing a scientific process for evaluating the effectiveness of the program. The process includes data collection, modeling, and evaluation.

The monitoring and data collection component consist of a watershed monitoring program to monitor sediment and nutrient for selected watersheds within the Illinois River basin and also to collect and analyze land use data throughout the river basin. Historically, there are a limited number of sediment and nutrient monitoring stations within the Illinois River basin, and most of the available records are of short duration. To fill the data gap and to generate reliable data for small watersheds, the Illinois Department of Natural Resources funded the Illinois State Water Survey to initiate a monitoring program that will collect precipitation, hydrologic, sediment, and nutrient data for selected small watersheds in the Illinois River basin that will assist in making a more accurate assessment of sediment and nutrient delivery to the Illinois River. Five small watersheds located within the Spoon and Sangamon River watersheds were selected for intensively monitoring sediment and nutrient within the Illinois River basin. The Spoon River watershed generates the highest sediment per unit area in the Illinois River basin, while the Sangamon River watershed is the largest tributary watershed to the Illinois River and delivers the largest total amount of sediment to the Illinois River.

As outlined in the Illinois River Basin Restoration Plan, the alternative of no-action in the Illinois River watershed would have resulted in increased sediment delivery to the Illinois River and habitats and ecosystem would continue to degrade. However, analysis of the available long term data from different sources and the most recent data from the CREP monitoring program, indicate that sediment and nutrient loads from the tributary watersheds are gradually decreasing or stabilizing as a result of implementation of conservation practices in the watershed. With the knowledge that reduction in sediment delivery from large watersheds takes time to move through the system, the indication of stabilized sediment delivery shows progress is being made in restoring the Illinois River watershed. If the present trends continue for the next 10 to 15 years, sediment and nutrient delivery to the Illinois River will be significantly reduced, and lead to improved ecosystem in the river and tributary watersheds in the long-term.

8. References

- Allgire, R.L., and M. Demissie. 1995. *Benchmark Sediment Monitoring Program for Illinois Streams: Program Summary and Site Descriptions*. Illinois State Water Survey Circular 181.
- Bicknell, B.R., J. C. Imhoff, J. L. Kittle, Jr., T. H. Jobes, and A. S. Donigian, Jr., 2001. Hydrological Simulation Program – FORTRAN (HSPF), User's Manual for version 12.0, USEPA, Athens, GA, 30605.
- Demissie, M., L. Keefer, and J. Slowikowski, A. Russell, T. Snider, and K. Stevenson. 2001. Sediment and Nutrient Monitoring at Selected Watersheds within the Illinois River Watershed for Evaluating the Effectiveness of the Illinois River Conservation Reserve Enhancement Program (CREP). Illinois State Water Survey Contract Report 2001-12.
- Demissie, M., R. Xia, L. Keefer, and N. Bhowmik. 2004. The Sediment Budget of the Illinois River. Illinois State Water Survey Contract Report 2004-13.
- Demissie, M., and N.G. Bhowmik. 1986. *Peoria Lake Sediment Investigation*. Illinois State Water Survey Contract Report 371.
- Demissie, M. R. Xia, L. Keefer, and N. Bhowmik. 2004. *The Sediment Budget of the Illinois River*. Illinois State Water Survey Contract Report 2004-13.
- Knapp, H.V. 2005. Analysis of Streamflow Trends in the Upper Midwest Using Long-Term Flow Records. *Proceedings of the 2005 Water and Environmental Resources Congress*. American Society of Civil Engineers, May 15-19, 2005, Anchorage, AK.
- Luman, D.E., 2008, Assessment & Potential of the 2007 USDA-NASS Cropland Data Layer for Statewide Annual Land Cover Applications, unpublished report of investigations including five appendices, Illinois State Geological Survey, 17 pp.
<http://crystal.igs.uiuc.edu/nsdihome/webdocs/landcover/nass07.html>.
- Luman, D. and T. Weicherding. *Land Cover of Illinois: 1999-2000*. 1999 January 1999 [cited 2003 November 1, 2003]; Available from: <http://www.agr.state.il.us/gis/index.html>.
- Ramamurthy, G.S., K.P. Singh, and M.L. Terstriep. 1989. *Increased Durations of High Flows Along the Illinois and Mississippi Rivers: Trends and Agricultural Impacts (Final Report)*. Illinois State Water Survey Contract Report 478.
- Short, M.B. 1999. *Baseline Loadings of Nitrogen, Phosphorus and Sediments from the Illinois Watersheds*. Illinois Environmental Protection Agency, IEPA/BOW/99-020, Springfield, IL.
- Singh, J., H. V. Knapp, Y. Lian, and M. Demissie. 2003. The Illinois River Basin Hydrologic Model. *Proceedings, 2003 Governor's Conference on the Management of the Illinois River System*. Oct. 2003, Peoria, IL, pp.73-84.
- State of Illinois, Department of Natural Resources, 2002. Illinois Conservation Reserve Enhancement Program (CREP) 2002. Annual Report for Reporting Period October 2000 through September 2001, Springfield, IL.
- U.S. Army Corps of Engineers. 2006. *(Draft) Illinois River Basin Restoration Comprehensive Plan with Integrated Environmental Assessment*. Rock Island District, Rock Island, IL.

APPENDIX D

**Sediment and Nutrient Monitoring at Selected
Watersheds within the Kaskaskia River
Watershed for Evaluating the Effectiveness of
the Illinois River Conservation Reserve
Enhancement Program (CREP)**

by
Illinois State Water Survey
Prairie Research Institute, University of Illinois
Champaign, Illinois

Prepared for the
Office of Resource Conservation,
Illinois Department of Natural Resources
Springfield, Illinois

December 2014

This report was printed on recycled and recyclable papers.

Contents

	Page
1. Introduction.....	1
Acknowledgments.....	1
2. Watershed Characteristics and Background	3
Hydrology	3
Geology.....	3
3. Monitoring and Data Collection	9
Monitoring Stations	10
Stream Stage and Flow	15
Sediment and Nutrient Data.....	16
Sediment Data.....	16
Nutrient Data.....	16
4. Land Use Practices.....	25
Land Cover.....	25
Conservation Practices.....	34
5. Summary and Conclusions	39
6. References.....	41

List of Tables

	Page
2-1 Kaskaskia tributary watersheds and drainage areas	6
3-1 Sediment and Nutrient Monitoring Stations Established for the Kaskaskia River CREP	10
3-2 Summary Statistics for Water Year 2000–2014	24

List of Figures

	Page
2-1	Kaskaskia River Basin, sub-basins, and major tributary watersheds..... 5
2-2	Physiographic regions and loess thicknesses in Kaskaskia Basin 7
3-1	General location of monitoring stations in the Kaskaskia River watershed 11
3-2	Detailed location of monitoring stations in Lost Creek (402) watershed 12
3-3	Detailed location of monitoring stations in North Fork Kaskaskia River (403) watershed 12
3-4	Detailed location of monitoring stations in Hurricane Creek (404) and East Fork Shoal Creek watersheds 13
3-5	Streamgage monitoring stations in Kaskaskia River Basin: a) Lost Creek, b) North Fork Kaskaskia River, c) Hurricane Creek, and d) East Fork Shoal Creek..... 14
3-6	Raingage stations in Kaskaskia River Basin: a) Witt, Illinois (43) and Shattuc, Illinois (44) 15
3-7	Suspended sediment concentrations and water stage at Lost Creek (402) and North Fork Kaskaskia River (403) for Water Year 2014..... 18
3-8	Suspended sediment concentrations and water stage at Hurricane Creek (404) and East Fork Shoal Creek (405) for Water Year 2014..... 19
3-9	Nitrogen concentrations and water stage at Lost Creek (402) and North Fork Kaskaskia River (403) for Water Year 2014..... 20
3-10	Nitrogen concentrations and water stage at Hurricane Creek (404) and East Fork Shoal Creek (405) for Water Year 2014..... 21
3-11	Phosphorus concentrations and water stage at Lost Creek (402) and North Fork Kaskaskia River (403) for Water Year 2014..... 22
3-12	Phosphorus concentrations and water stage at Hurricane Creek (404) and East Fork Shoal Creek (405) for Water Year 2014..... 23
4-1	Types of land cover in Kaskaskia River Basin 26
4-2	Percent watershed area of types of land cover in Kaskaskia River Basin 27
4-3	Conservation Reserve Program (CRP) in Kaskaskia Basin..... 28
4-4	Watershed being monitored for hydrology, sediment, and nutrients..... 29
4-5	Kaskaskia River Basin Watersheds: Percent change in generalized NASS land-use from 2007-2013 30
4-6	Lost Creek watershed from ISWS Station 402: Generalized NASS Cropland Data Layer Acreage Totals, 2007-2013 31

List of Figures (concluded)

	Page
4-7	North Fork Kaskaskia River watershed from ISWS Station 403: Generalized NASS Cropland Data Layer Acreage Totals, 2007-2013..... 32
4-8	Hurricane Creek watershed from ISWS Station 404: Generalized NASS Cropland Data Layer Acreage Totals, 2007-2013 32
4-9	East Fork Shoal Creek watershed from ISWS Station 402: Generalized NASS Cropland Data Layer Acreage Totals, 2007-2013 33
4-10	State CREP contract locations (IDNR, 2014)..... 35
4-11	Cumulative acres of conservation practices installed in Lost Creek watershed at monitoring station ISWS #402 from 1999-2013 36
4-12	Cumulative acres of conservation practices installed in North Fork Kaskaskia River watershed at monitoring station ISWS #403 from 1999-2013 36
4-13	Cumulative acres of conservation practices installed in Hurricane Creek watershed at monitoring station ISWS #404 from 1999-2013 37
4-14	Cumulative acres of conservation practices installed in East Fork Shoal Creek watershed at monitoring station ISWS #405 from 1999-2013 37

Monitoring and Evaluation of Sediment and Nutrient Delivery to the Illinois River: Illinois River Conservation Reserve Enhancement Program (CREP)

by
Illinois State Water Survey
Prairie Research Institute, University of Illinois

1. Introduction

The Conservation Reserve Enhancement Program (CREP) was initiated as a joint federal/state program with the goal of improving water quality and wildlife habitat in the Illinois and Kaskaskia River basins. Based on numerous research and long-term data, the two main causes of water quality and habitat degradations in the rivers of Illinois were known to be related to sedimentation and nutrient loads. Based on this understanding, the two main objectives of the Illinois CREP were stated as follows:

- 1) Reduce the amount of silt and sediment entering the main stem of the Illinois and Kaskaskia River by 20 percent.
- 2) Reduce the amount of phosphorous and nitrogen loadings to the Illinois and Kaskaskia River by 10 percent.

To assess the progress of the program towards meeting the two goals, the Illinois Department of Natural Resources (IDNR) and the Illinois State Water Survey (ISWS) are developing a scientific process for evaluating the effectiveness of the program. The process includes data collection, modeling, and evaluation. Progress made so far in each of these efforts is presented in this report.

Acknowledgments

The work upon which this report is based is supported by funds provided by the Office of Resource Conservation, Illinois Department of Natural Resources, under the guidance of Luke Garver, IDNR CREP Program Coordinator. The project is also supported as part of Laura Keefer's regular duties at the Illinois State Water Survey under the guidance of Mike Demissie, Director, Illinois State Water Survey and ISWS Principal Investigator for the CREP studies.

Several Illinois State Water Survey staff worked diligently to meet project objectives and their tireless dedication is much appreciated. Erin Bauer developed and implemented the data collection computer programming and protocols for the field instrumentation. John Beardsley and Jim Osborne designed, fabricated, and installed the instrument shelters. Erin Bauer designed, developed and manages the project databases, processes data, performs quality control, and produced the tables and figures for this report. Yao Hu, University of Illinois Ph.D. candidate, wrote the MatLab code for the project data processing. Lara Seek and John Beardsley are responsible for the field data collection efforts that amount to many long, wet, and tiring days – a herculean effort given all the rain this past year. Joyce Wyse performs data entry as well as

track and organize project records. Erin Bauer compiled, investigated and analyzed the land cover data (Illinois and Kaskaskia Basins). Phil Graff, ISWS, and Lisa Beja, IDNR, compiled the CREP contract conservation practices database (Illinois and Kaskaskia Basins). Laura Keefer is responsible for the overall investigation, implementation, management and analyses of the ISWS Kaskaskia monitoring activities. Leon Hinz, Illinois Natural History Survey (INHS), collaborated in the ISWS/INHS co-location site selection of intensive monitoring stations and his assistance was most appreciated.

2. Watershed Characteristics

The Kaskaskia River watershed has a drainage area of 5,810 mi², is generally located in the southwest region of the State of Illinois, and occupies all or portions of 15 counties (Figure 1). The headwaters begin in Champaign and Piatt Counties in east-central Illinois and flows in a southwesterly direction to join the Mississippi River in Randolph County. Table 1 lists the tributary watersheds and associated drainage areas. Figure 1 illustrates approximately 22 tributary watersheds in the basin that range in drainage area from 53 to 917 mi². The two largest tributary watersheds are Shoal Creek (917 mi²) and Silver Creek (480 mi²) and together occupy nearly 25 percent of the Kaskaskia River watershed drainage area. In general, the Kaskaskia River watershed is divided into four sub-watersheds (Upper, Middle, Lower, and Shoal Creek) that are associated with the outlets at the two main reservoirs, Lake Shelbyville and Carlyle Reservoir, and confluence with the Mississippi River. The Shoal Creek tributary watershed is distinguished due to its large drainage area. See (Illinois Department of Natural Resources 2000) for further information.

Hydrology

Knapp and others (2012) describe the Kaskaskia River as one of the more highly managed rivers in Illinois. The streamflow on the main stem of the Kaskaskia River is controlled by two federal reservoirs (Shelbyville and Carlyle Reservoirs) and the navigation pools in the lower reaches of the river are maintained by a lock and dam. Water is withdrawn for industry and public water supplies from several reservoirs constructed throughout the watershed. Other inflows come from effluent discharges throughout the drainage system by municipal systems and industries, as well as power plant cooling water returns. A detailed water supply assessment of the Kaskaskia River watershed can be found in (Knapp, Roadcap et al. 2012).

Geology

The surficial geology plays a role in the types of land cover in the Kaskaskia River watershed. Figure 5 illustrates the boundaries of the physiographic regions, loess (windblown silt) thicknesses, and shaded relief for the Kaskaskia River watershed. The watershed is predominantly in the Bloomington Ridged Plain and Springfield Plain of the Till Plains Section.

The Upper sub-watershed is entirely in the Bloomington Ridged Plain and characterized by low, broad ridges with intervening wide stretches of relatively flat or gently undulatory ground (Leighton, Ekblaw et al. (1948). These alternating ridges with flat ground are indicative of the most recent glacial period, referred to as the Wisconsin. Therefore, the drainage system is more recent than the Springfield Plain which is older and more developed.

The Middle, Shoal Creek, and most of the Lower sub-watersheds are in the Springfield Plain which is part of the Illinoian glacial drift period that occurred before the Wisconsin. The Illinoian is characteristically flat with low and broad ridges (moraines) but some areas in the Kaskaskia watershed have ridges and hills with irregular assemblages of gravel with small intervening plains. The drainage system is characterized by major rivers in low gradient and broad terraced valleys and tributaries in wide v-shaped valleys with headwaters originating from the low gradient, broad shallow valleys of the till plains. Basically, the Springfield Plain occupies the older Illinoian glacial drift with older drainage development, whereas the

Bloomington Ridge Plain occupies the Wisconsin, which overlies the Illinoian, and is flat with sequences of ridges and initial stages of drainage.

Another geologic characteristic that controls drainage development and is a factor in erosion is the thickness of the windblown silt (loess) that overlies the glacial drift, similar to frosting on a layer cake (Illinoian and Wisconsin glacial drift), somewhat smoothing out imperfections on the surface. As seen in Figure 5, the loess in the Upper sub-watershed is between 0-5 feet thick and lies in the ridged and wide flat valleys of the Bloomington Ridged Plain. Most of the Middle sub-watershed and upper reaches of the Shoal Creek sub-watershed the loess is 0-5 feet thick and lies in the more developed drainage landscape of the Springfield Plain. The lower reaches of Shoal Creek and most of the Lower sub-watersheds have thicknesses that can range from 5 to greater than 20 feet proceeding from east to west toward the Mississippi River. However, many of the stream valleys in these areas do not have loess present and is considered to have been eroded. Areas with thick loess are considered prone to erosion under steep conditions which can result in unstable stream channels. A more extensive discussion on the geology and surficial materials in the Kaskaskia River watershed can be found in (Illinois Department of Natural Resources 2000).

In summary, the four sub-watersheds of the Kaskaskia River watershed are fairly distinct from each other based on geology and land cover features. These features have an influence on water quality, erosion, and aquatic habitat. Agriculture production is dominant in the Upper sub-watershed due to the consistent, relatively flat and wide valleys between gentle ridges, as well as the highly productive soil developed in the loess cap. Large areas dominated by highly productive soil and agriculture tend to have elevated nutrient levels in the stream system. The Middle and upper-Shoal sub-watersheds have a mix of agriculture and woodlands/grasses, where the agriculture is in the flatter uplands and woodlands in the deeper valleys. Nutrients may be slightly more elevated in the drainage system but some erosion issues may play a factor in the valleys. The lower-Shoal and Lower sub-watersheds are similar in land use to the Middle sub-watershed, slightly more agriculture but the loess thicknesses in combination with higher relief result in erosion being more of an issue in these areas.




Figure 2-1. Kaskaskia River Basin, sub-basins, and major tributary watersheds

Table 2-1. Kaskaskia tributary watersheds and drainage areas

<i>Tributary Name</i>	<i>Drainage Area</i>	
	<i>(acres)</i>	<i>(mi²)</i>
Ash Creek	89,610	140
Beck Creek	130,771	204
Crooked Creek	224,663	351
East Fork Kaskaskia River	132,477	207
Elkhorn Creek	56,760	89
Hickory Creek	92,224	144
Hoffman Creek	67,428	105
Horse Creek	60,175	94
Hurricane Creek	128,822	201
Johnathan Creek	36,896	58
Kaskaskia-L. Shelbyville	122,705	192
Kaskaskia Ditch	103,474	162
Kaskaskia River	658,183	1,028
Lake Fork	109,537	171
Little Crooked Creek	73,254	114
Mud Creek	87,207	136
Plum Creek	57,399	90
Richland Creek	213,431	333
Robinson Creek	79,112	124
Shoal Creek	586,584	917
Silver Creek	307,171	480
Sugar Creek	112,775	176
West Okaw River	154,219	241
Whitley Creek	33,687	53
<i>Total</i>	<i>3,718,563</i>	<i>5,810</i>




Figure 2-5. Physiographic regions and loess thicknesses in Kaskaskia Basin

3. Monitoring and Data Collection

The monitoring and data collection component consists of a watershed monitoring program to monitor sediment and nutrient for selected sub-watersheds within the Kaskaskia River basin and also to collect and analyze land use data throughout the river basin. Currently available data is insufficient to monitor long-term trends especially in small watersheds where changes can be observed and quantified more easily than in larger watersheds. To fill the data gap and to generate reliable data for small watersheds, the Illinois Department of Natural Resources funded the Illinois State Water Survey to initiate a monitoring program that will collect precipitation, hydrologic, sediment, and nutrient data for selected small watersheds in the Kaskaskia River basin that will assist in making a more accurate assessment of sediment and nutrient delivery.

The monitoring strategy for the project was to select small Kaskaskia River tributary watersheds to establish an intensive monitoring program to detect any changes in sediment and nutrient transport characteristics that could be attributed to changes in land use or other factors. The project is designed to measure the cumulative impact within the watershed on sediment and nutrient yield and is not designed to measure the impact of specific BMPs on water quality or sediment yield. Several factors were evaluated to determine the final locations of the intensive monitoring sites, such as artificial inflow and outflow of water due to water supply, industrial, and recreation needs, geology, land use, currently available water quality data for more prescriptive monitoring plans, areas likely to have appreciable CREP sign-ups, and co-location with other physical, biological, and water quality program stations. Co-locating and/or supplementing monitoring stations with other water quality and aquatic sampling stations in the watershed is an integrated approach that contributes to understanding the mechanisms that link hydrologic, sediment, nutrient, biological, and physical information for application in other watersheds in Illinois.

Due to the highly managed nature of the Kaskaskia River watershed hydrology, this project assessed locations of water inflows and outflows that could mask monitoring results by affecting the normal balance of the sediment and nutrient loading character. For example, the streamflow in the main stem of the Kaskaskia River and several tributaries are significantly controlled by the periodic releases from reservoirs. Also, the water from those releases are more of a reflection of the water quality from lake processes rather than the transport of water and nutrients from the upper portions of the drainage system. Also assessed were locations of waste water treatment plant (WWTP) effluent, NPDES discharges, and other smaller reservoirs in the tributary watersheds. This project capitalized on a recently completed water supply assessment for the Kaskaskia River watershed, which assembled existing water availability and supply information mentioned above by Knapp, Roadcap et al. (2012).

To effectively monitor any changes in sediment and nutrient loading due to CREP, small-scale intensive monitoring in several places improves the ability to monitor changes over time. Ideally, these small-scale study watersheds should be in areas that will have the highest likelihood of CREP sign-ups. The ISWS contacted several local stakeholder groups, county Soil and Water Conservation Districts and CREP program staff to estimate areas likely to have appreciable CREP sign-ups within the Kaskaskia River watershed. This assessment period overlapped with the 2012 drought which appeared to have appreciably reduced CREP sign-ups for the first year of the project. Consequently, in collaboration with Illinois Natural History Survey (INHS) investigators, an analysis was made based on land cover, geology, hydrology,

biology and conservation reserve programs (CRP) already in the watershed. This allowed for comparing and contrasting watershed land uses with physical character that allowed for selection of watersheds estimated to be likely and unlikely for CREP signups. Four watersheds were then selected to represent combinations of physical watershed character and land cover.

Monitoring Stations

The four small watersheds selected for intensively monitoring sediment and nutrient within the Kaskaskia River basin are located within the Crooked Creek, North Fork Kaskaskia River, Hurricane Creek and Shoal Creek watersheds. In addition, two continuous recording raingages were established near the monitored watersheds. The general locations of the watersheds, monitoring stations and raingages are shown in figure 3-1 and more detailed station maps are shown in figures 3-2 through 3-4. Information about the stations is provided in table 3-1. Lost Creek (402) is a tributary of Crooked Creek which, in turn is a direct tributary of the Kaskaskia River with its confluence downstream of Carlyle Reservoir. The Carlyle Reservoir is a U.S. Army Corps of Engineers impoundment on the Kaskaskia River. The North Fork Kaskaskia River (403) and Hurricane Creek (404) are a direct tributaries of the Kaskaskia River and discharged directly into the upstream end of Carlyle Reservoir. East Fork Shoal Creek (405) is a tributary of Shoal Creek, the largest direct tributary of the Kaskaskia River, with its confluence downstream of Carlyle Reservoir. The type of data collected and the data collection methods have been presented in detail in the first progress report for the CREP monitoring program (Demissie et al., 2001) and in the Quality Assurance Project Plan (QAPP) given in Appendix A. The data collected at each of the monitoring stations follows these protocols.

Table 3-1. Sediment and Nutrient Monitoring Stations and Raingages Established for the Kaskaskia River CREP

<i>Station ID</i>	<i>Name</i>	<i>Drainage area</i>	<i>Watershed</i>
402	Lost Creek	38.0 sq mi (24,320 acres)	Crooked Creek
403	North Fork Kaskaskia River	35.5 sq mi (22,701 acres)	North Fork Kaskaskia River
404	Hurricane Creek	27.7 sq mi (17,753 acres)	Hurricane Creek
405	East Fork Shoal Creek	30.9 sq mi (19,820 acres)	Shoal Creek
43	Witt, IL	--	East Fork Shoal & Hurricane Creeks
44	Shattuc, IL	--	Lost Creek




Figure 3-1. General location of monitoring stations in the Kaskaskia River watershed




Figure 3-2. Detailed location of monitoring stations in Lost Creek (402) watershed




Figure 3-3. Detailed location of monitoring station in North Fork Kaskaskia River (403) watershed




Figure 3-4. Detailed location of monitoring stations in Hurricane (404) and East Fork Shoal Creek (405) watersheds

Each of the four monitoring stations are instrumented with a Campbell Scientific CR850 datalogger, CS476 radar water level sensor, ISCO automatic water sampler, cell modem, antenna, solar panel, and batteries. All instruments, except the ISCO sampler, are housed in a stainless steel shelter to protect them from weather and vandalism. The ISCO sampler is housed in a modified 55-gallon steel drum with a hinged lid for access. The two raingages are instrumented with a modified Belfort weighing-bucket raingage, Campbell Scientific CR200 datalogger, cell modem, antenna, solar panel, and battery. The shelter and instrument configurations of the four streamgage monitoring stations are shown in Figure 3-5 and raingage stations in Figure 3-6. All data is retrieved from the station dataloggers via cell modem every hour to ISWS computer databases.



Figure 3-5. Streamgage monitoring stations in Kaskaskia River Basin: a) Lost Creek, b) North Fork Kaskaskia River, c) Hurricane Creek, and d) East Fork Shoal Creek




Figure 3-6. Raingaging stations in Kaskaskia River Basin: a) Witt, Illinois (43) and b) Shattuc, Illinois (44)

Stream Stage and Flow

The “stage” of a stream is the measurement of the water surface of a stream from an arbitrary datum. The stage record is collected continuously and makes it possible to determine the volume of water carried by a stream past a streamgaging station. Through the application of a stage discharge rating curve, the continuous stage is converted to streamflow. Streamflow data are generated from the 15-minute stage record at a streamgaging station. The stage data are converted to discharge (streamflow) by applying a stage-discharge calibration curve. The calibration is developed by taking several detailed field measurements of the streamflow at known stages.

Methods used in this study for determining stream discharge follow established USGS procedures as outlined by Rantz (1982a, 1982b). Stream discharge is determined by measuring the mean velocity along a stream cross section. Each vertical represents the velocity of a flow area (substation), which is defined as the sum of half the distance between verticals by the water depth at the vertical. At each vertical the velocity is sampled at 20 and 80 percent of the total depth (for total depths ≥ 2.5 feet) or at 60 percent of the total depth (for total depths < 2.5 feet). The average of the 20 and 80 measurements or the single 60 percent measurement is assumed to be the mean velocity for that subsection. Each subsection discharge is calculated by multiplying the average velocity by the flow area, and then the sum of all the subsections equals the total discharge of the stream cross section. Every discharge is then plotted against the corresponding stage at which the discharge measurement occurred. After sufficient measurements have been collected, a curve is developed to express the relationship between stage and discharge. Using this stage-discharge curve, the stage data files are then converted to discharge. The discharge data can then be used to develop nutrient and sediment load data.

All data are compiled in to what is referred to as “water years”, which begins on October 1st and ends September 30th of the following year. The year is associated with the close of the period. For example, water year 2014 (WY2014) begins October 1, 2013 and ends September 30, 2014.

The process of collecting a sufficient number of streamflow measurements to adequately develop a stage-discharge calibration takes time. This usually takes 1-2 water years into a monitoring study. Therefore, stream discharge values are not available at this time, as well as sediment and nutrient load calculations. It is anticipated that calibration curves will be sufficient for producing preliminary data at the next annual progress report.

Sediment and Nutrient Data

Sediment Data

Suspended sediment samples are collected either manually or by ISCO automated pump sampler. The suspended sediment sampling methods used in this study followed established USGS procedures as outlined by Edwards and Glysson (1999) and FISP (1952). The manual sampling method used depth-integrating samplers for all but the shallowest conditions. The second method used to collect suspended sediment samples was the ISCO automated pump sampler. The programming of the CR850 datalogger controls the ISCO sampling schedule. This program allows automated sampling during high-flow events and is triggered by changes in stage over time. Manual suspended sediment samples were taken at all four stations during weekly station visits and during storms when possible.

Suspended sediment concentration (SSC) data and stream stage for all stations are shown in figures 3-7 and 3-8 for WY2014. Summary statistics for SSC samples can be found in table 3-2. As can be seen in the figures, suspended sediment concentrations are highly variable throughout a year depending on the climatic conditions and location of the stations in the watershed. The distance between monitoring stations ranges from 10 to 45 miles and subject to rainfall and storm variability and tracking through the region. It is also evident that sediment concentrations are the highest during storm events resulting in the transport of most of the sediment during storm events. Therefore, it is extremely important that samples are collected frequently during storm events to accurately measure sediment loads at monitoring stations. Approximately 463 SSC samples were collected at all stations. The highest maximum SSC occurred at Lost Creek (402) at 4616 mg/L and lowest at North Fork Kaskaskia (403) with 2487 mg/L. All stations had minimum SSC below 10 mg/L.

Nutrient Data

The nutrient data are organized into two groups: nitrogen species and phosphorous species. The nitrogen species include nitrate-nitrogen (NO₃-N), ammonium-nitrogen (NH₄-N), and total Kjeldahl nitrogen (TKN). The phosphorous species include total phosphorous (TP), total dissolved phosphorous (TDP), and orthophosphate (P-ortho). Approximately 1,023 samples have been collected for nitrogen and 888 for phosphorus. Nitrogen and phosphorus sample results with stream stage are shown in figures 3-9 to 3-10 and 3-11 to 3-12, respectively. A summary statistics for all stations showing the sample count, mean, median, minimum, maximum, 25th percentile, and 75th percentile are given in table 3-2.

Data for the nitrogen species at all four monitoring stations show that the dominant form of nitrogen transported by the streams is total Kjeldahl nitrogen (TKN). During storm events, the concentration of TKN rises significantly, exceeding the nitrate-N concentration ($\text{NO}_3\text{-N}$). TKN is highly correlated to suspended sediment concentrations. Ammonium-nitrogen ($\text{NH}_4\text{-N}$) concentrations are low at all stations except East Fork Shoal Creek (405) where concentrations were exceptionally high during the deep winter months of January and February 2014. TKN maximum concentrations were at Lost Creek (402) and East Fork Shoal (405).

As can be seen in figures 3-11 and 3-12 phosphorous species at all monitoring stations show that most of the phosphorous load is transported during storm events. Total phosphorous (t-P) concentrations are the highest during storm events and relatively low most of the time. This is very similar to that shown by sediment and thus implies high correlations between sediment and phosphorous concentrations and loads. Ortho-phosphate (P-ortho) and total dissolved phosphorus (TDP) concentrations also increase during storm events, not as strongly as TKN, and does stay slightly elevated when stream stage is elevated for long periods of time as seen, for example, at Lost Creek (402) during April-July 2014.





Figure 3-7. Suspended sediment concentrations and water stage at Lost Creek (402) and North Fork Kaskaskia River (403) for Water Year 2014





Figure 3-8. Suspended sediment concentrations and water stage at Hurricane Creek (404) and East Fork Shoal Creek (405) for Water Year 2014





Figure 3-9. Nitrogen concentrations and water stage at Lost Creek (402) and North Fork Kaskaskia River (403) for Water Year 2014





Figure 3-10. Nitrogen concentrations and water stage at Lost Creek (402) and North Fork Kaskaskia River (403) for Water Year 2014





Figure 3-11. Phosphorus concentrations and water stage at Lost Creek (402) and North Fork Kaskaskia River (403) for Water Year 2014





Figure 3-12. Phosphorus concentrations and water stage at Hurricane Creek (404) and East Fork Shoal Creek (405) for Water Year 2014

Table 3-2. Summary Statistics for Water Year 2014 (all concentrations in mg/L).

	<i>NO3-N</i>	<i>oPO4-P</i>	<i>NH4-N</i>	<i>TKN</i>	<i>t-P</i>	<i>t-P-Dissolved</i>	<i>SSC</i>
Lost Creek (402)							
Count	157	156	157	129	129	129	167
Mean	0.83	0.40	0.13	2.57	0.95	0.46	493.79
Median	0.60	0.42	0.07	2.25	0.87	0.46	234.39
Min	0.04	0.05	0.03	0.75	0.17	0.07	9.39
Max	3.82	1.20	0.93	12.15	2.56	1.41	4616.09
25 th Percentile	0.35	0.28	0.03	1.57	0.69	0.36	104.52
75 th Percentile	1.05	0.51	0.20	3.08	1.12	0.57	608.54
North Fork Kaskaskia River (403)							
Count	103	102	102	73	73	73	109
Mean	0.71	0.29	0.11	2.12	0.71	0.34	325.71
Median	0.44	0.30	0.07	1.91	0.67	0.36	205.07
Min	0.04	0.01	0.03	0.39	0.11	0.04	4.02
Max	3.28	0.88	0.48	7.21	2.01	0.93	2486.96
25 th Percentile	0.18	0.18	0.03	1.42	0.52	0.26	23.78
75 th Percentile	0.84	0.37	0.17	2.53	0.83	0.41	429.74
Hurricane Creek (404)							
Count	63	63	63	25	25	25	60
Mean	1.68	0.15	0.15	2.29	0.62	0.23	544.96
Median	1.44	0.09	0.05	1.93	0.46	0.19	71.63
Min	0.35	0.03	0.03	0.25	0.05	0.04	4.97
Max	5.51	0.72	1.06	6.66	2.05	0.73	3111.74
25 th Percentile	1.20	0.06	0.03	0.52	0.09	0.06	23.10
75 th Percentile	2.09	0.20	0.14	3.90	0.98	0.35	780.51
East Fork Shoal Creek (405)							
Count	63	63	63	25	25	25	127
Mean	1.68	0.15	0.15	2.29	0.62	0.23	653.29
Median	1.44	0.09	0.05	1.93	0.46	0.19	418.97
Min	0.35	0.03	0.03	0.25	0.05	0.04	6.56
Max	5.51	0.72	1.06	6.66	2.05	0.73	4252.03
25 th Percentile	1.20	0.06	0.03	0.52	0.09	0.06	33.00
75 th Percentile	2.09	0.20	0.14	3.90	0.98	0.35	998.17

4. Land Use Practices

Land Cover

The distribution of land cover and croplands in the basin are shown in Figure 4-1 and graphically summarized in Figure 4-2. The data is provided by the National Agricultural Statistics Service (NASS). In general, Figure 4-1 illustrates areas in agriculture production as represented by bright yellow and green colors, whereas woodlands, grassland, and wetlands in lighter greens and blues. Developed, urban types of areas are in gray. The Upper sub-watershed is dominated by agriculture production, the Middle sub-watershed is relatively equal between agriculture and all other land covers, and Shoal Creek and Lower sub-watersheds have agriculture in the flat upland areas and woodland, grassland, and wetlands predominantly in the stream valleys. Approximately 60 percent of the Kaskaskia River watershed area is in agriculture production, 30 percent in woodlands, grassland, and wetlands, 8 percent in developed (urban) areas, and 2 percent in water.

Figure 4-3 illustrates the distribution of Conservation Reserve Program (CRP) areas throughout the Kaskaskia River watershed and counties. Based on a visual inspection, a few observations can be made. A majority of CRP areas are in close proximity to streams and waterbodies. Middle and Shoal Creek sub-watersheds appear to have higher concentrations of CRP acres than the other two sub-watersheds. The Upper sub-watershed is less dense but CRP seems to be evenly distributed. The Lower sub-watershed is similar to the Upper except Clinton and Marion Counties are denser.

The USDA National Agriculture Statistics Service (NASS) land cover data has been available since 1999. In 2006 an evaluation of the usefulness of the crop data layers for annual land cover information in Illinois was undertaken by the Illinois State Geological Survey (ISGS) and NASS (Luman, 2008). Based on inherent errors associated with satellite data, irreparable mechanical problems with older multispectral imagery satellites and land cover classification methods used to interpret that imagery, new enhanced CDL protocols were established in 2007 for Illinois. Consequently, land cover misclassifications were identified prior to the new protocol, which became more apparent when evaluating the land cover in the monitored watersheds (figure 4-4): Lost Creek (402), North Fork Kaskaskia River (403), Hurricane Creek (404), and East Fork Shoal Creek (405). Therefore, any changes in land cover will be evaluated for this report beginning in 2007 through 2013 which is the most currently available NASS CDL data.

The four monitored watersheds have somewhat different ratios of land cover types. The Lost and East Fork Shoal Creek watersheds have 72 and 64 percent area in agriculture and 28 and 36 percent in non-agriculture land covers, respectively (table 4-1). One observation is North Fork Kaskaskia has over 15 percent more land in forest/shrubland than Lost Creek. Agriculture land cover is 57 and 59 percent in North Fork Kaskaskia and Hurricane Creeks, respectively, while the non-agriculture area is the inverse. Figure 4-5 illustrates the percent change in total watershed acres between 2007 and 2013 for six generalized land cover categories in each of the four monitored tributary watersheds in the Kaskaskia River Basin. Agriculture land covers were categorized into Corn, Soybeans, Double Crop with Soybeans and Other Cropland, as well as

Kaskaskia River Basin
Cropland Data - 2010




Figure 4-1. Types of land cover in Kaskaskia River Basin




Figure 4-2. Percent watershed area of types of land cover in Kaskaskia River Basin




Figure 4-3. Conservation Reserve Program (CRP) in Kaskaskia Basin




Figure 4-4. Watersheds being monitored for hydrology, sediment and nutrients.

Table 4-1. 7-year average (2007-2013) percent acres of land cover area by watershed

	<i>ISWS Station Number</i>			
	402	403	404	405
Corn	33	26	30	31
Soybeans	38	30	28	32
Other Crops	1	1	1	1
Grasslands	16	20	22	18
Forest/Shrubland	8	22	18	14
Developed, Barren, Open Space, Water, Wetlands	4	1	2	4
AGRICULTURE	72	57	59	64
NON-AGRICULTURE	28	43	41	36




Figure 4-5. Kaskaskia River Basin Watersheds: Percent Change in Generalized NASS Land-Use from 2007-2013.

summed in one category identified as Agriculture. Non-agriculture land covers were categorized into Grassland and Forest/Shrubland, and summed as Non-Agriculture. All four watersheds had a 5 percent reduction in Double Crop with Soybeans and non-agricultural land cover area (Grasslands and Forest/Shrubland) between 2007 and 2013. An increase in agricultural land cover area (Corn and Soybeans) ranged from 2 to nearly 7 percent occurred on all four watersheds. Lost, Hurricane, and East Fork Shoal Creek watersheds had marked percent increases in soybean acres and North Fork Kaskaskia watershed had an increase of corn greater than soybeans. The Hurricane Creek watershed experiences the largest decrease in non-agriculture land cover mostly occurring with losses in grasslands. Grasslands decreased on the average of 3.5 percent over all four monitored watersheds.

Figures 4-6 to 4-9 show the changes in each land cover for each year between 2007 and 2013. For this report, NASS Cropland Data Layer (CDL) categories for the monitored watersheds were combined into 6 general land cover categories: 1) corn, 2) soybean, 3) other cultivated crops, 4) grassland, 5) forest/shrubland and 6) developed, barren, open space, water and wetlands. Land cover area changes between years is represented in acres. Therefore, some watersheds may appear to have greater changes in acreage from year to year but may only represent a small percentage of the watershed depending on the total watershed acres.

Lost Creek watershed (figure 4-6) acres varied for corn, soybeans, and grasslands with corn and soybeans increasing in acres when comparing 2007 and 2013. All other land covers remained constant over the 7-year period. North Fork Kaskaskia watershed (figure 4-7) saw similar variability as Lost Creek watershed in most corn, soybeans, and grasslands acreage. Only minor increases in acres for Forest/shrubland. North Fork Kaskaskia has equal acres in Grasslands and Forest/Shrubland, as well as slightly more Soybean acres than Corn. Lost Creek has the lowest acres of Forest/Shrubland of the four watersheds. Hurricane Creek (figure 4-8) appeared to have a significant decrease in Grasslands and Developed land covers and increase in Forest/Shrubland from 2007 to 2008. . Finally, East Fork Shoal Creek (figure 4-9) exhibits the same annual variability in land cover acres between 2007 and 2013 as the other three monitored watersheds. Lost Creek and East Fork Shoal Creek watersheds have the most agriculture land covers of the four monitored watersheds, whereas North Fork Kaskaskia and Hurricane Creek watersheds are more evenly distributed of Corn, Soybean, Grassland, and Forest/Shrubland land covers. All four watersheds have extremely low acres devoted to other cultivated crops.




Figure 4-6. Lost Creek watershed from ISWS Station 402: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013.




Figure 4-7. North Fork Kaskaskia River watershed from ISWS Station 403: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013.




Figure 4-8. Hurricane Creek watershed from ISWS Station 404: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013.




Figure 4-9. East Fork Shoal Creek watershed from ISWS Station 405: Generalized NASS Cropland Data Layer Acreage Totals: 2007-2013.

Conservation Practices

There has been a significant increase in the implementation of conservation practices in Illinois in recent years with CREP making a major contribution. Figure 4-10 shows the location of approved CREP contracts for the state of Illinois as of 2013. With this type of information it will be possible to identify areas where there has been significant participation in the CREP program and where changes in sediment and nutrient delivery should be expected. The information will provide important input data to the watershed models that are being developed to evaluate the impact of CREP practices.

There are many conservation practices implemented throughout the watersheds as a result of federal and state conservation reserve programs. In order to evaluate watershed monitoring efforts, knowing when and what conservation practices are implemented in the watershed is important. Figures 4-11 to 4-14 show the cumulative acres of conservation practices installed in the four monitored watersheds from 1999 through 2013. The most installed conservation practice is new grasses/legumes in the North Fork Kaskaskia watershed. Other popular practices are existing grasses/legumes, permanent wildlife habitat, new hardwood trees, upland bird habitat buffers, and grass filter strips. North Fork Kaskaskia and Hurricane Creek watersheds have the most variety of practices installed.




Figure 4-10. State CREP contract locations (IDNR, 2014).




Figure 4-11. Cumulative acres of conservation practices installed in Lost Creek watershed at monitoring station ISWS #402 from 1999-2013.




Figure 4-12. Cumulative acres of conservation practices installed in North Fork Kaskaskia River watershed at monitoring station ISWS #403 from 1999-2013.




Figure 4-13. Cumulative acres of conservation practices installed in Hurricane Creek watershed at monitoring station ISWS #404 from 1999-2013.




Figure 4-14. Cumulative acres of conservation practices installed in East Fork Shoal Creek watershed at monitoring station ISWS #405 from 1999-2013.

5. Summary and Conclusions

The Conservation Reserve Enhancement Program (CREP) was initiated as a joint federal/state program with the goal of improving water quality and wildlife habitat in the Illinois and Kaskaskia River basins. Based on numerous research and long-term data in the Illinois River basin, the two main causes of water quality and habitat degradations in major river corridors were known to be related to sedimentation and nutrient loads. Based on this understanding, the two main objectives of the CREP were to reduce the amount of silt and sediment entering the main stem of the Illinois and Kaskaskia Rivers by 20 percent; and to reduce the amount of phosphorous and nitrogen loadings to by 10 percent. To assess the progress of the program towards meeting the two goals, the Illinois Department of Natural Resources (IDNR) and the Illinois State Water Survey (ISWS) are developing a scientific process for evaluating the effectiveness of the program.

The monitoring and data collection component consists of a sediment and nutrient watershed monitoring program for selected sub-watersheds within the Kaskaskia River basin and also to collect and analyze land use data throughout the basin. Currently available data is insufficient to monitor long-term trends especially in small watersheds where changes can be observed and quantified more easily than in larger watersheds. To fill the data gap and to generate reliable data for small watersheds, the Illinois Department of Natural Resources funded the Illinois State Water Survey to establish a monitoring program to collect precipitation, hydrologic, sediment, nutrient and land cover data for selected small watersheds in the Kaskaskia River basin that will assist in making a more accurate assessment of sediment and nutrient delivery.

The four small watersheds selected for intensively monitoring sediment and nutrient in the Kaskaskia River basin are located within the Crooked Creek, North Fork Kaskaskia River, Hurricane Creek and Shoal Creek watersheds. In addition, two continuous recording raingages were established near the monitored watersheds. Lost Creek (402) is a tributary of Crooked Creek which, in turn is a direct tributary of the Kaskaskia River with its confluence downstream of Carlyle Reservoir. The Carlyle Reservoir is a U.S. Army Corps of Engineers impoundment on the Kaskaskia River. The North Fork Kaskaskia River (403) and Hurricane Creek (404) are direct tributaries of the Kaskaskia River and discharge directly into the upstream end of Carlyle Reservoir. East Fork Shoal Creek (405) is a tributary of Shoal Creek, the largest direct tributary of the Kaskaskia River, with its confluence downstream of Carlyle Reservoir.

After assessing and evaluating many physical, geological, biological, land cover and CRP program data and information, as well as impacts of the 2012 drought, four intensive monitoring stations were selected and the sediment and nutrient monitoring network was established for the 2014 water year. The year started in one of the coldest winters recorded in the region for some time. This was followed by a particularly wet spring and summer. Nitrogen and phosphorus species concentrations more associated with particulate forms (TKN, t-P) appear to be higher than concentrations of the dissolved forms ($\text{NO}_3\text{-N}$ and TDP). Further monitoring will determine if this is due to the unusually wet seasons. Suspended sediment concentrations were higher in watersheds with higher percent area devoted to agriculture production or higher upland slopes. Continued data collection (sediment, nutrient, and precipitation), rating curve development, and compilation of available historical data will be performed in the coming year.

6. References

- Demissie, M., L. Keefer, and J. Slowikowski, A. Russell, T. Snider, and K. Stevenson. 2001. Sediment and Nutrient Monitoring at Selected Watersheds within the Illinois River Watershed for Evaluating the Effectiveness of the Illinois River Conservation Reserve Enhancement Program (CREP). Illinois State Water Survey Contract Report 2001-12.
- Edwards, T.K. and D.G. Glysson. 1999. Field Methods for Measurement of Fluvial Sediment. U.S. Geological Survey, Techniques of Water-Resources Investigations, Book 3, Chapter 32, 89p.
- Federal Interagency Sedimentation Project (FISP). 1952. The Design of Improved Types of Suspended Sediment Samplers. Interagency Report 6, St. Anthony Falls Hydraulics Laboratory, Minneapolis, MN.
- Illinois Department of Natural Resources. 2000a. *Kaskaskia River Area Assessment, Volume 1: Geology* Springfield, IL, 127 p.
- Illinois Department of Natural Resources. 2000b. *Kaskaskia River Area Assessment, Volume 2: Water Resources* Springfield, IL, 127 p.
- Knapp, H. V., G. S. Roadcap, E. G. Bekele, H. A. Wehrmann, W. E. Gillespie, J. S. Hecht and F. J. Pisani. 2012. *Water Supply Assessment for Kaskaskia River Watershed Development: Phase I Technical Report*. Illinois State Water Survey Contract Report 2012-12, 29 +Appendices p.
- Leighton, M. M., G. E. Ekblaw and L. Horberg. 1948. *Physiographic Divisions of Illinois*. Illinois State Geological Survey Report of Investigations 129, Urbana, ILp.
- Luman, D.E., 2008, Assessment & Potential of the 2007 USDA-NASS Cropland Data Layer for Statewide Annual Land Cover Applications, unpublished report of investigations including five appendices, Illinois State Geological Survey, 17 pp.
<http://crystal.isgs.uiuc.edu/nsdihome/webdocs/landcover/nass07.html>.
- Luman, D. and T. Weicherding. *Land Cover of Illinois: 1999-2000*. 1999 January 1999 [cited 2003 November 1, 2003]; Available from: <http://www.agr.state.il.us/gis/index.html>.
- Rantz, S. E. (1982). Measurement and Computation of Streamflow. Volume 1. Measurement of Stage and Discharge. Government Printing Office, Washington, D.C.
- Rantz, S. E. (1982). Measurement and Computation of Streamflow. Volume 2. Computation of Discharge. Government Printing Office, Washington, D.C.

