### black vulture

## Coragyps atratus

#### **FEATURES**

The black vulture averages 23 to 27 inches in length. It has a short tail. Except for white patches in the wing tips which can be seen as it flies, this bird is entirely black, including its bare head and legs.

#### **BEHAVIORS**

The black vulture is an uncommon, permanent resident in extreme southern Illinois. It is occasionally seen as a vagrant elsewhere in the state. Nesting occurs in April and May. No nest is built. The female lays two eggs in a shallow cave or tree cavity. The gray-green to dull-white eggs have brown marks. Both the male and female take turns incubating during the 28- to 39-day incubation period. This bird lives in swampy woodlands and bluffs. It feeds by scavenging for dead organisms.

#### **TAXONOMY**

Kingdom: Animalia Phylum: Chordata

Class: Aves

Order: Cathartiformes Family: Cathartidae

#### **ILLINOIS STATUS**

common, native

#### **ILLINOIS RANGE**





© Illinois Department of Natural Resources. 2021. *Biodiversity of Illinois*. Unless otherwise noted, photos and images © Illinois Department of Natural Resources.



# **Aquatic Habitats**

bottomland forests

### **Woodland Habitats**

bottomland forests; southern Illinois lowlands

# **Prairie and Edge Habitats**

none