

# gray catbird

*Dumetella carolinensis*

## FEATURES

The gray catbird averages about nine inches in length (tail tip to bill tip in preserved specimen). This bird has gray feathers all over except for the black cap and the rust-colored patch under the long tail. The large bill is turned slightly downward.

## BEHAVIORS

The gray catbird is a common migrant, summer resident and rare winter resident statewide. It lives in woodland edges and urban areas, staying low in brush. It has a varied song with each phrase repeated only once and including catlike sounds. Spring migrants arrive in Illinois in May. The breeding season lasts from May through July. The nest is built low to the ground in thickets, briars, shrubs or small trees. Both male and female construct the nest of twigs, leaves, grasses and other plant materials over a period of five to six days. Four, dark-green eggs are deposited by the female, and she incubates them for the entire 12- to 13-day incubation period. The nest is often parasitized by the brown-headed cowbird that deposits an egg that the gray catbird will hatch and raise, taking food and care away from its own young. Unlike most birds, the catbird female has been known to throw the cowbird egg out of the nest. Fall migration starts in mid-October. The gray catbird winters as far south as Panama. It eats insects in spring and fruits at other times in the year.

## TAXONOMY

Kingdom: Animalia

Phylum: Chordata

Class: Aves

Order: Passeriformes

Family: Mimidae

## ILLINOIS STATUS

common, native

## ILLINOIS RANGE



© Illinois Department of Natural Resources. 2024. *Biodiversity of Illinois*.

Unless otherwise noted, photos and images © Illinois Department of Natural Resources.

photo provided by [stevebyland/pond5.com](https://pond5.com)





**Aquatic Habitats**

none

**Woodland Habitats**

none

**Prairie and Edge Habitats**

edge