

io moth

Automeris io

FEATURES

The wingspan of this species is two to three and one-fourth inches. The male and female have different coloration. For males, the upperside of the wings is yellow with brown spots, and a faint line can be seen in the middle of the forewing. The line parallels the edge of the forewing. The upperside of the female's wings is raspberry colored with markings similar to those of the male. The hindwing of both sexes contains a large black spot with a blue and white center surrounded by a yellow band, then a black line followed by another yellow band. The underside of the male's wings is yellow, while the underside of the female's wings is red-brown. The large spot is visible on the hindwing of both sexes.

BEHAVIORS

The Io moth is found statewide. The adults are active starting in late May and can be found in wooded areas and savannas. The adult does not feed, while the larvae feed on many species of plants. One generation is produced in northern Illinois while two generations are produced annually in the southern part of the state.

TAXONOMY

Kingdom: Animalia
Phylum: Arthropoda
Class: Insecta
Order: Lepidoptera
Family: Saturniidae

ILLINOIS STATUS

common, native

ILLINOIS RANGE



provided by Brett Hondow/pond5.com



Aquatic Habitats

bottomland forests

Woodland Habitats

bottomland forests; southern Illinois lowlands; upland deciduous forests

Prairie and Edge Habitats

edge; shrub prairie