

# jade clubtail

## *Arigomphus submedianus*

### FEATURES

The adult jade clubtail is about two inches long and has a pale green body with a dark spot on the end of its long abdomen. It has small, widely separated, green eyes and a pale, green face. Females are slightly paler in color than males and lack the swollen region at the end of the abdomen that gives the species its name. They are easily mistaken for the stillwater clubtail (*Arigomphus lentulus*) but have smaller stripes on their thorax; and while the jade clubtail is found statewide, the stillwater clubtail has only been observed on the southeastern side of the state.

### BEHAVIORS

Larval species of the family Gomphidae are burrowers and have short thick antennae with wedge-shaped heads. Adults can be seen in late May to July statewide near slow-moving rivers, ponds, and lakes. They make short, sporadic flights and perch on low lying vegetation and brush near shorelines. Copulating pairs find shelter on shoreline vegetation. Females lay eggs without help of male by flying along the surface of the water and releasing eggs beneath the surface at irregular intervals.

### TAXONOMY

Kingdom: Animalia  
Phylum: Arthropoda  
Class: Insecta  
Order: Odonata  
Family: Gomphidae

### ILLINOIS STATUS

common, native

### ILLINOIS RANGE



© Mary Kay Rubey


female



© Mary Kay Rubey

male





### **Aquatic Habitats**

lakes, ponds and reservoirs; marshes; rivers and streams; swamps; temporary water supplies; wet prairies and fens

### **Woodland Habitats**

none

### **Prairie and Edge Habitats**

edge