## lesser scaup

# Aythya affinis

### **FEATURES**

The lesser scaup averages 15 to 18 inches in length (tail tip to bill tip in preserved specimen). This bird has a white stripe of feathers on the back edge of the wing that is visible when it flies. The male scaup has a blue bill, purple head feathers, black chest feathers and black tail feathers. All other areas on the male have light gray feathers with small, dark bars. The female has dark brown feathers with a white area between the bill and the eye.

### **BEHAVIORS**

The lesser scaup is a common migrant, an uncommon winter resident and a rare summer resident in Illinois. This diving duck may be seen on lakes, ponds and sewage lagoons during migration. Spring migrants begin arriving in Illinois in late March and April. The males move through before the females. Summers are spent in Canada and Alaska. Fall migration through Illinois starts in September. The lesser scaup winters in Florida and Louisiana. A few of these birds may winter in Illinois if open water is present. The scaup eats snails, mussels, mayflies and aquatic plants. It makes a "quack" or "scaup" sound.

#### **TAXONOMY**

Kingdom: Animalia Phylum: Chordata


Class: Aves

Order: Anseriformes Family: Anatidae

### **ILLINOIS STATUS**

common, native

### **ILLINOIS RANGE**



© Illinois Department of Natural Resources. 2024. *Biodiversity of Illinois*. Unless otherwise noted, photos and images © Illinois Department of Natural Resources.




adult male preserved specimen



adult female preserved specimen



adult drake (male) with nasal tag



## **Aquatic Habitats**

bottomland forests; lakes, ponds and reservoirs; Lake Michigan; marshes; peatlands; rivers and streams; swamps; wet prairies and fens

### **Woodland Habitats**

none

## **Prairie and Edge Habitats**

none