

Aquatic Species Fact Cards



The *Aquatic Species Fact Cards* can be used to support the lessons in Units 1 and 2. They can also be used independently for students to learn more about these species and to encourage further research.

STUDENT ACTIVITY PAGE | Aquatic Species Fact Cards

blue-green algae



BLUE-GREEN ALGAE

Kingdom Bacteria
Phylum Cyanobacteria


LOCATION - statewide - native

FEATURES - chlorophyll a; some have gas bubbles that allow them to float; may form filaments or clusters

HABITAT - any moist environment; tolerant of extreme conditions

HABITS - most are photosynthetic; carry out nitrogen fixation

water mold



WATER MOLD

Kingdom Chromalveolate
Phylum Heterokontophyta

LOCATION - statewide - native

FEATURES - saprophytic or parasitic; cellulose in cell walls; may cause disease in fishes and their eggs

HABITAT - some aquatic, some terrestrial


HABITS - often found on dead organisms in water; adds nutrients to surrounding area

ILLINOIS Equal opportunity to participate in programs of the Illinois Department of Natural Resources (IDNR) and those funded by the U.S. Fish and Wildlife Service and other agencies is available to all individuals regardless of race, sex, national origin, disability, age, religion or other non-merit factors. If you believe you have been discriminated against, contact the funding source's civil rights office and/or the Equal Employment Opportunity Officer, IDNR, One Natural Resources Way, Springfield, IL 62702-1271; 217/785-0067; TTY 217/782-9175. This information may be provided in an alternative format if required. Contact the DNR Clearinghouse at 217/782-7498 for assistance.



Aquatic Illinois © 2011, Illinois Department of Natural Resources
DNR 156 – 6/12 • IOCI 12-0831

protozoa



Kingdom Protozoa

LOCATION - statewide - native

FEATURES - lack chloroplasts and cell walls; unicellular; heterotrophs

HABITAT - aquatic or parasitic

HABITS - amoebas and relatives may secrete shells; some are covered with cilia; capture other organisms

sponge



Kingdom Animalia

Phylum Porifera

LOCATION - only where water is clear and free of debris - native

FEATURES - simple; multicellular; sessile

HABITAT - clear water

HABITS - filter-feeder, eating small organisms and particles in the water that passes through its body

hydra



Kingdom Animalia

Phylum Cnidaria

Class Hydrozoa

LOCATION - statewide - native

FEATURES - polyp is the typical life form; radial symmetry; nematocysts or "stinging cells" are found in the tentacles

HABITAT - bottom substrate of slow-moving water

HABITS - carnivore; eats small organisms trapped by its tentacles

leech

Kingdom Animalia

Phylum Annelida


Class Clitellata

LOCATION - statewide - native

FEATURES - sucker at each end of the body; segmented, flattened body; hermaphrodites

HABITAT - water or moist areas on land

HABITS - many are external parasites, feeding on the blood of fishes, turtles, amphibians, water birds and snails; others are predators or scavengers; can survive for months between meals



common cat-tail



CAT-TAIL

Typha latifolia

Kingdom Plantae
Division Tracheophyta

LOCATION - statewide - native

FEATURES - smooth, upright, unbranched stems up to 12

feet tall; leaves elongated and pointed at tip; many flowers in a dense spike

HABITAT - marshes or other wet areas

HABITS - spreads by seeds or rhizomes; flowers June-September

pin oak



PIN OAK

Quercus palustris

Kingdom Plantae
Division Tracheophyta


LOCATION - scattered throughout Illinois - native

FEATURES - may grow to 75 feet; lower branches droop; acorns 1-4 together

HABITAT - flood plain woods, along streams, edges of swamps and ponds

HABITS - flowers April-May with the male and female flowers on the same tree

water lotus



AMERICAN LOTUS

Nelumbo lutea

Kingdom Plantae
Division Tracheophyta

LOCATION - scattered throughout Illinois - native

FEATURES - round leaves up to 2½ feet across; large, solitary, pale yellow flowers

HABITAT - standing water, lakes, ponds, rivers

HABITS - flowers July-September; may form dense colonies

silver maple



SILVER MAPLE

Acer saccharinum

Kingdom Plantae
Division Tracheophyta

LOCATION - statewide - native

FEATURES - may reach 100 feet tall; bark gray or silver; leaves white on lower surface; winged seeds

HABITAT - bottomland woods

HABITS - flowers late February to early April

eastern cottonwood



Populus deltoides

Kingdom Plantae

Division Tracheophyta

LOCATION - statewide - native

FEATURES - may reach 100 feet tall; bark gray and furrowed at maturity; leaves triangular; male and female flowers on separate trees

HABITAT - bottomland woods

HABITS - flowers February-March; fruits are ellipsoid capsules in clusters

pitcher-plant



Sarracenia purpurea

Kingdom Plantae

Division Tracheophyta


LOCATION - northeastern Illinois - native

FEATURES - tubular green leaves streaked with purple and red; leaf is a pitfall trap for insects

HABITAT - bogs and fens

HABITS - insects and other small organisms are attracted to the leaf where they may be trapped and digested

dragonfly



Kingdom Animalia

Phylum Arthropoda

Class Insecta


LOCATION - statewide - native

FEATURES - wings held outstretched at rest; chewing mouthparts; long abdomen

HABITAT - adults usually found near water (in which the nymphs live)

HABITS - adults feed on insects they catch while in flight; nymphs feed on aquatic insects

crayfish



Kingdom Animalia

Phylum Arthropoda

Class Malacostraca

LOCATION - statewide - native

FEATURES - 5 pairs of legs; large claw on each of the first pair of legs

HABITAT - in water or in burrows constructed below ground

HABITS - eats dead vegetation and small invertebrates

mosquito



Kingdom Animalia
Phylum Arthropoda
Class Insecta

LOCATION - statewide - native

FEATURES - one pair of membranous wings; sucking mouthparts; complete metamorphosis

HABITAT - adults wide-ranging; larvae in still or stagnant water

HABITS - larvae feed on organic debris; adult females feed on blood; adult males feed on nectar and plant juices

caddisfly

Kingdom Animalia
Phylum Arthropoda
Class Insecta

LOCATION - statewide - native

FEATURES - wings held rooflike over body while resting; slender, elongate, mothlike insect

HABITAT - larvae in ponds and streams; adults near water

HABITS - larvae build a portable case and eat plant material; adults feed on liquids and live about one month



mayfly



Kingdom Animalia
Phylum Arthropoda
Class Insecta


LOCATION - statewide - native

FEATURES - wings held above body at rest; soft-bodied; nymph has gills along abdomen and three tails

HABITAT - nymphs aquatic; adults near water

HABITS - nymphs eat algae and aquatic plants; adults live for 1-2 days and do not feed

mussel



Kingdom Animalia
Phylum Mollusca
Class Bivalvia

LOCATION - statewide - native

FEATURES - soft inner body and hard outer shell

HABITAT - streams and lakes

HABITS - feeds on plankton and detritus that it filters from water; may live for 100 years

largemouth bass



LARGEMOUTH
BASS

Micropterus salmoides

Kingdom Animalia

Phylum Chordata

Superclass Osteichthyes

Class Actinopterygii


LOCATION - statewide - native

FEATURES - upper jaw extends beyond back of eye when mouth is closed; dark lateral stripe along length of body

HABITAT - lakes, ponds, sluggish rivers

HABITS - eats mainly fishes and crayfish; females may lay up to 100,000 eggs

bluegill



BLUEGILL

Lepomis macrochirus

Kingdom Animalia

Phylum Chordata

Superclass Osteichthyes

Class Actinopterygii

LOCATION - statewide - native

FEATURES - short, black gill flap with no margin; black spot at posterior end of dorsal fin

HABITAT - sluggish streams and impoundments

HABITS - eats insects and their larvae, crayfish, algae and fishes; females may lay up to 67,000 eggs

muskellunge



MUSKELLUNGE

Esox masquinongy

Kingdom Animalia

Phylum Chordata

Superclass Osteichthyes

Class Actinopterygii

LOCATION - hatchery-stocked lakes - exotic

FEATURES - body elongated; head large with duckbill jaws and canine teeth

HABITAT - lakes

HABITS - eats fishes; female may lay up to 265,000 eggs

channel catfish



CHANNEL
CATFISH

Ictalurus punctatus

Kingdom Animalia

Phylum Chordata

Superclass Osteichthyes

Class Actinopterygii

LOCATION - statewide - native

FEATURES - no scales; eight barbels; adipose fin; deeply forked tail; rounded anal fin; single hard spine in pectoral and dorsal fins

HABITAT - lakes, ponds, rivers

HABITS - eats plant and animal material; females may lay up to 70,000 eggs

shovelnose sturgeon

Scaphirhynchus platorynchus

Kingdom Animalia
Phylum Chordata
Superclass Osteichthyes
Class Actinopterygii

LOCATION - rivers - native

FEATURES - snout flattened and shovel-shaped; fringed barbels; upper lobe of tail fin with a long, slender filament

HABITAT - rivers

HABITS - eats insect larvae; females may lay as many as 200,000 eggs



SHOVELNOSE STURGEON

blackstripe topminnow

Fundulus notatus

Kingdom Animalia
Phylum Chordata
Superclass Osteichthyes
Class Actinopterygii

LOCATION - statewide - native

FEATURES - front of dorsal fin base situated above anal fin base; brown with a broad, black horizontal stripe


HABITAT - large lowland rivers and pools of streams

HABITS - eats insects, crustaceans and snails; female may lay up to 30 eggs at once



TOPMINNOW

bullfrog



BULLFROG

Lithobates catesbeianus

Kingdom Animalia
Phylum Chordata
Class Amphibia

LOCATION - statewide - native

FEATURES - 3½-6 inches; no dorsolateral folds; tympanum larger than eye; olive, green or brown dorsum; call is "jug-o-rum"

HABITAT - any type of permanent water

HABITS - solitary animal; eats most anything it can swallow

mudpuppy

Necturus maculosus


Kingdom Animalia
Phylum Chordata
Class Amphibia

LOCATION - statewide - native

FEATURES - 8-13 inches; aquatic; four toes per foot; permanent bushy gills; slimy

HABITAT - lakes, lagoons, rivers, large creeks

HABITS - eats fishes, arthropods, annelids and mollusks



MUDPUPPY

northern water snake

Nerodia sipedon

Kingdom Animalia

Phylum Chordata

Class Reptilia

LOCATION - northern ½ of the state - native

FEATURES - 24-42 inches; keeled scales; heavy-bodied; venter with numerous semilunar brown or red markings

HABITAT - streams, lakes, ponds

HABITS - active day and night; eats fishes and amphibians



WATER SNAKE

spotted salamander

Ambystoma maculatum

Kingdom Animalia

Phylum Chordata

Class Amphibia

LOCATION - extreme northeastern and northwestern Illinois, the eastern border and the southern ⅓ of the state - native

FEATURES - 6-7¼ inches; blue-black, slate or brownish colored with two rows of yellow-white or orange-yellow spots from head onto tail

HABITAT - high quality forested areas with permanent or temporary ponds

HABITS - active at night; eats arthropods, mollusks and annelids



SPOTTED SALAMANDER

spiny softshell turtle

Apalone spinifera

Kingdom Animalia

Phylum Chordata


Class Reptilia

LOCATION - statewide - native

FEATURES - 5-17 inches; pointed nose; leathery shell; anterior edge of carapace with bumps


HABITAT - lakes, sloughs, rivers, streams

HABITS - may bask on sandbar or bury itself in sand at stream's edge; eats crustaceans, fishes, amphibians, insects and mollusks



SOFTSHELL TURTLE

snapping turtle



SNAPPING TURTLE

Chelydra serpentina

Kingdom Animalia

Phylum Chordata

Class Reptilia


LOCATION - statewide - native

FEATURES - 8-12 inches; 10-35 pounds; strong jaws; enormous head, thick legs

HABITAT - all types of water bodies

HABITS - spends much time on the bottom waiting for prey; eats most anything

wood duck



Aix sponsa

Kingdom Animalia

Phylum Chordata

Class Aves

LOCATION - common migrant and summer resident statewide - native

FEATURES - male is highly colored, female is brown; long, square dark tail

HABITAT - wooded swamps, rivers, ponds

HABITS - perches in trees; eats acorns, seeds and corn in fall and insects in summer

red-winged blackbird

Agelaius phoeniceus

Kingdom Animalia

Phylum Chordata

Class Aves

LOCATION - migrant and summer resident statewide; winter resident in southern Illinois - native

FEATURES - male has bright red shoulder patches; female is brown streaked; song is "konk-ka-reee"

HABITAT - cultivated fields, drainage ditches, marshes, wood edges

HABITS - eats insects, seeds and waste grain; forms huge roosts in winter



belted kingfisher



Megasceryle alcyon

Kingdom Animalia

Phylum Chordata

Class Aves

LOCATION - common migrant statewide; uncommon summer resident - native

FEATURES - blue and white coloration; harsh, rattling call

HABITAT - around ponds, lakes, rivers, streams

HABITS - catches small fishes by diving; burrows into banks to lay eggs

great blue heron

Ardea herodias

Kingdom Animalia

Phylum Chordata

Class Aves

LOCATION - statewide migrant and summer resident - native

FEATURES - 42-52 inches; blue-gray color; long legs, long neck, daggerlike bill

HABITAT - major rivers, marshes, lakes, ponds

HABITS - usually nests in colonies; eats fishes, crayfish, frogs and other aquatic creatures



bank swallow



Riparia riparia

Kingdom Animalia

Phylum Chordata

Class Aves

LOCATION - common migrant and summer resident statewide - native

FEATURES - brown back; brown neck band; call is a dry, weak rattle

HABITAT - open areas, over fields, along rivers and over lakes and ponds

HABITS - needs banks associated with waterways, roadways or mounds of sand for nest sites; nests in colonies; eats insects

osprey



Pandion haliaetus

Kingdom Animalia

Phylum Chordata

Class Aves

LOCATION - statewide migrant - native

FEATURES - shows a crook in wings while flying; broad, black cheek patch

HABITAT - lakes, but may be found at rivers and ponds

HABITS - eats fishes that it catches with its feet

raccoon



Procyon lotor

Kingdom Animalia

Phylum Chordata

Class Mammalia

LOCATION - statewide - native

FEATURES - long tail banded with light and dark rings; black facial mask

HABITAT - any as long as a permanent water supply, den sites and food are available

HABITS - climbs and swims well; eats a variety of plant and animal matter

American mink



Mustela vison

Kingdom Animalia

Phylum Chordata

Class Mammalia

LOCATION - statewide - native

FEATURES - red-brown back; dark tip on tail; toes webbed

HABITAT - where permanent water supply is available

HABITS - nocturnal; good swimmer; eats muskrats, fishes, voles, mice, rabbits, crayfish, birds, reptiles, amphibians and insects

North American river otter

Lontra canadensis

Kingdom Animalia

Phylum Chordata

Class Mammalia

LOCATION - statewide - native


FEATURES - webbed toes; thick, short, brown fur; can close ears

HABITAT - streams, rivers, lakes, ponds, wetlands, drainage ditches, backwater areas

HABITS - active day or night; eats fishes, frogs and crayfishes; excellent swimmer



American beaver



Castor canadensis

Kingdom Animalia

Phylum Chordata

Class Mammalia

LOCATION - statewide - native

FEATURES - broad, flattened paddlelike tail; webbed hind feet; lips close behind front teeth

HABITAT - streams, rivers, ponds, lakes, drainage ditches, backwater areas

HABITS - eats bark of shrubs and trees, grasses, corn and aquatic plants; nocturnal

muskrat



Ondatra zibethicus

Kingdom Animalia

Phylum Chordata

Class Mammalia


LOCATION - statewide - native

FEATURES - tail elongated and nearly hairless; hind feet elongate and partly webbed

HABITAT - ponds, lakes, streams, drainage ditches, swamps

HABITS - nocturnal; eats cat-tails and other aquatic plants; may build houses of vegetation or burrow in banks

white-tailed deer



Odocoileus virginianus

Kingdom Animalia

Phylum Chordata

Class Mammalia

LOCATION - statewide - native

FEATURES - antlers on males; conspicuous white underside of tail

HABITAT - wooded areas along streams and swamps

HABITS - active at night and at dusk; eats waste grain, acorns and leaves and shoots of some trees