

**A Site Inventory of
Nature Preserves and State Forest in the
Illinois River Section of the
Illinois River and Mississippi River
Sand Areas Natural Division**

John E. Ebinger

Illinois Natural History Survey
607 East Peabody Drive
Champaign, Illinois 61820

Center for Biodiversity
Technical Report 2005 (4)

Prepared for and funded by:

Illinois Wildlife Preservation Fund
Illinois Department of Natural Resources
One Natural Resources Way
Springfield, Illinois 62702

RECEIVED
JUN 16 2005
FEDERAL AID

Vascular Flora of Manito Prairie Nature Preserve, Tazewell County, Illinois

William E. McClain, Adjunct Research Associate in Botany
Illinois State Museum, Springfield, Illinois 62701
and

Loy R. Phillippe and John E. Ebinger, Center for Biodiversity
Illinois Natural History Survey, Champaign, Illinois 61820

ABSTRACT

The vascular flora of Manito Prairie Nature Preserve was studied during the growing seasons of 2001 and 2002. Located on a gravel terrace of the Illinois River valley 11 km southwest of Pekin, Illinois, much of the preserve has been disturbed by past plowing and grazing, although small gravel prairie remnants exist. On the gravel prairie remnants *Schizachyrium scoparium* (Michx.) Nash was the leading dominant with an Importance Value of 61.8. Other common prairie species include *Dichanthelium oligosanthes* (Schult.) Gould, *Dalea purpurea* Vent., and *Echinacea pallida* Nutt. A total of 223 vascular plant species were encountered in the preserve, two fern and fern-allies, 40 monocots, and 181 dicots. Non-native species were common with 56 taxa, representing about 25% of the flora. The Floristic Quality Index for the preserve is 40.51.

INTRODUCTION

At the time of European settlement prairie vegetation covered about 60% of Illinois (Iverson et al. 1991). Most was "black soil" tall-grass prairie of the Grand Prairie and much of the Southern Till Plain Natural Divisions (Schwegman 1973). Other prairie community types were also found in Illinois, including sand prairies, gravel prairies, loess hill prairies, and glacial drift prairies. Of these, gravel prairies are extremely uncommon (Fell and Fell 1956).

Gravel prairies are rare in the midwestern United States, many having been destroyed by mining and farming operations. Some occur on kames or eskers mostly in the Northeastern Morainal Division of Illinois, but most occur in the northern half of Illinois on glacial outwash plains that resulted from erosional events during Wisconsin glaciation (Willman and Frye 1970, Willman 1973, King 1981). Here they are associated with valley train deposits along major river systems. Fell and Fell (1956) listed plant species and associations of a few gravel prairies along the Rock River in Winnebago County, while McFall (1984) listed plants found on Manito gravel prairie in Tazewell County. More recently the flora of a reconstructed gravel prairie in the Wabash River valley, Lawrence County, Illinois was studied (Edgin et al. 2003). Also, three gravel prairies were studied along Wea Creek, a tributary of the Wabash River in north-central Indiana (Post et al. 1985).

As little is known about the flora of gravel prairies in Illinois, a study of one of the few remaining examples of this community type was undertaken. Because Manito Prairie Nature Preserve (MPNP) still contains small, intact remnants of gravel prairie, and there is an earlier study giving the floristic composition of the preserve (McFall 1984), it was decided to examine the vegetation of this site. This study was undertaken to determine vascular plant species composition, vegetation structure, and floristic quality of the major plant communities at the MPNP.

STUDY SITE

MPNP is located in southern Tazewell County, about 11 km southwest of Pekin, Illinois on a terrace above the Illinois River floodplain (SW1/4 S15 T24N R6W). This site lies at the edge of the Illinois River Section of the Illinois River and Mississippi River Sand Areas Natural Division (Schwegman 1973). The gravels and sands that form this terrace were deposited during the post-glacial period of Wisconsin glaciation about 14,500 years ago. At that time glacial deposits in northeastern Illinois were breached causing the Kankakee Torrent (Willman 1973). These flood waters carried huge amounts of sand and gravel which were deposited along the broad floodplain of the Illinois River starting below Hennepin, Illinois. Subsequent erosion created these extensive gravel bluffs.

MPNP was dedicated in 1985, is 7.94 ha in size, and is situated on a sand and gravel terrace which forms a 6-10 m bluff above the Illinois River floodplain (Hunter 1966). Evaluated by the Illinois Natural Areas Inventory in 1977, it was identified as a site of state-wide significance (White 1978). Originally called Shoop Prairie, it presently consists of about 3.2 ha of Grade B dry gravel hill prairie (McFall 1984, White and Madany 1978). Most of the prairie is along the bluff of County Road 850E, and consists of three parts separated by ravines. The remainder of the preserve consists of successional upland fields that, in some places, have a high concentrations of prairie species, and a few ravines that are mostly dominated by woody species.

Climate at MPNP is continental with warm summers and cold winters. Based on weather data from Peoria, 25 km to the northwest, the mean annual precipitation is 91.5 cm, with the month of May having the highest rainfall (10.6 cm). Mean annual temperature is 10.4°C with the hottest month being July with an average of 23.9°C, and the coldest January with an average of -5.3°C. Frost-free days range from 150 to 204, the average being 176 (Midwestern Regional Climate Center 2002).

METHODS

MPNP was visited at various times during the growing seasons of 2001 to 2002. Voucher specimens of each plant species were collected, identified, and deposited in the Stover-Ebinger Herbarium of Eastern Illinois University, Charleston, Illinois (EIU), and the Illinois Natural History Survey, Champaign, Illinois (ILLS). Criteria for designating non-native species followed Mohlenbrock (1986) and Gleason and Cronquist (1991) while nomenclature follows Mohlenbrock (1986).

Four transects 25 m long were located randomly along cardinal compass directions in the mature gravel prairie remnants, and two transects 25 m long was located in the succes-

sional field. Along each transect, m^2 quadrats were located at 1 m intervals. Odd numbered quadrats were located on the right side of the transect line; even numbered quadrats on the left side of the transect line. Cover of each species was determined by using the Daubenmire cover class system (Daubenmire 1959) as modified by Bailey and Poulton (1968). The modified Daubenmire cover scale is as follows: class 1 = 0 to 1%; class 2 = >1 to 5%; class 3 = >5 to 25%; class 4 = >25 to 50%; class 5 = >50 to 75%; class 6 = >75 to 95%; class 7 = >95 to 100%. Importance value (IV) for ground layer species was determined by summing relative cover and relative frequency.

The Floristic Quality Index (FQI) was determined using the coefficient of conservatism (CC) assigned to each species by Taft et al. (1997). The CC for each taxon was determined by assigning an integer from 0 to 10 based on the species tolerance to disturbance and its fidelity to habitat integrity. The FQI is a weighted index of species richness (N), and is the arithmetic product of the mean CC, multiplied by the square-root of the species richness (\sqrt{N}) of an inventory sites: $FQI = \text{mean CC}(\sqrt{N})$.

For relatively small areas the FQI gives a rapid means of comparison, and an indication of the floristic integrity of the site. When used along with other floristic measures, such as quadrat-based sampling methods, it provides a method of making comparisons among sites. Prairie sites with an FQI of 35 or higher are considered good quality (Taft et al. 1997).

RESULTS

A total of 223 plant species within 169 genera and 63 families were documented (Appendix I). Fern and fern-allies were represented by only two species. Of the remainder, 40 were monocots in 5 families and 27 genera, and 181 were dicots in 56 families and 140 genera. Non-native exotic species were common, 56 being found, representing about 25% of the flora. Though an obvious feature of the preserve, these non-native species were mostly restricted to disturbed areas in and at the edge of the preserve, and were rarely encountered in the high quality gravel prairie community. Woody species were also common with 40 being found, 18% of the flora. Though mostly native, many of the woody species were invading the gravel prairie, with a few confined to wooded ravines that traverse the preserve. The most important plant families were Poaceae with 30 species, and Asteraceae with 28 species, followed by the Rosaceae and Fabaceae.

Mature dry gravel prairie

Located on the west- and southwest-facing slopes of the preserve, this prairie was, in many places, overgrown with numerous woody species that formed dense thickets. Between the thickets and toward the crests of the steep hillside were dry prairie remnants of good quality. Some state endangered and threatened species occurred here, including *Astragalus tennesseensis* (Tennessee milk vetch) and *Besseyia bullii* (kitten tails), along with *Hymenoxys acaulis* (lakeside daisy) which was planted and has persisted (Herkert and Ebinger 2002).

The bunch-grass *Schizachyrium scoparium* (little bluestem) was the dominant species with an IV of 61.8, being more than four times as abundant as the next most important species; *Dichanthelium oligosanthes* (panic-grass) with an IV of 12.3 (Table 1). Other

common grasses included *Sorghastrum nutans* (Indian grass), *Sporobolus cladestinus* (dropseed), and *Bouteloua curtipendula* (sideoats grama). Common forbs included *Dalea purpurea* (purple prairie clover), *Echinacea pallida* (pale coneflower), *Ambrosia psilostachya* (western ragweed) and *Opuntia humifusa* (prickly-pear).

Forty-one taxa were found in the plots, four of which were non-native species, while two were native woody invaders (Table 1). Of the remainder, ten were grasses and sedges and the rest were prairie forbs commonly associated with dry gravel prairies (White and Madany 1978). Bare ground and litter accounted for about 25% of the cover (Table 1). Most forbs were growing between the clumps of grasses. These clumps were commonly 10-35 cm across, bare ground being common between them.

Upland old field

All uplands in the preserve had been plowed before the area was acquired. Old plow-lines and distinct changes in vegetation determined disturbance. Though not diverse floristically, prairie plants were common in much of the uplands, and were the dominant species in many areas. Nineteen taxa were encountered in the plots (Table 2). Indian grass dominated with an IV of 86.5, while other common prairie grasses included *Dichanthelium oligosanthes* (IV of 17.9), and little bluestem (IV of 4.7). Only a few native prairie forbs were present, but non-native species were common components. Woody invasion was obvious with the presence of *Malus ioensis* (Iowa crab apple) which ranked second in IV (Table 2).

DISCUSSION

The floristic integrity of the entire nature preserve, as measured by the FQI of Taft et al. (1997) was 40.71, while the mean Coefficient of Conservation (CC) was 2.71. Twenty-three species had a CC of seven or greater. If non-native species were excluded from the calculation, the FQI was 46.81 and the mean CC was 3.62. Though exotic species were common site components, most were restricted to areas of disturbance. Only four exotic species were encountered in the plots of the high quality prairie and these all had IV's of less than 1.4 (Table 1). The high species diversity along with the high FQI qualifies this site as being regionally noteworthy (Taft et al. 1997).

The small remnants of dry gravel hill prairie within the MPNP represent a rare community type in the midwest. For this reason steps should be taken to maintain and expand this community. Extensive brush removal and prescribed burning will be needed to restore this gravel prairie to its former extent and quality. This need becomes obvious when comparisons are made with the species list of McFall (1984) and the present study. Some obvious losses have already occurred, and it is likely that more will occur in the future as the size of the remnants decrease. At least six prairie species appear to have been extirpated, including *Asclepias amplexicaulis* (sand milkweed), *Froelichia gracilis* (cottonweed), *Helianthus tuberosus* (Jerusalem artichoke), *Oenothera rhombipetala* (sand primrose), *Psoralea onobrychis* (French grass), and *Tephrosia virginiana* (goat's-rue). Other species appear to have declined in abundance. Both *Besseyia bullii* and *Muhlenbergia cuspidata* are restricted to one small population each. *Hymenoxys acaulis*, though originally planted in six spots, is reduced to one small population with only one

individual flowering in 2002. It is likely that more species will be extirpated in the future as remnant size decreases.

McFall (1984) recorded 212 vascular plant species from the MPNP, while during the present study 223 taxa were recorded. Much of the increase is in non-native species, though some woody taxa reported by McFall (1984) could not be relocated. Recent management activities on the preserve probably account for this decrease. Presently most of the gullies have been cleared of woody vegetation, and some of the thickets surrounding the gravel prairies have been removed. Also, the use of occasional fires on various parts of the preserve has help reduce woody invasion. Extensive thickets remain, however, and these will need to be removed.

ACKNOWLEDGMENTS

The authors would like to thank Dr. Gordon Tucker, Eastern Illinois University, for help with *Carex* identifications, and Michelle Simone, Natural Heritage Biologist, Illinois Department of Natural Resources for her help and advice.

LITERATURE CITED

- Bailey, A. W., and C. E. Poulton. 1968. Plant communities and environmental relationships in a portion of the Tillamook burn, northwestern Oregon. *Ecology* 49:1-13.
- Daubenmire, R. 1959. A canopy coverage method of vegetation analysis. *Northwest Science* 33:43-64.
- Edgin, B., B. Garrard, G. C. Tucker, and J. E. Ebinger. 2003. Vegetation of Allison Prairie - a gravel prairie reconstruction in Lawrence County, Illinois. *Erigenia* 19:52-59.
- Fell, E. W. and G. B. Fell. 1956. The gravel-hill prairies of Rock River Valley in Illinois. *Transactions of the Illinois State Academy of Science*. 49:47-62.
- Gleason, H. A., and A. Cronquist. 1991. *Manual of the vascular flora of northeastern United States and adjacent Canada*. Second Edition. The New York Botanical Garden, Bronx, New York. lxxv + 910 pp.
- Herkert, J. R. and J. E. Ebinger, 2002. editors. *Endangered and threatened species of Illinois: Status and distribution*. Volume 1. Plants. Illinois Endangered Species Protection Board, Springfield, Illinois. 161 pp.
- Hunter, R. E. 1966. Sand and gravel resources of Tazewell County, Illinois. *State Geological Survey Circular* 399. 22 pp.
- Iverson, L. R., G. L. Rolfe, T. J. Jacob, A. S. Hodgins, and M. R. Jeffords. 1991. *Forests of Illinois*. Illinois Council on Forest development, Urbana, and Illinois Natural History Survey, Champaign, Illinois.
- King, J. E. 1981. Late Quaternary vegetational history of Illinois. *Ecological Monographs* 51:43-62.
- McFall, D. W. 1984. Vascular plants of the Manito Gravel Prairie, Tazewell County, Illinois. *Transaction of the Illinois State Academy of Science* 77:9-14.
- Midwestern Regional Climate Center. 2002. <http://mcc.sws.uiuc.edu>
- Mohlenbrock, R. H. 1986. *Guide to the Vascular Flora of Illinois, revised and enlarged edition*, Southern Illinois University Press, Carbondale, Illinois. viii + 507 pp.
- Post, T. W., J. A. Bacone, and J. R. Aldrich. 1985. Gravel prairies of Indiana. *Proceedings of the Indiana Academy* 94:457-463.
- Schwegman, J. E. 1973. *Comprehensive plan for the Illinois nature preserves system*. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, Illinois. 32 pp + map.
- Taft, J. B., G. S. Wilhelm, D. M. Ladd, and L. A. Masters. 1997. Floristic quality assessment for vegetation in Illinois, a method for assessing vegetation integrity. *Erigenia* 15:3-95.
- White J. 1978. Illinois Natural Areas inventory. Technical report. Volume I. Survey methods and results. Illinois Natural Areas Inventory, Urbana. Xix + 426 pp.
- White, J., and M. H. Madany. 1978. Classification of natural communities in Illinois. Pp. 310-405 in Illinois natural areas inventory. Technical report. (J. White, Editor). Illinois Natural Areas Inventory, Urbana, Illinois.
- Willman, H. B. 1973. Geology along the Illinois waterway - a basis for environmental planning. *Illinois State Geological Survey Circular* 478. Urbana. 48 pp.
- Willman, H. B. and J. C. Frye. 1970. Pleistocene stratigraphy of Illinois. *Illinois State Geological Survey Bulletin* 94:1-204.

Table 1. Frequency (%), average cover, relative frequency, relative cover, and importance value of ground layer species in a xeric gravel prairie at Manito Prairie Nature Preserve, Tazewell County, Illinois. (* = non-native)

Species	Freq. %	Average Cover	Rel. Freq.	Rel. Cover	I. V.
<i>Schizachyrium scoparium</i>	100	35.43	13.2	48.6	61.8
<i>Dichanthelium oligosanthes</i>	52	3.91	6.9	5.4	12.3
<i>Dalea purpurea</i>	53	3.55	7.0	4.9	11.9
<i>Echinacea pallida</i>	35	4.99	4.6	6.8	11.4
<i>Ambrosia psilostachya</i>	40	2.60	5.3	3.6	8.9
<i>Sorghastrum nutans</i>	31	2.97	4.0	4.1	8.1
<i>Opuntia humifusa</i>	45	1.54	6.0	2.1	8.1
<i>Lespedeza capitata</i>	40	1.99	5.3	2.7	8.0
<i>Heterotheca camporum</i>	29	2.93	3.9	4.0	7.9
<i>Sporobolus clandestinus</i>	31	1.65	4.0	2.3	6.3
<i>Senecio plattensis</i>	20	1.88	2.6	2.6	5.2
<i>Bouteloua curtipendula</i>	28	0.95	3.7	1.3	5.0
<i>Malus ioensis</i>	16	2.03	2.1	2.8	4.9
<i>Sporobolus asper</i>	20	1.37	2.6	1.9	4.5
<i>Euphorbia corollata</i>	17	1.09	2.3	1.5	3.8
<i>Asclepias verticillata</i>	27	0.13	3.5	0.2	3.7
<i>Bouteloua hirsuta</i>	17	0.93	2.3	1.3	3.6
<i>Cyperus filiculmis</i>	24	0.15	3.2	0.2	3.4
<i>Desmodium illinoense</i>	15	0.56	1.9	0.8	2.7
<i>Ruellia humilis</i>	15	0.37	1.9	0.5	2.4
<i>Potentilla arguta</i>	13	0.33	1.8	0.5	2.3
<i>Oxalis dillenii</i>	13	0.07	1.8	0.1	1.9
<i>Brickellia eupatorioides</i>	9	0.21	1.1	0.3	1.4
* <i>Poa pratensis</i>	9	0.15	1.1	0.2	1.3
* <i>Achillea millefolium</i>	9	0.11	1.1	0.1	1.2
<i>Oenothera biennis</i>	8	0.11	1.1	0.1	1.2
* <i>Mirabilis nyctaginea</i>	7	0.10	0.9	0.1	1.0
<i>Crataegus mollis</i>	3	0.08	0.4	0.1	0.5
<i>Eupatorium altissimum</i>	3	0.08	0.4	0.1	0.5
<i>Gaura biennis</i>	3	0.08	0.4	0.1	0.5
<i>Silene stellata</i>	3	0.05	0.4	0.1	0.5
<i>Solidago juncea</i>	3	0.08	0.4	0.1	0.5
<i>Carex</i> spp.	4	0.02	0.5	--	0.5
<i>Cassia fasciculata</i>	4	0.02	0.5	--	0.5
<i>Antennaria neglecta</i>	1	0.20	0.2	0.2	0.4
* <i>Kummerowia stipulacea</i>	3	0.01	0.4	--	0.4
<i>Phlox bifida</i>	3	0.01	0.4	--	0.4
<i>Aster pilosus</i>	1	0.04	0.2	0.1	0.3
<i>Callirhoe triangulata</i>	1	0.04	0.2	0.1	0.3
<i>Lactuca canadensis</i>	1	0.04	0.2	0.1	0.3
<i>Oxalis violacea</i>	1	0.01	0.2	--	0.2
Totals		72.86	100.0	100.0	200.0
Average bare ground and litter		25.10			

Table 2. Frequency (%), average cover, relative frequency, relative cover, and importance value of ground layer species in an upland old field at Manito Prairie Nature Preserve, Tazewell County, Illinois. (* = non-native)

Species	Freq. %	Average Cover	Rel. Freq.	Rel. Cover	Freq. I. V.
<i>Sorghastrum nutans</i>	100	71.10	14.3	72.2	86.5
<i>Malus ioensis</i>	92	10.34	13.1	10.5	23.6
<i>Dichanthelium oligosanthes</i>	96	4.10	13.7	4.2	17.9
* <i>Poa pratensis</i>	100	3.48	14.3	3.5	17.8
* <i>Rumex acetosella</i>	76	0.48	10.8	0.5	11.3
<i>Fragaria virginiana</i>	56	3.12	8.0	3.2	11.2
<i>Oxalis dillenii</i>	44	0.32	6.3	0.3	6.6
<i>Asclepias verticillata</i>	32	0.46	4.6	0.5	5.1
<i>Schizachyrium scoparium</i>	16	2.40	2.3	2.4	4.7
* <i>Bromus inermis</i>	12	1.22	1.7	1.2	2.9
<i>Carex</i> spp.	16	0.48	2.3	0.5	2.8
<i>Oenothera biennis</i>	12	0.26	1.7	0.3	2.0
<i>Potentilla simplex</i>	12	0.16	1.7	0.2	1.9
* <i>Potentilla recta</i>	12	0.06	1.7	0.1	1.8
* <i>Kummerowia stipulacea</i>	8	0.04	1.1	0.1	1.2
<i>Potentilla arguta</i>	4	0.12	0.6	0.1	0.7
<i>Rubus occidentalis</i>	4	0.12	0.6	0.1	0.7
* <i>Verbascum thapsus</i>	4	0.12	0.6	0.1	0.7
<i>Conyza canadensis</i>	4	0.02	0.6	—	0.6
Totals		98.40	100.0	100.0	200.0
Average bare ground and litter		0.50			

APPENDIX I.

Vascular taxa encountered at Manito Prairie Nature Preserve, Tazewell County, Illinois, are listed alphabetically by family under major plant groups. Non-native (exotic) species are indicated by an asterisk (*). For each species the author's collection number (JEE) is given, and the specimens are deposited in the Stover-Ebinger Herbarium of Eastern Illinois University (EIU). Collecting number preceded by P were collected by Loy R. Philippe, and the specimens are deposited in the Illinois Natural History Survey Herbarium (ILLS).

FERN AND FERN-ALLIES

Aspleniaceae

Asplenium platyneuron (L.) BSP. P35679

Equisetaceae

Equisetum laevigatum A.Br. 30720

MONOCOTS

Commelinaceae

**Commelina communis* L. 31168

Tradescantia ohiensis Raf. 30721

Cyperaceae

Carex aggregata Mack. P35681

Carex bebbii (Bailey) Fern. 30722

Carex blanda Dewey 30551

Carex muhlenbergii Willd. 30723

Carex pennsylvanica Lam. 30552

Cyperus filiculmis Vahl 30906

Liliaceae

**Asparagus officinalis* L. 30861

Poaceae

**Agropyron repens* (L.) Beauv. 30862

Agrostis hyemalis (Walt.) BSP. P35684

Andropogon gerardii Vitman 31169

Bouteloua curtipendula (Michx.) Torr.
30907

Bouteloua hirsuta Lag. 31122

**Bromus inermis* Leyss. 30724

**Bromus tectorum* L. 30553

**Chloris verticillata* Nutt. 31170

**Dactylis glomerata* L. 30725

Dichanthelium oligosanthes (Schult.) Gould
30727

**Digitaria sanguinalis* (L.) Scop. 31171

Elymus canadensis L. 30908

Elymus virginicus L. 30909

Eragrostis spectabilis (Pursh) Steud. 31172

Eragrostis trichodes (Nutt.) Wood 31067

**Festuca pratensis* Huds. 30726

Leptoloma cognatum (Schult.) Chase
31083

Muhlenbergia cuspidata (Torr.) Rydb.
31084

Muhlenbergia frondosa (Poir.) Fern. 31173

Muhlenbergia schreberi J.F. Gmel. P36147

**Poa compressa* L. 30728

**Poa pratensis* L. 30729

Schizachyrium scoparium (Michx.) Nash
31068

**Setaria viridis* (L.) Beauv. 31069

Sorghastrum nutans (L.) Nash 31085

Sporobolus asper (Michx.) Kunth 31121

Sporobolus clandenstinus (Biehler) Hitchc.
31120

Sporobolus heterolepis (Gray) Gray 31174

Tridens flavus (L.) Hitchcock 31070

Vulpia octoflora (Walt.) Rydb. 31086

Smilacaceae

Smilax hispida Muhl. 30730

DICOTS

Acanthaceae

Ruellia humilis Nutt. 30863

Anacardiaceae

Rhus aromatica Ait. 30554

Rhus glabra L. 31087

Toxicodendron radicans (L.) Kuntze 30731

Apiaceae

Sanicula canadensis L. 30864

Spermolepis inermis (Nutt.) Math. & Con-
stance P35685

Apocynaceae

Apocynum cannabinum L. 30865

Asclepiadaceae

Asclepias syriaca L. 30866

Asclepias tuberosa L. 30867

Asclepias verticillata L. 30868

Asclepias viridiflora Raf. P35686

Asteraceae

**Achillea millefolium* L. 30732

Ambrosia artemisiifolia L. 31071
Ambrosia psilostachya DC. 31072
Ambrosia trifida L. 31073
Antennaria neglecta Greene 30555
 **Arctium minus* Bernh. 31175
Aster ericoides L. 31177
Aster pilosus Willd. 31176
Brickellia eupatorioides (L.) Shinnars 31074
Cirsium discolor (Muhl.) Spreng. 31088
 **Cirsium vulgare* (Savi) Tenore 30910
Conyza canadensis (L.) Cronq. 31075
Echinacea pallida Nutt. 30869
Erigeron strigosus Muhl. 30733
Eupatorium altissimum L. 31090
Eupatorium rugosum Houtt. 31089
Gnaphalium obtusifolium L. 31178
Heterotheca camporum (Greene) Shinnars
 30870
Hymenoxys acaulis (Pursh) Parker 30591
Lactuca canadensis L. 30911
 **Lactuca serriola* L. 30912
Rudbeckia hirta L. 30871
Senecio plattensis Nutt. 30556
Solidago canadensis L. 31179
Solidago juncea Ait. 30913
 **Sonchus oleraceus* L. 30914
 **Taraxacum officinale* Weber 30557
 **Tragopogon dubius* Scop. 30734

Boraginaceae

**Buglossoides arvensis* (L.) I.M. Johnston
 30558
Hackelia virginiana (L.) I.M. Johnston
 30872

Lithospermum incisum Lehm. 31180
Mertensia virginica (L.) Pers. 30559
Onosmodium hispidissimum Mack. 30873

Brassicaceae

**Alliaria petiolata* (Bieb.) Cavara & Grande
 30560
Arabis glabra (L.) Bernh. 30735
Arabis shortii (Fern.) Gl. 30564
 **Capsella bursa-pastoris* (L.) Medic. 30561
Draba reptans (Lam.) Fern. 30562
 **Lepidium campestre* (L.) R. Br. 30736
 **Lepidium densiflorum* Schrad. 30563
Lepidium virginicum L. 30737

Cactaceae

Opuntia humifusa (Raf.) Raf. 30877

Caesalpiniaceae

Cassia fasciculata Michx. 30917
Gleditsia triacanthos L. 30918
Gymnocladus dioica (L.) K. Koch. 31185

Campanulaceae

Campanula americana L. 30874
Triodanis perfoliata (L.) Nieuwl. 30738

Caprifoliaceae

**Lonicera maackii* (Rupr.) Maxim. 30739
Sambucus canadensis L. 31091

Caryophyllaceae

**Arenaria serpyllifolia* L. P35683
 **Cerastium glomeratum* Thuill. 30565
 **Cerastium vulgatum* L. 30566
 **Dianthus armeria* L. 30916
 **Holosteum umbellatum* L. 30567
 **Lychnis alba* Mill. 30568
Minuartia stricta (Michx.) Hiern. 30740
 **Saponaria officinalis* L. 30875
Silene antirrhina L. 30569
Silene stellata (L.) Ait.f. 30876
 **Stellaria media* (L.) Vill. 30570

Celastraceae

Celastrus scandens L. 31092

Chenopodiaceae

**Chenopodium album* L. 31181

Cornaceae

Cornus drummondii C.A. Mey. 30878

Elaeagnaceae

**Elaeagnus umbellata* Thunb. 31182

Euphorbiaceae

Acalypha rhomboidea Raf. 31183
Chamaesyce maculata (L.) Small 31093
Croton glandulosus L. 31184
Euphorbia corollata L. 30879
Poinsettia cyathophora (Murr.) Kl. &
 Garcke 31095
Poinsettia dentata (Michx.) Kl. & Garcke
 31094

Fabaceae

Amorpha canescens Pursh 30880
Amorpha fruticosa L. 30741
Astragalus tennesseensis Gray 30571
Dalea purpurea Vent. 30882
Desmodium illinoense Gray 30883
Lespedeza capitata Michx. 31076
 **Kummerowia stipulacea* (Maxim.) Makino
 31123
 **Medicago lupulina* L. 30742
 **Melilotus alba* Medic. 30881
 **Melilotus officinalis* (L.) Pallas 30743
 **Trifolium arvense* L. P36145

Fagaceae

- Quercus imbricaria* Michx. 30919
Quercus macrocarpa Michx. 30920
Quercus rubra L. 31077

Geraniaceae

- Geranium carolinianum* L. 30744

Grossulariaceae

- Ribes missouriense* Nutt. 30745

Hydrophyllaceae

- Ellisia nyctelea* L. 30572
Hydrophyllum virginianum L. 30746

Hypericaceae

- Hypericum sphaerocarpum* Michx. 30884

Juglandaceae

- Carya tomentosa* (Poir.) Nutt. 31186
Juglans nigra L. 31078

Lamiaceae

- Agastache nepetoides* (L.) Ktze. 31096
Hedeoma hispida Pursh P35677
 **Lamium amplexicaule* L. 30573
 **Leonurus cardiaca* L. 30747
 **Nepeta cataria* L. 30885
Scutellaria parvula Michx. 30748
Teucrium canadense L. 30886
Trichostema brachiatum L. 31079

Lauraceae

- Sassafras albidum* (Nutt.) Nees 30921

Lythraceae

- Cuphea viscosissima* Jacq. 31097

Malvaceae

- Callirhoe triangulata* (Leavenw.) Gray
 30887

Menispermaceae

- Menispermum canadense* L. 30749

Moraceae

- **Cannabis sativa* L. 31080
 **Morus alba* L. 30922
Morus rubra L. 31098

Nyctaginaceae

- **Mirabilis nyctaginea* (Michx.) MacM.
 30750

Oleaceae

- Fraxinus americana* L. 31187

Onagraceae

- Gaura biennis* L. 30923
Oenothera biennis L. 30888

Oxalidaceae

- Oxalis dillenii* Jacq. 30751
Oxalis stricta L. P35680
Oxalis violacea L. 30574

Papaveraceae

- Corydalis micrantha* (Engelm.) Gray 30575

Phytolaccaceae

- Phytolacca americana* L. 31099

Plantaginaceae

- **Plantago lanceolata* L. 30924
Plantago rugelii Dcne. 31081
Plantago virginica L. 30752

Platanaceae

- Platanus occidentalis* L. 31188

Polemoniaceae

- Phlox bifida* Beck 30576

Polygalaceae

- Polygala verticillata* L. P36144

Polygonaceae

- **Polygonum convolvulus* L. P36149
Polygonum scandens L. 31189
Polygonum tenue Michx. 31100
 **Rumex acetosella* L. 30577
Rumex altissimus Wood P35687
 **Rumex crispus* L. 30753

Primulaceae

- Androsace occidentalis* Pursh 30578
Dodecatheon meadia L. 30579

Ranunculaceae

- Anemone caroliniana* Walt. 30581
Anemone cylindrica Gray 30754
Aquilegia canadensis L. 30582
Clematis pitcheri Torr. & Gray 30755
Ranunculus abortivus L. 30583

Rhamnaceae

- Rhamnus lanceolata* Pursh 29544

Rosaceae

- Crataegus mollis* (T. & G.) Scheele 30584
Fragaria virginiana Duchesne 30585
Geum canadense Jacq. 30889
Malus ioensis (Wood) Britt. 30586
Potentilla arguta Pursh 30890

~~**Potentilla recta* L. 30891~~*Potentilla simplex* Michx. 31124*Prunus americana* Marsh. 31298*Prunus serotina* Ehrh. 30756**Rosa multiflora* Thunb. 30757*Rubus flagellaris* Willd. 30587*Rubus occidentalis* L. 30758

Rubiaceae

Galium aparine L. 30759*Galium circaezans* Michx. 30892

Rutaceae

Ptelea trifoliata L. 30760*Zanthoxylum americanum* Mill. 30925

Saxifragaceae

Heuchera richardsonii R. Br. 30761

Scrophulariaceae

Besseyia bullii (Eat.) Rydb. 30590*Penstemon pallidus* Small 30762*Scrophularia lanceolata* Pursh 30893**Verbascum thapsus* L. 30894**Veronica arvensis* L. 30580

Solanaceae*Physalis heterophylla* Nees 30895*Physalis virginiana* Mill. P35678*Solanum carolinense* L. 30926

Ulmaceae

Celtis occidentalis L. 30927**Ulmus pumila* L. 31101*Ulmus rubra* Muhl. 30928

Urticaceae

Parietaria pensylvanica Muhl. 31082

Verbenaceae

Verbena stricta Vent. 30896*Verbena urticifolia* L. P36150

Violaceae

**Viola rafinesquii* Greene 30588*Viola sororia* Willd. 30589

Vitaceae

Parthenocissus quinquefolia (L.) Planch.

30929

Vitis riparia Michx. 30763

Vascular Flora of Long Branch Nature Preserve, Mason County, Illinois

Loy R. Phillippe, Mary Ann Feist, John E. Ebinger
Illinois Natural History Survey, Champaign, Illinois 61820
and
William E. McClain
Illinois State Museum, Springfield, Illinois 62706

ABSTRACT

Long Branch Nature Preserve, Mason County, Illinois is located in the Illinois River sand deposits in the central part of the state. Located on a large stabilized dune, 18 ha of the Preserve is dominated by a mature dry sand prairie community along with small savanna/woodland community. In the dry sand prairie community *Schizachyrium scoparium* (Michx.) Nash had the highest importance value (IV of 55.7 out of 200), followed by *Ambrosia psilostachya* DC. (IV of 28.5), and *Opuntia humifusa* (Raf.) Raf. (IV of 19.0). The savanna/woodland community was dominated by *Quercus marilandica* L. (IV of 180.7) followed by *Q. velutina* Lam. A total of 251 vascular plant species were found on the Preserve including six fern and fern-allies, three gymnosperms, 171 dicots, and 71 monocots. Forty-two non-native species were found, comprising nearly 17% of the flora. The Preserve had a Floristic Quality Index of 48.76 an indication of its high natural quality.

INTRODUCTION

Prairie vegetation was common in Illinois at the time of European settlement and covered about 60% of the state (Iverson et al. 1991). Most was tall-grass, black soil prairie that occurred in the prairie peninsula of northeastern Illinois (Transeau 1935, Schwegman 1973, Ebinger and McClain 1991). Depending upon soil and topography, other prairie types were common, including loess hill prairies, glacial till prairies, sand prairies, and gravel prairies (Schwegman 1973). Sand prairies were relatively common in the northern half of Illinois, most occurring on outwash plains that resulted from erosional events associated with Wisconsin glaciation (Willman and Frye 1970, King 1981).

Two extensive sand regions are the Kankakee sand deposits of northeastern Illinois, and the Illinois River sand deposits in the central part of the state (Gleason 1910, Schwegman 1973). The Kankakee sand deposits were formed when glacial lakes drained about 14,500 years ago after glacial moraines were breached, resulting in the Kankakee Torrent (Willman 1973). The Illinois River sand deposits were formed when waters of the Kankakee Torrent slowed as they entered the broad lowlands of the Illinois River below present day Hennepin.

The structure and composition of forest, woodland, and savanna communities of the Illinois River sand deposits have been studied by various workers (Jenkins et al. 1991, Coates et al. 1992, McClain et al. 2002). Also, Rodgers and Anderson (1979) examined presettlement vegetation, while Anderson and Brown (1983, 1986) determined effects of fire on sand savannas and adjacent forest. Since the studies of Gleason (1910) little information is available concerning the structure and composition of ground layer vegetation of the sand deposits. The only information available are studies of the dry sand prairie remnants at the Henry Allan Gleason Nature Preserve (McClain et al. 2005), and wetland sand communities at Matanzas Prairie Nature Preserve (Feist et al. 2005). The present study was undertaken to determine vascular plant species composition, vegetation structure, and floristic quality of major plant communities at Long Branch Nature Preserve (LBNP) and associated sand prairie remnants nearby.

DESCRIPTION OF THE STUDY SITE

LBNP is located in southwestern Mason County, about 8 km south of Havana, Illinois (NW1/4 S31 T21N R8W). Dedicated in 1989, this Preserve lies within the Illinois River Section of the Mississippi River and Illinois River Sand Area Natural Division (Schwegman 1973). This 38 ha Preserve is situated on a large dune, and though once grazed, an 18 ha section of the dune had never been subjected to major disturbances. The remainder of the Preserve had been farmed and about 4 ha had been planted to pines that have since been removed. The 18 ha dry sand prairie was designated "grade B" by the Illinois Natural Areas Inventory (White 1978). The soils are excessively drained Plainfield sands (Calsyn 1995) that are part of the dune and swale topography known as the Parkland Formation (Willman and Frye 1970).

LBNP has a continental climate with warm summers and cold winters. Based on weather data from Havana, mean annual precipitation is 96.0 cm, with May having the highest rainfall (11.3 cm). Mean annual temperature is 10.8°C with the hottest month being July (average of 24.6°C), and the coldest January (average of -5.0°C). Frost-free days range from 140 to 206, with the average being 173 day per year (Midwestern Regional Climate Center 2002).

METHODS

Between 1991 and 2003 the LBNP has been visited by scientists from the Illinois Natural History Survey. During these visits voucher specimens were collected, identified, and deposited in the herbarium of the Illinois Natural History Survey, Champaign, Illinois (ILLS), and the Stover-Ebinger Herbarium, Eastern Illinois University, Charleston, Illinois (EIU). Determination of non-native species followed Mohlenbrock (2002) and Gleason and Cronquist (1991). Nomenclature follows Mohlenbrock (2002) while the community classification follows those of White and Madany (1978).

In late summer of 2003 six 25 m transects were located randomly along cardinal compass directions in the dry sand prairie of the LBNP. Along each transect, 1 m² quadrates were spaced at 1 m intervals (n=25/transect), odd-numbered quadrates to the right, even-numbered quadrates to the left. A random numbers table was used to determine the number of meters (0 to 9) the quadrate was located from the transect line. Species cover was deter-

mined using the Daubenmire cover class system (Daubenmire 1959) as modified by Bailey and Poulton (1968). Importance value (IV) for ground layer species was determined by summing relative cover and relative frequency. In late summer of 2001 three sites were surveyed on private property just north of the Preserve, two mature dry sand prairie remnants (150 plots), and a disturbed dry sand prairie remnant (50 plots). The same procedure described above was used except the quadrates were 1/4 m². These sites on private land were studied as they contained some dry sand prairie remnants that differed from those found on the LBNP.

During the early summer of 2001, a 25 m by 50 m section of open woods along the north edge of LBNP was surveyed. In this small woodland all living woody individuals >10.0 cm dbh were identified and the diameters recorded. From this data, the living-stem density (stems/ha), basal area (m²/ha), relative density, relative dominance, importance value (IV), and average diameter (cm) were calculated for each species. Determination of the IV follows the procedure used by McIntosh (1957), and is the sum of the relative density and relative dominance (basal area).

The Floristic Quality Index (FQI) was determined using the coefficient of conservatism (CC) assigned to each species by Taft et al. (1997). The CC was determined by assigning each species an integer from 0 to 10 based on the species tolerance to disturbance and its fidelity to habitat integrity. Therefore, the FQI is a weighted index of species richness (N = number of species present), and is the arithmetic product of the average coefficient of conservatism (C-Value = the average of all species CC's) multiplied by the square root of the species richness (\sqrt{N}) of an inventory site: $FQI = C\text{-Value} (\sqrt{N})$. For relatively small areas that are intensively studied, the FQI gives a rapid means of comparison and an indication of the floristic integrity of the site. Using the FQI along with other floristic measures, such as quadrat-based sampling methods, provides a method of making comparisons among sites. Prairies with an FQI of 35 or higher are considered good quality natural areas (Taft et al. 1997).

RESULTS

A total of 251 vascular plant species within 179 genera and 69 families were documented for LBNP. Of these, six were fern and fern-allies, three gymnosperms, 171 dicots in 131 genera and 55 families, and 71 monocots in 41 genera and eight families (Appendix I). Forty-two non-native (exotic) species were encountered, about 17% of the species present. One threatened species, *Cyperus grayoides*, was encountered (Herkert and Ebinger 2002). The predominant plant families were the Poaceae with 39 species, the Asteraceae with 35 species, and the Cyperaceae with 19 species.

Mature dry sand prairie communities

Of the species found on the LBNP, 45 were present in the 150 quadrats sampled. Of these taxa *Schizachyrium scoparium* (little bluestem) was most important, having a frequency of 93%, an average cover of 26.77, and an IV of 55.7 (Table 1). Also common, *Ambrosia psilostachya* (western ragweed) was second with an IV of 28.5, while *Opuntia humifusa* (common prickly pear) was third with an IV of 19.0. Overall, five native prairie species, that are typical components of dry sand prairies, had IV's greater than 10. All would be expected in good quality dry sand prairie communities in Illinois. No exotic species were

encountered in the plots, and none were observed in the general area of the transects. The FQI for this site was 48.76 with a mean C-Value of 3.084 when all native and exotic species were included in the calculations.

In mature sand prairie remnants just north of LBNP on private land, similar results were obtained. In the larger of these remnants (Dry Sand Prairie # 1), about 3 ha in size, the same dominants were present, little bluestem having an IV of 73.9, followed by western ragweed (IV of 35.1), with common prickly pear being third (IV of 22.7) (Table 1). Most of the subordinate species were the same as in the LBNP mature sand prairie. One non-native species was encountered, *Rumex acetosella* (sour dock), with an IV of 3.5 (Table 1). On the second sand prairie remnant, also on private land and about 0.5 ha in size, the same three species were among the dominants, but *Helianthus occidentalis* (western sunflower) was second with an IV of 37.1, followed by western ragweed and common prickly pear (Table 1).

Disturbed dry sand prairie community

A disturbed sand prairie community that still contained much of its natural character was sampled. This community on private land just north of LBNP, was about 3 ha in size, had been disturbed by past grazing and probably off-road vehicles. Clumps of *Rhus aromatica* (fragrant sumac) and successional trees and shrubs dominated this site. Western ragweed (IV of 32.5) and common prickly pear (IV of 31.6) were important components, but *Conyza canadensis* (horseweed) with an IV of 36.7 was dominant, while *Eragrostis trichodes* (thread love grass), with an IV of 31.7 was third (Table 1). No non-native species were encountered in the plots.

Savanna/woodland communities

The small woodlots on LBNP had closed canopies, but trees became scattered and the canopy open near woodland edges. In the small remnant studied, *Quercus marilandica* (blackjack oak) dominated with an IV of 180.7 (Table 2). *Quercus velutina* (black Oak) was also present, and was also scattered throughout the dry sand prairie.

DISCUSSION

Though 42 non-native, adventive species were collected, most were restricted to disturbances at the edges of the Preserve, in the recently removed pine plantation, or the fallow field on the southern half of the Preserve. One exotic species was found in the study plots, sour dock, which is a pervasive species in most Illinois sand prairies. The fallow field also contained some woody species, most of which were exotics or invasive native species.

Dry sand prairie at LBNP is very similar to that at Henry Allan Gleason Nature Preserve 22 km to the northeast in northern Mason County, Illinois (McClain et al. 2005). Three of the top four dominants were the same with little bluestem dominant and western ragweed and common prickly pear important subdominants. *Tephrosia virginiana* (goat's-rue) was second in IV at Gleason Nature Preserve but was not found in the plots at LBNP. This species is rare at LBNP and had a clumped distribution (Table 1).

Gleason (1910) referred to dry sand prairie communities as the bunch-grass association. At LBNP the dry sand prairie was dominated by 20-40 cm clumps of little bluestem. Nearly circular in outline these clumps formed a dense mass through which few other species could grow. Mostly, other species grew in spaces between clumps. The lack of exotic species between these clumps, the high FQI, the high species diversity, and the large number of conservative prairie species present, indicate that the mature dry sand prairie remnant at the LBNP is of high natural quality.

ACKNOWLEDGMENTS

The authors thank Dr. Gordon Tucker, Eastern Illinois University, for help in the identification of species of Cyperaceae, and John Wilker, Natural Heritage Biologist, Illinois Department of Natural Resources, for his help and advice. The Illinois Department of Natural Resources, Wildlife Preservation Funds supported this project.

LITERATURE CITED

- Anderson, R.C. and L.E. Brown. 1983. Comparative effects of fire on trees in a midwestern savannah and an adjacent forest. *Bulletin of the Torrey Botanical Club* 110:87-90.
- Anderson, R.C. and L.E. Brown. 1986. Stability and instability in plant communities following fire. *American Journal of Botany* 73:364-368.
- Bailey, A.W. and C.E. Poulton. 1968. Plant communities and environmental relationships in a portion of the Tillamook burn, northwestern Oregon. *Ecology* 49:1-13.
- Calsyn, D.E. 1995. Soil survey of Mason County, Illinois. Soil Report 146, University of Illinois Agricultural Experiment Station, Urbana. ix+211 pp.
- Coates, D.T., S.E. Jenkins, and J.E. Ebinger. 1992. Woody vegetation survey of Barkhausen Woods, a closed canopy sand forest in Mason County, Illinois. *Erigenia* 12:1-6.
- Daubenmire, R. 1959. A canopy coverage method of vegetation analysis. *Northwest Science* 33:43-64.
- Ebinger, J.E. and W.E. McClain. 1991. Forest succession in the prairie peninsula of Illinois. *Illinois Natural History Survey Bulletin* 34:375-381.
- Feist, M.A., M.J. Morris, L.R. Phillippe, J.E. Ebinger, and W.E. McClain. 2005. Sand prairie communities of Matanzas Nature Preserve, Mason County, Illinois. (in review)
- Gleason, H.A. 1910. The vegetation of the inland sand deposits of Illinois. *Bulletin of the Illinois State Laboratory of Natural History* 9:21-174.
- Gleason, H.A., and A. Cronquist. 1991. *Manual of the vascular flora of northeastern United States and adjacent Canada*. Second Edition. The New York Botanical Garden, Bronx, New York. lxxv+910 pp.
- Herkert, J.R. and J.E. Ebinger. 2002. editors. *Endangered and Threatened Species of Illinois: Status and Distribution*. Volume 1: Plants. Endangered Species Protection Board, Springfield, Illinois. 161 pp.
- Iverson, L.R., G.L. Rolfe, T.J. Jacob, A.S. Hodgins, and M.R. Jeffords. 1991. *Forests of Illinois*. Illinois Council on Forest Development, Urbana, and Illinois Natural History Survey, Champaign, Illinois.
- Jenkins, S.E., J.E. Ebinger, and W.E. McClain. 1991. Woody vegetation survey of Bishop's Woods, a sand forest in Mason County, Illinois. *Transactions of the Illinois State Academy of Science* 84:20-27.
- King, J.E. 1981. Late Quaternary vegetational history of Illinois. *Ecological Monographs* 51:43-62.

- McClain, W.E., L.R. Phillippe, and J.E. Ebinger. 2005. Floristic assessment of the Henry Allan Gleason Nature Preserve, Mason County, Illinois. *Castanea* (in press)
- McClain, W.E., S.D. Turner, and J.E. Ebinger. 2002. Vegetation of forest communities at the Sand Prairie-Scrub Oak Nature Preserve, Mason County, Illinois. *Transactions of the Illinois State Academy of Science* 95:37-46.
- McIntosh, R.P. 1957. The York Woods. A case history in forest succession in southern Wisconsin. *Ecology* 38:29-37.
- Midwestern Regional Climate Center. 2002. <http://mcc.sws.uiuc.edu>
- Mohlenbrock, R.H. 2002. *Vascular Flora of Illinois*. Southern Illinois University Press, Carbondale, Illinois. xi+490 pp.
- Rodgers, C.S. and R.C. Anderson. 1979. Presettlement vegetation of two prairie peninsula counties. *Botanical Gazette* 140:232-240.
- Schwegman, J.E. 1973. Comprehensive plan for the Illinois nature preserves system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, Illinois. map+32 pp.
- Taft, J.B., G.S. Wilhelm, D. M. Ladd, and L.A. Masters. 1997. Floristic quality assessment for vegetation in Illinois, a method for assessing vegetation integrity. *Erigenia* 15:1-95.
- Transeau, E.N. 1935. The prairie peninsula. *Ecology* 16:423-437.
- White, J. 1978. Illinois Natural Areas Inventory. Technical report. Volume I. Survey methods and results. Illinois Natural Areas Inventory, Urbana. xix+426 pp.
- White, J. and M.H. Madany. 1978. Classification of natural communities in Illinois. Pp. 310-405 in Illinois natural areas inventory. Technical report. (J. White, Editor). Illinois Natural Areas Inventory, Urbana, Illinois.
- Willman, H.B. 1973. Geology along the Illinois waterway - a basis for environmental planning. Illinois State Geological Survey Circular 478. Urbana. 48 pp.
- Willman, H.B. and J.C. Frye. 1970. Pleistocene stratigraphy of Illinois. Illinois State Geological Survey Bulletin 94:1-204.

Table 1. Relative cover and importance values of ground layer species encountered in prairie communities at Long Branch Nature Preserve and adjacent private ground just north of the Preserve.

Species	LBNP		Private Property North of LBNP					
	Dry Sand Prairie (n=100)		Dry Sand Prairie #1 (n=100)		Dry Sand Prairie #2 (n=50)		Disturbed Sand Prairie (n=50)	
	Avg. Cover	I.V.	Avg. Cover	I.V.	Avg. Cover	I.V.	Avg. Cover	I.V.
<i>Schizachyrium scoparium</i>	26.77	55.7	26.68	73.9	23.38	57.8	--	--
<i>Ambrosia psilostachya</i>	10.81	28.5	9.42	35.1	9.07	30.0	9.74	32.5
<i>Opuntia humifusa</i>	5.74	19.0	5.86	22.7	4.53	17.1	9.13	31.6
<i>Leptoloma cognatum</i>	5.89	16.3	1.01	4.8	0.07	0.8	0.30	1.0
<i>Calamovilfa longifolia</i>	2.43	10.6	--	--	--	--	0.36	1.4
<i>Dichanthelium villosissimum</i>	1.37	6.8	1.32	8.7	3.61	15.1	3.66	14.4
<i>Coryza canadensis</i>	0.85	6.5	1.15	12.5	0.96	9.6	10.77	36.7
<i>Carex muhlenbergii</i>	0.34	6.2	0.42	3.7	0.35	4.2	1.17	10.5
<i>Crotonopsis linearis</i>	0.19	5.0	0.07	2.9	0.11	4.1	0.02	0.8
<i>Aristida tuberculosa</i>	0.61	4.9	0.32	3.8	--	--	0.03	1.2
<i>Commelina erecta</i>	0.24	3.6	--	--	--	--	--	--
<i>Cyperus lupulinus</i>	0.22	3.5	0.06	1.4	--	--	--	--
<i>Cyperus schweinitzii</i>	0.31	3.4	0.09	3.0	--	--	0.15	2.1
<i>Chamaechrista fasciculata</i>	0.31	3.2	0.03	1.2	0.02	0.8	--	--
<i>Euthamia gymnospermoides</i>	0.78	2.9	--	--	--	--	--	--
<i>Carex tomsa</i>	0.21	2.2	0.03	0.3	--	--	0.84	3.7
<i>Lespedeza capitata</i>	0.28	2.2	--	--	--	--	--	--
<i>Cyperus grayoides</i>	0.10	1.5	--	--	--	--	--	--
<i>Eragrostis spectabilis</i>	0.41	1.5	0.34	1.8	0.07	0.8	0.06	0.5
<i>Panicum virgatum</i>	0.63	1.5	--	--	--	--	--	--
<i>Paspalum bushii</i>	0.32	1.5	0.52	3.3	0.12	0.9	2.17	10.0
<i>Croton glandulosus</i>	0.07	1.4	0.03	1.1	0.04	1.5	0.06	2.1
<i>Rhus aromatica</i>	0.62	1.3	--	--	--	--	--	--
<i>Eragrostis trichodes</i>	0.25	1.2	1.40	5.1	1.63	5.7	8.82	31.7
<i>Oenothera rhombipetala</i>	0.06	1.2	0.09	3.0	1.23	6.0	0.87	8.1
<i>Froelichia floridana</i>	0.07	1.1	0.18	1.9	0.03	1.2	0.12	2.7
<i>Antennaria plantaginifolia</i>	0.03	0.9	--	--	--	--	--	--
<i>Chrysopsis camporum</i>	0.22	0.7	0.38	1.2	--	--	0.30	1.0
<i>Hieracium longipilum</i>	0.04	0.7	--	--	--	--	--	--
<i>Andropogon gerardii</i>	0.14	0.5	--	--	--	--	--	--
<i>Polygonum tenue</i>	0.02	0.5	--	--	0.02	0.7	--	--
<i>Pseudognaphalium obtusifolium</i>	0.02	0.4	0.20	2.3	0.07	0.8	0.36	1.4
<i>Solidago speciosa</i>	0.05	0.4	--	--	--	--	--	--
<i>Heterostipa spartea</i>	0.04	0.4	--	--	--	--	--	--
<i>Triplasis purpurea</i>	0.03	0.4	0.02	0.6	0.01	0.4	0.01	0.4
<i>Asclepias hirtella</i>	0.10	0.3	--	--	--	--	--	--
<i>Baptisia bracteata</i>	0.10	0.3	--	--	--	--	--	--
<i>Brickellia eupatorioides</i>	0.03	0.3	--	--	--	--	--	--
<i>Lactuca canadensis</i>	0.03	0.3	--	--	--	--	0.06	0.5
<i>Liarris aspera</i>	0.03	0.3	--	--	--	--	--	--
<i>Phlox bifida</i>	0.01	0.3	--	--	--	--	--	--
<i>Sorghastrum nutans</i>	0.03	0.3	--	--	--	--	--	--
<i>Chamaesyce geyeri</i>	0.01	0.1	--	--	0.02	0.8	--	--
<i>Chenopodium desiccatum</i>	0.01	0.1	--	--	--	--	0.01	0.4

Table 1. continued

Species	LBNP		Private Property North of LBNP					
	Dry Sand Prairie (n=100)		Dry Sand Prairie #1 (n=100)		Dry Sand Prairie #2 (n=50)		Disturbed Sand Prairie (n=50)	
	Avg. Cover	I.V.	Avg. Cover	I.V.	Avg. Cover	I.V.	Avg. Cover	I.V.
<i>Poinsettia dentata</i>	0.01	0.1	--	--	--	--	--	--
<i>Rumex acetosella</i>	--	--	0.32	3.5	--	--	--	--
<i>Euphorbia corollata</i>	--	--	0.22	1.5	--	--	--	--
<i>Teucrium canadense</i>	--	--	0.04	0.7	--	--	--	--
<i>Helianthus occidentalis</i>	--	--	--	--	12.02	37.1	--	--
<i>Fallopia cristatum</i>	--	--	--	--	0.22	2.9	--	--
<i>Koeleria macrantha</i>	--	--	--	--	0.36	1.3	--	--
<i>Lithospermum croceum</i>	--	--	--	--	0.01	0.4	--	--
<i>Tephrosia virginiana</i>	--	--	--	--	--	--	1.38	5.3
Totals	60.83	200.0	50.20	200.0	57.95	200.0	50.39	200.0
Average bare ground	42.45		44.85		38.90		59.45	

Table 2. Density (#/ha), basal area (m²/ha), relative values, and importance values of woody species in a small woodland at the north edge of Long Branch Nature Preserve, Mason County, Illinois

Species	Density (#/ha)	Basal Area (m ² /ha)	Relative Density	Relative Dominance	Importance Value	Avg. Diam. (cm)
<i>Quercus marilandica</i>	256	15.256	88.9	91.8	180.7	24.6
<i>Quercus velutina</i>	16	1.064	5.7	6.4	12.1	25.0
<i>Juniperus virginiana</i>	8	.208	2.7	1.3	4.0	18.1
<i>Prunus serotina</i>	8	.088	2.7	0.5	3.2	11.6
Totals	288	16.616	100.0	100.0	200.0	

APPENDIX I.

Vascular plant species encountered at Long Branch Nature Preserve, Mason County, Illinois, listed alphabetically by family in major plant groups. An asterisk indicates non-native species. John E. Ebinger (E) collections are deposited in the Stover-Ebinger Herbarium, Eastern Illinois University, Charleston, Illinois (EIU). Loy R. Phillippe (P) collections are deposited in the Illinois Natural History Survey Herbarium, Champaign, Illinois (ILLS).

FERN AND FERN-ALLIES

Aspleniaceae

Asplenium platyneuron (L.) Oakes P13288

Dryopteridaceae

Woodsia obtusa (Spreng.) Torr. E30370

Equisetaceae

Equisetum hyemale L. P13421

Equisetum laevigatum A. Br. P33301

Ophioglossaceae

Botrychium virginianum (L.) Sw. P13162

Thelypteridaceae

Thelypteris palustris Schott P13309

GYMNOSPERMAE

Cupressaceae

Juniperus virginiana L. P13300

Pinaceae

**Pinus banksiana* Lamb. P13327

**Pinus sylvestris* L. E31198

DICOTS

Acanthaceae

Ruellia humilis Nutt. P13279

Aceraceae

Acer negundo L. P13073

Amaranthaceae

Amaranthus albus L. P13640

Froelichia floridana (Nutt.) Moq. P13555

Froelichia gracilis (Hook.) Moq. P13433

Anacardiaceae

Rhus aromatica Ait. E28425

Toxicodendron radicans (L.) Kuntze
P13561

Apiaceae

Osmorhiza longistylis (Torr.) DC. P13291

Sanicula canadensis L. P13289

Apocynaceae

Apocynum cannabinum L. P33296

Apocynum sibiricum Jacq. P15299

Asclepiadaceae

Ampelamus albidus (Nutt.) Britt. P13414

Asclepias amplexicaulis Small P13274

Asclepias hirtella (Pennell) Woodson
P13314

Asclepias syriaca L. P13315

Asclepias verticillata L. P13419

Asteraceae

**Achillea millefolium* L. E31399

Ageratina altissima (L.) King & Robins.
P13647

Ambrosia artemisiifolia L. E29188

Ambrosia psilostachya DC. E29208

Antennaria plantaginifolia (L.) Hook.
E28416

Arnoglossum atriplicifolia (L.) H. Robins.
P13416

Aster ericoides L. E29411

Aster oblongifolius Nutt. P13629

Aster pilosus Willd. E29412

Bidens bipinnata L. E29201

Bidens comosa (Gray) Wieg. E29185

Bidens connata Muhl. P31290

Brickellia eupatorioides (L.) Shinnery
P13566

Chrysopsis camporum Greene P13319

Cirsium discolor (Muhl.) Spreng. P13613

Conyza canadensis (L.) Cronq. E29209

Coreopsis lanceolata L. P13310

Erigeron annuus (L.) Pers. P13278

Erigeron strigosus Muhl. E28621

Eupatorium serotinum Michx. E30371

Euthamia graminifolia (L.) Nutt. P31291

Euthamia gymnospermoides Greene E29196

Helianthus occidentalis Riddell E28836

**Helianthus petiolaris* Nutt. P13626

Heliopsis helianthoides (L.) Sweet P13646

Hieracium longipilum Torr. P13554

Krigia virginica (L.) Willd. P13140

Lactuca canadensis L. P13553

Liatris aspera Michx. P13603

Pseudognaphalium obtusifolium (L.)

Hilliard & Burt E29206

Senecio platensis Nutt. P13156

Solidago canadensis L. P13645

Solidago speciosa Nutt. P13628

**Taraxacum officinale* Weber P13074

**Tragopogon dubius* Scop. P13139

Boraginaceae

- Hackelia virginiana* (L.) I. M. Johnston
P33299
Lithospermum croceum Fern. P33302
Lithospermum incisum Lehm. P33306

Brassicaceae

- **Alliaria petiolata* (Bieb.) Cavara & Grande
E28420
Descurainia pinnata (Walt.) Britt. P13055
Draba reptans (Lam.) Fern. P13054
Erysimum capitatum (Dougl.) Greene
P13057
**Lepidium densiflorum* Schrad. P35696
Lepidium virginicum L. P13142

Cactaceae

- Opuntia humifusa* (Raf.) Raf. P13326

Caesalpinaceae

- Chamaechrista fasciculata* (Michx.) Greene
E28838
Gleditsia triacanthos L. P13312
Senna marilandica (L.) Link E30372

Campanulaceae

- Campanulastrum americana* (L.) Small
P33459
Triodanis perfoliata (L.) Nieuwl. P13282

Cannabinaceae

- **Cannabis sativa* L. P13624

Caprifoliaceae

- **Lonicera maackii* (Rupr.) Maxim. P33295
Sambucus canadensis L. P13290

Caryophyllaceae

- **Holosteum umbellatum* L. P13052
Silene stellata (L.) Ait. f. E30901
**Stellaria media* (L.) Cyrillo P13413

Celastraceae

- Celastrus scandens* L. P33458
Euonymus atropurpurea Jacq. P13062

Chenopodiaceae

- **Chenopodium album* L. E31398
**Chenopodium ambrosioides* L. P13642
Chenopodium desiccatum A. Neis. P13563
Cycloloma atriplicifolium (Spreng.) Coult.
P13415

Convolvulaceae

- **Ipomoea hederacea* (L.) Jacq. P13632

Cornaceae

- Cornus drummondii* C. A. Mey. P13155

Euphorbiaceae

- Chamaesyce geveri* (Engelm.) Small P13562
Croton glandulosus L. E28828
Crotonopsis linearis Michx. P13558
Euphorbia corollata L. P13283

Phyllanthus caroliniensis Walt. E30373

- Poinsettia cyathophora* (Murr.) Kl. &
Gracke E30374
Poinsettia dentata (Michx.) Kl. & Gracke
P13639

Fabaceae

- Amorpha canescens* Pursh P13427
Baptisia bracteata Ell. P13148
Desmodium illinoensis Gray P33309
Desmodium sessilifolium (Torr.) Torr. &
Gray P33461
Lespedeza capitata Michx. P13612
**Melilotus officinalis* (L.) Pallas. P13321
**Robinia pseudoacacia* L. P13175
Tephrosia virginiana (L.) Pers. P13306

Fagaceae

- Quercus x bushii* Sarg. E29409
Quercus marilandica Muenchh. P13154
Quercus velutina Lam. P13552

Geraniaceae

- Geranium carolinianum* L. P33300

Grossulariaceae

- Ribes missouriense* Nutt. P13071

Hydrophyllaceae

- Ellisia nyctelea* L. P13165

Hypericaceae

- Hypericum majus* (Gray) Britt. P33287
Hypericum mutilum L. P33456

Juglandaceae

- Juglans nigra* L. P13161

Lamiaceae

- Agastache nepetoides* (L.) Ktze. E29191
**Leonurus cardiaca* L. P13295
Lycopus americanus Muhl. P13644
Monarda punctata L. P13564
**Nepeta cataria* L. P13418
Teucrium canadense L. P13428

Lauraceae

- Sassafras albidum* (Nutt.) Nees P13060

Malvaceae

- Callirhoe triangulata* (Leavenw.) Gray
P13407

Menispermaceae

- Menispermum canadense* L. P13302

Molluginaceae

- **Mollugo verticillata* L. P13324

Moraceae

- **Maclura pomifera* (Raf.) Schneider P33281
**Morus alba* L. P13292

Nyctaginaceae

**Mirabilis nyctaginea* (Michx.) MacM.
P13286

Onagraceae

Circaea lutetiana Aschers. & Magnus
P33282
Ludwigia alternifolia L. E29194
Oenothera laciniata Hill P13143
Oenothera rhombipetala Nutt. P13406

Oxalidaceae

Oxalis stricta L. P13273

Phytolaccaceae

Phytolacca americana L. P13325

Plantaginaceae

Plantago patagonica Jacq. P13269

Polemoniaceae

Phlox bifida Beck P13065

Polygalaceae

Polygala polygama Walt. P33310
Polygala verucillata L. P33303

Polygonaceae

Fallopia cristata (Engelm. & Gray) Holub.
P31292

Persicaria coccinea (Muhl.) Greene P33297
**Persicaria hydropiper* (L.) Opiz. P33286
Persicaria pensylvanicum (L.) Small
P33284
Persicaria punctata (Ell.) Small E29193
Polygonum tenue Michx. P31293
**Rumex acetosella* L. P13276

Primulaceae

Androsace occidentalis Pursh P13053

Ranunculaceae

Anemone virginiana L. P13301
Clematis virginiana L. P33308
Ranunculus abortivus L. P13164

Rhamnaceae

Ceanothus americanus L. P13272

Rosaceae

Fragaria virginiana Duchesne P13146
Geum canadense Jacq. P13287
**Potentilla recta* L. P13320
Potentilla simplex Michx. P13147
Prunus serotina Ehrh. P13160b
Prunus virginiana L. P13072
Rosa carolina L. P13313
Rosa suffulta Greene P13275
Rubus argutus Link P13437
Rubus flagellaris Willd. P13149
Rubus occidentalis L. P13150

Rubiaceae

Diodia teres Walt. P13606

Galium aparine L. P13169

Galium circaezans Michx. P13305

Rutaceae

Ptelea trifoliata L. P13303
Zanthoxylum americanum Mill. P13067

Salicaceae

Populus tremuloides Michx. P13144
Salix humilis Marsh. P13064

Scrophulariaceae

Nuttallanthus canadensis (L.) D. Sutton
P13141
Penstemon pallidus Small P13145
**Verbascum thapsus* L. P13420

Solanaceae

Physalis heterophylla Nees P13304
Solanum carolinense L. P13318

Ulmaceae

Celtis occidentalis L. P13063
Ulmus americana L. P13056
Ulmus rubra Muhl. P13061

Urticaceae

Parietaria pensylvanica Muhl. P13294

Verbenaceae

Verbena stricta Vent. P13432

Violaceae

**Viola rafinesquii* Greene P13051
Viola sororia Willd. P13069

Vitaceae

Parthenocissus quinquefolia (L.) Planch.
P33307
Vitis riparia Michx. P13151

MONOCOTS**Araceae**

Arisaema triphyllum (L.) Schott P13070

Commelinaceae

Commelina erecta L. P13617
Tradescantia ohioensis Raf. P13163

Cyperaceae

Bulbostylis capillaris (L.) C. B. Clarke
P13410
Carex amphibola Steud. P13171
Carex blanda Dewey E28422
Carex jamesii Schwein. P13166
Carex lurida Wahl. P33289
Carex muhlenbergii Schk. P13172
Carex oligocarpa Schk. P13296
Carex pensylvanica Lam. P13059
Carex scoparia Schk. P35695
Carex stipata Muhl. P35694
Carex tomsa (Fern.) Bickn. P13308
Carex tribuloides Wahl. P33291
Cyperus esculentus L. P13408

Cyperus grayoides Mohlenbr. P13638
Cyperus lupulinus (Spreng.) Marcks P13633
Cyperus schweinitzii Torr. P36151
Cyperus strigosus L. P33288
Eleocharis ovata (Roth) Roem. & Schultes
 P33457
Scirpus atrovirens Willd. P33283

Iridaceae

**Belamcanda chinensis* (L.) DC. P13625
Sisyrinchium albidum Raf. E28415
Sisyrinchium campestre Bickn. P13422

Juncaceae

Juncus acuminatus Michx. P33293
Juncus interior Wieg. P13316
Juncus marginatus Rostk. P33292

Liliaceae

**Allium vineale* L. P35698
 **Asparagus officinalis* L. P13160
Smilacina stellata (L.) Desf. P13170

Poaceae

Agrostis hyemalis (Walt.) BSP. E28424
Andropogon gerardii Vitman P13615
Aristida desmantha Trin. & Ripr. P13567
Aristida tuberculosa Nutt. P13621
Bouteloua hirsuta Lag. P33305
 **Bromus commutatus* Schrad. E28624
 **Bromus inermis* Leyss. P13270
 **Bromus tectorum* L. P13159
Calamovilfa longifolia (Hook.) Scribn.
 P13550
Cenchrus longispinus (Hack.) Fern. P13409
Dichantherium acuminatum (Sw.) Gould &
 Clark P33285
Dichantherium depauperatum (Muhl.)
 Gould E28429
Dichantherium oligosanthes (Schult.) Gould
 P13284

Dichantherium perlongum (Nash) Freckm.
 P13297

Dichantherium villosissimum (Nash)
 Freckm. P13285

**Digitaria ischaemum* (Schreb.) Schreb.
 P13607

**Digitaria sanguinalis* (L.) Scop. P13641

**Echinochloa crus-galli* (L.) Beauv. E30375

Elymus villosus Muhl. P13293

**Eragrostis cilianensis* (All.) Vign. P13429

Eragrostis spectabilis (Pursh) Steud.
 P13551

Eragrostis trichodes (Nutt.) Wood. P13549

Heterostipa spartea (Trin.) Barkworth
 P13271

Koeleria macrantha (Ledeb.) Spreng.
 P13307

Leptoloma cognatum (Schult.) Chase
 P31287

Panicum capillare L. P13434

Panicum virgatum L. P13609

Paspalum bushii Nash P13601

**Poa pratensis* L. P13167

Schizachyrium scoparium (Michx.) Nash
 P13610

**Secale cereale* L. P13322

**Setaria faberi* F. Herzm. P33294

**Setaria viridis* (L.) Beauv. P13435

Sorghastrum nutans (L.) Nash P13630

Sporobolus clandestinus (Biehler) Hitchc.
 P13634

Sporobolus cryptandrus (Torr.) Gray
 P13627

Tridens flavus (L.) Hitchc. P31288

Triplasis purpurea (Walt.) Chapm. P13623

Vulpia octoflora (Walt.) Rydb. E28414

Smilacaceae

Smilax hispida Muhl. P13152

Floristic assessment of the Henry Allan Gleason Nature Preserve, Mason County, Illinois

William E. McClain, Illinois State Museum, Springfield, Illinois 62706

and

Loy R. Phillippe and John E. Ebinger, Center for Biodiversity, Illinois Natural History Survey, Champaign, Illinois 61820

ABSTRACT--The Henry Allan Gleason Nature Preserve is located in the extensive glacial sand deposits associated with the Illinois River in central Illinois. An extensive sand dune is present within the preserve on which undisturbed dry sand prairie, disturbed dry sand prairie, and blow-out communities are present. The undisturbed sand prairie is dominated by *Schizachyrium scoparium* (42% of the importance value [IV]), along with *Tephrosia virginiana*, *Opuntia humifusa*, and *Ambrosia psilostachya*. The disturbed sand prairie is dominated by *Eragrostis trichodes* (24% of the IV), followed by *Heterotheca camporum*, *Ambrosia psilostachya*, and *Rhus aromatica*. Common species in an active blow-out includes *Aristida tuberculosa* and *Cyperus grayioides*, while nearly stabilized blow-outs have a high diversity characterized by *Bouteloua hirsuta*, *Ambrosia psilostachya*, and *Eragrostis trichodes*. A total of 172 plant species were found: 4 fern and fern-allies, 3 gymnosperms, 39 monocots, 126 dicots. Thirty-one non-native species were found, comprising about 18% of the flora. The Floristic Quality Index for the nature preserve is 41.33 when the non-native species are included in the calculations.

INTRODUCTION

At the time of European settlement prairie vegetation covered about 60% of Illinois (Iverson et al. 1991). Though mostly tall-grass prairie, other types of prairie were also common, particularly sand prairies (Schwegman 1973). Sand deposits are common in the northern half of Illinois, accounting for about 5% of the state's land area. Most occur on glacial outwash plains resulting from erosional events associated with Wisconsin glaciation (Willman and Frye 1970, King 1981). One of the most extensive sand deposits is along the Illinois River in the central part of the state (Schwegman 1973). This sand deposit was formed as glacial lakes were drained about 14,500 years ago when glacial moraines were breached, resulting in the Kankakee Torrent. These waters carried huge amounts of sand and gravel that were deposited when the Kankakee Torrent entered the broad lowlands of the Illinois River below present day Hennepin. These sands were subsequently shaped by winds, creating the dune and swale topography known as the Parkland Formation (Willman and Frye 1970, Willman 1973).

The flora of the Illinois River sand deposits was studied by Gleason (1910) in his extensive review of the inland sand deposits of Illinois. Rodgers and Anderson (1979), using General Land Office survey records, examined presettlement vegetation of the area, while Anderson and Brown (1983, 1986) determined effects of fire on sand savannas and adjacent forest. More recently Jenkins et al. (1991) and Coates et al. (1992) studied the composition and structure of two sand forest communities in Mason County. Except for Gleason (1910) little information has been published on the composition of the sand prairie communities of these sand deposits. The present study was undertaken to determine vascular plant species composition, vegetation structure, and floristic quality of the major plant communities of the Henry Allen Gleason Nature Preserve (HAGNP).

STUDY SITE

HAGNP is located in extreme northwestern Mason County, just southeast of the town of Goofy Ridge, and about 15km northeast of Havana, Illinois (SE1/4 S6, NE1/4 S7 T22N R7W). Officially dedicated in 1970, this site lies within the

Illinois River Section of the Mississippi River and Illinois River Sand Area Natural Division, and is within Sand Ridge State Forest (Schwegman 1973). The most significant physiographic feature at HAGNP is a large sand dune known as "Devil's Tower", which comprises about half of the preserve. This dune is more than 25 m tall, and has a large blow-out located near the summit.

When dedicated, much of the 44.5 ha prairie had been heavily disturbed. The entire site was designated "grade C" dry sand prairie by the Illinois Natural Areas Inventory, which indicates that the site has been disturbed, though some small remnants of original sand prairie still existed (White 1978). Most of the large dune had been planted in pine trees. When these were removed in 1978 many were 7 m tall with the diameter at breast height ranging from 20-25 cm. The first prescribed burn on the preserve was in the spring of 1980 with successive burns in 1982, 1993, 1997, and 1999. The soils of the preserve are excessively drained Plainfield sands (Calsyn 1995).

Climate at the HAGNP is continental with warm summers and cold winters. Based on weather data from Havana, 15km to the southwest, mean annual precipitation is 96.0cm, with May having the highest rainfall (11.3cm). Mean annual temperature is 10.8°C with the hottest month being July (average of 24.6°C), and the coldest January (average of -5.0°C). Frost-free days range from 140 to 206, with the average being 173 day per year (Midwestern Regional Climate Center 2002).

METHODS

The HAGNP was visited throughout the growing seasons of 1998 to 2002 and voucher specimens of each plant species were collected, identified, and deposited in the Stover-Ebinger Herbarium of Eastern Illinois University, Charleston, Illinois (EIU). Criteria for designating non-native species follows Mohlenbrock (1986) and Gleason and Cronquist (1991), while nomenclature follows Mohlenbrock (1986).

Within the undisturbed dry sand prairie, the disturbed dry sand prairie, and an active blow-out, line transects 25m long were placed, four in each of the three community types. The transects, surveyed in September of 2000, were stratified across each community, and oriented so that it occurred entirely within a single community type. In addition, three partially stabilized blow-outs having some areas of shifting sand were studied. Due to their small size, two 15 m transects were randomly placed in each. Along each transect, m^2 (1m x 1m) quadrats were located at 1m intervals ($n=25$ /transect), except in the partially stabilized blowout communities where $1/4m^2$ (50cm x 50cm) quadrats were used ($n=15$ /transect). Odd-numbered quadrats were located on the right side of the transect line, even-numbered quadrats on the left. Cover for each species was determined by using the Daubenmire canopy cover class system (Daubenmire 1959) as modified by Bailey and Poulton (1968) (class 1 = 0 to 1%; class 2 = 2 to 5%; class 3 = 6 to 25%; class 4 = 26 to 50%; class 5 = 51 to 75%; class 6 = 76 to 95%; class 7 = 96 to 100%). Importance value (IV) was determined by summing relative cover and relative frequency (total IV = 200), while average cover was determined by dividing the total cover of each species by the number of plots surveyed in each community type.

The Floristic Quality Index (FQI) was determined using the coefficient of conservatism (CC) assigned to each species (Taft et al. 1997). The CC for each taxon was determined by assigning each an integer from 0 to 10 based on the species tolerance to disturbance and its fidelity to habitat integrity. This index provides a measure of the floristic integrity or degree of disturbance to a site. The FQI is a weighted index of species richness (N), and is the arithmetic product of the mean CC, multiplied by the square-root of the species richness (\sqrt{N}) of an inventory site: $FQI = \text{mean CC}(\sqrt{N})$. For relatively small areas, FQI gives a rapid means of comparison and an indication of the floristic integrity of the site. When used along with other floristic measures, such as

quadrat-based sampling methods, it provides a method of making quantitative comparisons between sites. Prairies with an FQI of 35 or higher are considered good quality natural areas (Taft et al. 1997).

RESULTS AND DISCUSSION

A total of 172 vascular plant species representing 137 genera and 56 families were documented from the 44.5 ha preserve (Appendix I). Fern, fern-allies and gymnosperms were represented by only seven species. Of the remainder, 39 were monocots in 5 families and 28 genera, and 126 were dicots in 46 families and 103 genera. Thirty-one non-native species were found, and 31 species of trees and shrubs were collected. The predominant plant families were the Asteraceae and Poaceae, each with 28 species. The Illinois endangered *Lesquerella ludoviciana* and the Illinois threatened *Cyperus grayioides* were found on the preserve (Herkert and Ebinger (2002).

Active blow-out community: Only 12 plant species were encountered in the active blow-out. Average bare ground and litter accounted for 83% of the cover. *Aristida tuberculosa* was the dominant species, followed by *Cyperus grayioides*, and *Diodia teres* (Table 1). Only eight species were encountered in the plots and all but three had IV's lower than 10.

Partially stabilized blow-out communities: Three small, nearly stabilized blow-outs were examined, two located near the top of Devil's Tower, the third near the top of a low ridge in the southwestern part of the preserve. On these partially stabilized blow-outs bare ground and litter cover averaged 52% (Table 2). All had a similar flora and shared dominant species with at least three species with highest IVs in each blow-out occurring among the top five species when the data were combined (Table 2). *Bouteloua hirsuta* was usually the dominant species, followed by *Ambrosia psilostachya*, and commonly *Eragrostis trichodes*, while *Lesquerella ludoviciana* was an important component. In Illinois this endangered species is known only from the HAGNP where it occurs only in these three partially stabilized blow-out communities.

Disturbed dry sand prairie: Gleason (1910) indicated that Devil's Tower was originally covered with prairie but most had been destroyed by cultivation and pasturing, only a few small areas remained in their natural condition. Later the slopes of this dune were planted in pines that were removed in 1978. Presently these slopes contain a disturbed dry sand prairie having a high species diversity including some taxa commonly associated with mature sand prairies: *Schizachyrium scoparium*, *Opuntia humifusa*, and *Dichanthelium villosissimum*. In this disturbed sand prairie *Eragrostis trichodes* dominated followed by *Heterotheca camporum*, *Ambrosia psilostachya*, and *Rhus aromatica* (Table 3).

Mature dry sand prairie: Mature dry sand prairies are located on the east and northwest sides of the preserve on the lower flanks of Devil's Tower. Rarely exceeding 100 m by 75 m in size, these areas contain many species common to dry sand prairies (Gleason 1910, White and Madany 1978). *Schizachyrium scoparium* was the leading dominant. *Tephrosia virginiana* and *Opuntia humifusa* were second and third in IV respectively, while *Ambrosia psilostachya* and *Dichanthelium villosissimum* also had IVs that exceeded 15 (Table 4).

This dry sand prairie, referred to as the bunch-grass association by Gleason (1910), was dominated by many clumps of *Schizachyrium scoparium* that were 20-60 cm across, nearly circular in outline, and formed a dense mass through which few other plant species could grow. Some larger clumps had centers that had died, forming rings in which no other species were found. *Dichanthelium villosissimum* has a similar growth pattern but formed smaller clumps. Many of

the forbs, particularly *Ambrosia psilostachya*, *Commelina erecta*, and *Conyza canadensis* grew in the spaces between the clumps. The bunch-grass community described by Gleason (1910) is very similar to the undisturbed dry sand prairie community found during the present study.

Blowing sand was a common characteristic of the sand communities. Even on relatively calm days there was moving sand in the blow-outs. Here vegetation was sparse, and most of the area was exposed sand (bare ground cover of 83%). In the nearly stabilized blow-outs, ground cover was about 50%. Where bunch grasses were common, however, the amount of open sand decreased significantly (bare ground cover of 35%).

The Floristic Quality Index for the HAGNP was 41.33 with 31 non-native species included in the calculations. Removing these exotic species from the calculations increased the FQI to 45.64 with a mean coefficient of conservatism of 3.84. Of the native species encountered, 13 had a coefficient of conservatism of 8 or higher and were relatively common components of the flora. In contrast, non-native taxa were restricted to the margins of the preserve, in areas of human disturbance, and were not common components of study sites. None were encountered in the blow-out community (Table 1), or in the mature dry sand prairie (Table 4). In the stabilized blow-out community the only non-native was *Salsola iberica* (Table 2), while the only exotics in the disturbed dry sand prairie were *Poa pratensis* and *Helianthus petiolaris*, both being uncommon (Table 3).

Present management practices on the HAGNP consists of occasional fires to control wood encroachment. These burns have decreased litter accumulation and appear to have been helpful in increasing prairie quality and decreasing the number of exotic species on the prairie. Presently both the undisturbed and disturbed sand prairie communities contain many native sand prairie taxa and very few exotics. Though not apparent from this study, woody species are becoming common in parts of the preserve and more frequent fires will be needed to stop woody encroachment.

ACKNOWLEDGMENTS

The authors would like to thank Dr. Gordon Tucker, Eastern Illinois University, for help with identification of the Cyperaceae, and John Wilker, Natural Heritage Biologist, Illinois Department of Natural Resources, for his help and advice.

LITERATURE CITED

- Anderson, R.C. and L.E. Brown. 1983. Comparative effects of fire on trees in a midwestern savannah and an adjacent forest. *Bull. Torrey Bot. Club* 110:87-90.
- Anderson, R.C. and L.E. Brown. 1986. Stability and instability in plant communities following fire. *Amer. J. Bot.* 73:364-368.
- Bailey, A.W. and C.E. Poulton. 1968. Plant communities and environmental relationships in a portion of the Tillamook burn, northwestern Oregon. *Ecology* 49:1-13.
- Calsyn, D.E. 1995. Soil survey of Mason County, Illinois. Soil Report 146. University of Illinois Agricultural Experiment Station, Urbana, Illinois.
- Coates, D.T., S.E. Jenkins, and J.E. Ebinger. 1992. Woody vegetation survey of Barkhausen Woods, a closed canopy sand forest in Mason County, Illinois. *Erigenia* 12:1-6.
- Daubenmire, R. 1959. A canopy coverage method of vegetation analysis. *Northwest Sci.* 33:43-64.

Gleason, H.A. 1910. The vegetation of the inland sand deposits of Illinois. Bull. Illinois State Laboratory Nat. Hist. 9:21-174.

Gleason, H.A. and A. Cronquist. 1991. Manual of the vascular flora of northeastern United States and adjacent Canada. Second Edition. The New York Botanical Garden, Bronx, New York.

Herkert, J.R. and J.E. Ebinger. 2002. editors. Endangered and threatened species of Illinois: Status and distribution. Volume 1. Plants. Illinois Endangered Species Protection Board, Springfield, Illinois.

Iverson, L.R., G.L. Rolfe, T.J. Jacob, A.S. Hodgins, and M.R. Jeffords. 1991. Forests of Illinois. Illinois Council on Forest development, Urbana, and Illinois Natural History Survey, Champaign, Illinois.

Jenkins, S.E., J.E. Ebinger, and W.E. McClain. 1991. Woody vegetation survey of Bishop's Woods, a sand forest in Mason County, Illinois. Trans. Illinois State Acad. Sci. 84:20-27.

King, J.E. 1981. Late Quaternary vegetational history of Illinois. Ecol. Monogr. 51:43-62.

Midwestern Regional Climate Center. 2002. <http://mcc.sws.uiuc.edu>

Mohlenbrock, R.H. 1986. Guide to the vascular flora of Illinois, revised and enlarged edition, Southern Illinois University Press, Carbondale, Illinois.

Rodgers, C.S. and R.C. Anderson. 1979. Presettlement vegetation of two prairie peninsula counties. Botanical Gazette 140:232-240.

Schwegman, J.E. 1973. Comprehensive plan for the Illinois nature preserves system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, Illinois.

Taft, J.B., G.S. Wilhelm, D.M. Ladd, and L.A. Masters. 1997. Floristic quality assessment for vegetation in Illinois, a method for assessing vegetation integrity. *Erigenia* 15:1-95.

White J. 1978. Illinois natural areas inventory technical report. Volume I. Survey methods and results. Illinois Natural Areas Inventory, Urbana, Illinois.

White, J. and M.H. Madany. 1978. Classification of natural communities in Illinois. p. 310-405. In: White, J. (ed.) Illinois natural areas inventory technical report. Illinois Natural Areas Inventory, Urbana, Illinois.

Willman, H.B. 1973. Geology along the Illinois waterway - a basis for environmental planning. Illinois State Geological Survey Circular 478. Urbana, Illinois.

Willman, H.B. and J.C. Frye. 1970. Pleistocene stratigraphy of Illinois. Illinois State Geological Survey Bull. 94:1-204.

Table 1. Frequency (%), average cover (% of total area), relative frequency, relative cover, and importance value of ground layer species in an active blow-out at Henry Allan Gleason Nature Preserve, Mason County, Illinois.

Species	Freq.	Average Cover	Rel. Freq.	Rel. Cover	I. V.
<i>Aristida tuberculosa</i>	86	5.88	49.6	45.9	95.5
<i>Cyperus grayioides</i>	56	4.55	32.4	35.6	68.0
<i>Diodia teres</i>	14	0.85	8.1	6.7	14.8
<i>Panicum virgatum</i>	4	0.60	2.3	4.7	7.0
<i>Rhus aromatica</i>	2	0.53	1.2	4.1	5.3
<i>Dichanthelium villosissimum</i>	4	0.34	2.3	2.6	4.9
<i>Crotonopsis linearis</i>	6	0.03	3.5	0.2	3.7
<i>Ambrosia psilostachya</i>	1	0.03	0.6	0.2	0.8
Totals		12.81	100.0	100.0	200.0
Average bare ground and litter		83.75			

Table 2. Frequency (%), average cover (% of total area), relative frequency, relative cover, and importance value of ground layer species in stabilized blow-outs at Henry Allan Gleason Nature Preserve, Mason County, Illinois.

Species	Freq.	Average Cover	Rel. Freq.	Rel. Cover	I. V.
<i>Bouteloua hirsuta</i>	72	8.92	11.9	22.6	34.5
<i>Ambrosia psilostachya</i>	80	6.33	13.2	16.0	29.2
<i>Eragrostis trichodes</i>	67	5.69	11.0	14.4	25.4
<i>Aster oblongifolius</i>	31	3.57	5.2	9.1	14.3
<i>Rhus aromatica</i>	19	3.55	3.1	9.0	12.1
<i>Lesquerella ludoviciana</i>	43	1.04	7.2	2.6	9.8
<i>Koeleria macrantha</i>	21	2.23	3.5	5.6	9.1
<i>Oenothera rhombipetala</i>	40	0.37	6.6	0.9	7.5
<i>Opuntia humifusa</i>	22	1.49	3.7	3.8	7.5
<i>Heterotheca camporum</i>	17	1.03	2.8	2.6	5.4
<i>Paspalum bushii</i>	20	0.76	3.3	1.9	5.2
<i>Chamaesyce geyeri</i>	26	0.21	4.2	0.5	4.7
<i>Conyza canadensis</i>	22	0.38	3.7	1.0	4.7
<i>Commelina erecta</i>	20	0.35	3.3	0.9	4.2
<i>Calamovilfa longifolia</i>	20	0.24	3.3	0.6	3.9
<i>Schizachyrium scoparium</i>	8	1.03	1.3	2.6	3.9
<i>Brickellia eupatorioides</i>	7	0.73	1.1	1.9	3.0
<i>Cassia fasciculata</i>	14	0.13	2.4	0.3	2.7
<i>Phlox bifida</i>	10	0.24	1.7	0.6	2.3
<i>Liatris aspera</i>	8	0.28	1.3	0.7	2.0
<i>Leptoloma cognatum</i>	6	0.41	0.9	1.0	1.9
<i>Croton glandulosa</i>	9	0.04	1.5	0.1	1.6
<i>Aster ericoides</i>	4	0.21	0.7	0.5	1.2
<i>Asclepias viridiflora</i>	4	0.05	0.7	0.1	0.8
<i>Lespedeza capitata</i>	1	0.17	0.2	0.4	0.6
<i>Dichanthelium villosissimum</i>	2	0.07	0.4	0.2	0.6
<i>Poinsettia dentata</i>	3	0.02	0.6	--	0.6
<i>Salsola iberica</i>	2	0.01	0.4	--	0.4
<i>Teucrium canadense</i>	2	0.01	0.4	--	0.4
<i>Strophostyles helvola</i>	1	0.03	0.2	0.1	0.3
<i>Euphorbia corollata</i>	1	0.01	0.2	--	0.2
Totals		39.60	100.0	100.0	200.0
Average bare ground and litter		52.40			

Table 3. Frequency (%), average cover (% of total area), relative frequency, relative cover, and importance value of ground layer species in a disturbed sand prairie at Henry Allan Gleason Nature Preserve, Mason County, Illinois.

Species	Freq.	Average Cover	Rel. Freq.	Rel. Cover	I. V.
<i>Eragrostis trichodes</i>	98	16.15	21.0	26.8	47.8
<i>Heterotheca camporum</i>	67	17.64	14.3	29.2	43.5
<i>Ambrosia psilostachya</i>	82	9.51	17.6	15.7	33.3
<i>Rhus aromatica</i>	36	9.20	7.7	15.2	22.9
<i>Opuntia humifusa</i>	64	3.28	13.7	5.4	19.1
<i>Commelina erecta</i>	28	0.83	6.0	1.4	7.4
<i>Conyza canadensis</i>	12	0.53	2.8	0.9	3.7
<i>Calamovilfa longifolia</i>	10	0.56	2.1	0.9	3.0
<i>Sporobolus clandestinus</i>	7	0.69	1.5	1.1	2.6
<i>Dichanthelium oligosanthes</i>	8	0.34	1.7	0.6	2.3
<i>Aster oblongifolius</i>	4	0.60	0.9	1.0	1.9
<i>Dichanthelium villosissimum</i>	5	0.13	1.1	0.2	1.3
<i>Oenothera rhombipetala</i>	4	0.19	0.9	0.3	1.2
<i>Physalis heterophylla</i>	4	0.12	0.9	0.2	1.1
<i>Schizachyrium scoparium</i>	4	0.10	0.9	0.2	1.1
<i>Croton glandulosus</i>	4	0.05	0.9	0.1	1.0
<i>Poa pratensis</i>	4	0.05	0.9	0.1	1.0
<i>Aristida tuberculosa</i>	4	0.02	0.9	--	0.9
<i>Euphorbia corollata</i>	3	0.07	0.6	0.1	0.7
<i>Helianthus petiolaris</i>	3	0.18	0.4	0.3	0.7
<i>Sporobolus cryptandrus</i>	3	0.04	0.6	0.1	0.7
<i>Liatris aspera</i>	2	0.04	0.4	0.1	0.5
<i>Paspalum bushii</i>	2	0.06	0.4	0.1	0.5
<i>Cassia fasciculata</i>	2	0.01	0.4	--	0.4
<i>Chamaesyce geyeri</i>	2	0.01	0.4	--	0.4
<i>Asclepias verticillata</i>	1	0.01	0.2	--	0.2
<i>Chenopodium desiccatum</i>	1	0.01	0.2	--	0.2
<i>Cyperus schweinitzii</i>	1	0.01	0.2	--	0.2
<i>Lactuca canadensis</i>	1	0.01	0.2	--	0.2
<i>Triplasis purpurea</i>	1	0.01	0.2	--	0.2
Totals		60.45	100.0	100.0	200.0
Average bare ground and litter		47.98			

Table 4. Frequency (%), average cover (% of total area), relative frequency, relative cover, and importance value of ground layer species in a mature sand prairie at Henry Allan Gleason Nature Preserve, Mason County, Illinois.

Species	Freq.	Average Cover	Rel. Freq.	Rel. Cover	I. V.
<i>Schizachyrium scoparium</i>	97	37.06	25.5	59.1	84.6
<i>Tephrosia virginiana</i>	43	10.03	11.3	16.0	27.3
<i>Opuntia humifusa</i>	66	4.88	17.3	7.8	25.1
<i>Ambrosia psilostachya</i>	58	2.27	15.2	3.6	18.8
<i>Dichanthelium villosissimum</i>	50	3.37	13.1	5.4	18.5
<i>Rhus aromatica</i>	7	2.88	1.8	4.6	6.4
<i>Eragrostis trichodes</i>	11	1.40	2.9	2.2	5.1
<i>Commelina erecta</i>	13	0.14	3.4	0.2	3.6
<i>Panicum virgatum</i>	10	0.27	2.6	0.4	3.0
<i>Crotonopsis linearis</i>	10	0.05	2.6	0.1	2.7
<i>Conyza canadensis</i>	5	0.08	1.3	0.1	1.4
<i>Leptoloma cognatum</i>	3	0.07	0.8	0.1	0.9
<i>Stipa spartea</i>	2	0.18	0.5	0.3	0.8
<i>Aristida tuberculosa</i>	3	0.02	0.8	--	0.8
<i>Eragrostis spectabilis</i>	1	0.03	0.3	0.1	0.4
<i>Koeleria macrantha</i>	1	0.01	0.3	--	0.3
<i>Cyperus schweinitzii</i>	1	0.01	0.3	--	0.3
Totals		62.75	100.0	100.0	200.0
Average bare ground and litter		35.44			

APPENDIX I. Vascular species encountered at Henry Allen Gleason Nature Preserve, Mason County, Illinois, listed alphabetically by family under the major plant groups. An asterisk indicates non-native (exotic) species (*). For each species the author's collecting number (JEE) is given.

FERN AND FERN-ALLIES

Aspleniaceae

- Asplenium platyneuron* (L.) Oakes 28399
Woodsia obtusa (Spreng.) Torr. 29639

Equisetaceae

- Equisetum hyemale* L. 28243

Ophioglossaceae

- Botrychium virginianum* (L.) Sw. 29640

GYMNOSPERMES

Cupressaceae

- Juniperus virginiana* L. 28146

Pinaceae

- **Pinus banksiana* Lamb. 28646
 **Pinus strobus* L. 30366

MONOCOTS

Commelinaceae

- Commelina erecta* L. 28131
Tradescantia ohiensis Raf. 28145

Cyperaceae

- Carex meadii* Dewey 25791
Carex muhlenbergii Willd. 28110
Carex tonsa (Fern.) Bickn. 29223
Cyperus filiculmis Vahl. 25290
Cyperus grayioides Mohlenbr. 25291
Cyperus schweinitzii Torr. 25793

Liliaceae

- **Allium vineale* L. 28650
Smilacina stellata (L.) Desf. 28323

Poaceae

- Agrostis hyemalis* (Walt.) BSP. 25790
Aristida desmantha Trin. & Rupr. 28237
Aristida purpurascens Poir. 28239
Aristida tuberculosa Nutt. 28238
Bouteloua hirsuta Lag. 25293
 **Bromus commutatus* Schrad. 28655
 **Bromus inermis* Leyss. 30186
 **Bromus tectorum* L. 28400
Calamovilfa longifolia (Hook.) Scribn. 28124
Dichanthelium oligosanthos Schult. 25794
Dichanthelium villosissimum Nash 28128
 **Digitaria sanguinalis* (L.) Scop. 30902
Elymus canadensis L. 28121
Eragrostis spectabilis (Pursh) Steud. 28140
Eragrostis trichodes (Nutt.) Wood 28136

Koeleria macrantha (Ledeb.) Spreng. 25792
Leptoloma cognatum (Schult.) Chase 28154
Panicum virgatum L. 28114
Paspalum bushii Nash 25292
 **Poa pratensis* L. 25800
Schizachyrium scoparium (Michx.) Nash 28241
Sorghastrum nutans (L.) Nash 28240
Sporobolus clandestinus (Biehler) Hitchcock 28236
Sporobolus cryptandrus (Torr.) Gray 29417
Stipa spartea Trin. 25796
Tridens flavus (L.) Hitchcock 28143
Triplasis purpurea (Walt.) Chapm. 28235
Vulpia octoflora (Walt.) Rydb. 28401

Smilacaceae

Smilax hispida Muhl. 29416

DICOTS

Amaranthaceae

Froelichia floridana (Nutt.) Moq. 25289

Anacardiaceae

Rhus aromatica Ait. 28117

Rhus glabra L. 29414

Toxicodendron radicans (L.) Kuntze 28397

Apiaceae

Osmorhiza longistylis (Torr.) DC. 29642

Sanicula canadensis L. 28652

Apocynaceae

Apocynum sibiricum Jacq. 28649

Asclepiadaceae

Asclepias syriaca L. 28644

Asclepias verticillata L. 28123

Asclepias viridiflora Raf. 25799

Asteraceae

**Achillea millefolium* L. 28657

Ambrosia artemisiifolia L. 28148

Ambrosia psilostachya DC. 28147

Aster ericoides L. 28136

Aster oblongifolius Nutt. 28250

Aster pilosus Willd. 28251

Bidens bipinnata L. 28141

Brickellia eupatorioides (L.) Shinnars 28139

Cirsium discolor (Muhl.) Spreng. 29228

Conyza canadensis (L.) Cronq. 28246

Coreopsis lanceolata L. 25795

Coreopsis palmata Nutt. 28144

Erigeron annuus (L.) Pers. 28647

Erigeron stigosus Muhl. 25803

Eupatorium rugosum Houtt. 28149

Eupatorium serotinum Michx. 28247

Gnaphalium obtusifolium L. 28248

**Helianthus petiolaris* Nutt. 25287

Heterotheca camporum (Greene) Shinnars 28127
Krigia virginica (L.) Willd. 25788
Lactuca canadensis L. 28158
 **Lactuca serriola* L. 29227
Liatris aspera Michx. 28249
Rudbeckia hirta L. 28645
Senecio plattensis Nutt. 28326
Solidago canadensis L. 29415
Solidago nemoralis Ait. 28142
 **Tragopogon dubius* Scop. 25784

Boraginaceae

Hackelia virginiana (L.) I. M. Johnston 30903
Lithospermum caroliniense (J. F. Gmel.) MacM. 25789

Brassicaceae

**Alliaria petiolata* (Bieb.) Cavara & Grande 28407
Arabis canadensis L. 28402
 **Lepidium densiflorum* Schrad. 30854
Lepidium virginicum L. 28126
Lesquerella ludoviciana (Nutt.) S. Wats. 27791

Cactaceae

Opuntia humifusa (Raf.) Raf. 28129

Caesalpiniaceae

Cassia fasciculata Michx. 18132
Gymnocladus dioicus (L.) K. Koch 28818

Campanulaceae

Campanula americana L. 28159
Triodanis perfoliata (L.) Nieuwl. 25787

Caryophyllaceae

**Holosteum umbellatum* L. 28321
 **Saponaria officinalis* L. 25803
Silene antirrhina L. 25801

Celastraceae

Celastrus scandens L. 28157
Euonymus atropurpurea Jacq. 28327

Chenopodiaceae

Chenopodium desiccatum A. Nels. 28118
Cycloloma atriplicifolium (Spreng.) Coult. 25288
 **Salsola iberica* Sennen & Pav. 28134

Cistaceae

Helianthemum canadense (L.) Michx. 28244

Convolvulaceae

**Ipomoea hederacea* (L.) Jacq. 29226
Ipomoea lacunosa L. 29225

Cornaceae

Cornus drummondii C.A. Mey. 30188

Euphorbiaceae

- Chamaesyce geyeri* (Engelm. & Gray) Small 28160
Croton glandulosus L. 25286a
Crotonopsis linearis Michx. 28116
Euphorbia corollata L. 28122
Poinsettia dentata (Michx.) Kl. & Garcke 25286b

Fabaceae

- Desmodium sessilifolium* (Torr.) Torr. & Gray 30367
Lespedeza capitata Michx. 28242
 **Melilotus alba* Medic. 30189
 **Melilotus officinalis* (L.) Pallas 28398
 **Robinia pseudoacacia* L. 28651
Strophostyles helvola (L.) Ell. 30368
Tephrosia virginiana (L.) Pers. 28115

Fagaceae

- Quercus x bushii* Sarg. 28112
Quercus marilandica Muenchh. 28156
Quercus velutina Lam. 28643

Geraniaceae

- Geranium carolinianum* L. 28403

Grossulariaceae

- Ribes missouriense* Nutt. 30190

Juglandaceae

- Juglans nigra* L. 28648

Lamiaceae

- Monarda punctata* L. 28819
Physostegia virginiana (L.) Benth. 30369
Teucrium canadense L. 28119

Malvaceae

- Callirhoe triangulata* (Leavenw.) Gray 28155

Menispermaceae

- Menispermum canadense* L. 28816

Molluginaceae

- **Mollugo verticillatus* L. 28658

Moraceae

- **Cannabis sativa* L. 30192
 **Maclura pomifera* (Raf.) Schneider 30193
 **Morus alba* L. 28408

Nyctaginaceae

- **Mirabilis nyctaginea* (Michx.) MacM. 25797

Onagraceae

- ~~*Oenothera biennis* L. 30904~~
Oenothera rhombipetala Nutt. 28133

Oxalidaceae

- Oxalis stricta* L. 28659
- Papaveraceae
Corydalis micrantha (Engelm.) Gray 28410
- Phytolaccaceae
Phytolacca americana L. 28150
- Plantaginaceae
 **Plantago patagonica* Jacq. 25785
- Polemoniaceae
Phlox bifida Beck 28135
- Polygonaceae
Polygonum cristatum Engelm. & Gray 28120
Polygonum tenue Michx. 29229
 **Rumex acetosella* L. 28153
- Portulacaceae
Talinum rugospermum Holz. 29992
- Ranunculaceae
Anemone caroliniana Walt. 25208
-
- Rosaceae
Fragaria virginiana Duchesne 28245
Geum canadense Jacq. 28653
Malus ioensis (Wood) Britt. 30187
Prunus angustifolia Marsh. 28151
Prunus serotina Ehrh. 28656
Rosa carolina L. 28654
 **Rosa multiflora* Thunb. 28404
Rubus allegheniensis Porter 28406
Rubus occidentalis L. 28405
- Rubiaceae
Diodia teres Walt. 28130
Galium aparine L. 28660
- Rutaceae
Ptelea trifoliata L. 28152
Zanthoxylum americanum Mill. 28125
- Scrophulariaceae
Linaria canadensis (L.) Dum.-Cours. 25786
Penstemon pallidus Small 28114
 **Veronica arvensis* L. 29641
- Solanaceae
Physalis heterophylla Nees. 25798
Solanum carolinense L. 29224
-
- Verbenaceae
Verbena stricta Vent. 28111
Verbena urticifolia L. 30905

Violaceae


Viola pedata L. 28324

**Viola rafinesquii* Greene 28322

Vitaceae

Parthenocissus quinquefolia (L.) Planch. 28817

Vitis vulpina L. 28410



Sand prairie communities of Matanzas Nature Preserve, Mason County, Illinois

Mary Ann Feist, Marilyn J. Morris, Loy R. Phillippe, John E. Ebinger, Illinois
Natural History Survey, Champaign, Illinois 61920

and

William E. McClain, Illinois State Museum, Springfield, Illinois 62706

ABSTRACT--The Matanzas Prairie Nature Preserve is located in the extensive glacial sand deposits associated with the Illinois River near Bath, Mason County, Illinois. This preserve contains the only remaining high quality wet-mesic sand prairie and sedge meadow associated with the Illinois River sand deposits. The sedge meadow, about 5 ha in size, was dominated by *Carex stricta* (IV of 66.6 out of 200), *Calamagrostis canadensis*, and *Rosa palustris* (both with IV's of 28.3). The wet-mesic sand prairie, occurs on slightly higher ground, and was dominated by *Solidago canadensis*, *Andropogon gerardii*, *Carex* spp., *Poa pratensis*, and *Euthamia graminifolia*. In some parts of the preserve it gradually changes to shrub prairie with a similar ground layer. A total of 342 species of vascular plants species were encountered on the preserve; 5 fern and fern-allies, 100 monocots, and 237 dicots. Except for *Poa pratensis*, which was among the dominant species in the wet-mesic sand prairie and shrub prairie, the 38 exotic species were rarely encountered (11% of the flora). The Floristic Quality Index for the sedge meadow was _____, the wet-mesic sand prairie _____, and the shrub prairie _____.

INTRODUCTION

At the time of European settlement prairie vegetation covered about 60% of Illinois (Iverson et al. 1991). Most was black soil, tall-grass prairie in the prairie peninsula, though sand prairie was also relatively common (Transeau 1935, Schwegman 1973). In the northern half of Illinois extensive sand areas exist on glacial outwash plains associated with erosional events of the Wisconsin glaciation (Willman and Frye 1970, King 1981). The two most extensive are the Kankakee sand deposits in northeastern Illinois, and the Illinois River sand deposits in the central part of the state (Schwegman 1973). Both were formed about 14,500 years ago as a result of depositional events associated with the Kankakee Torrent (Willman and Frye 1970, Willman 1973).

The flora of the Illinois River sand deposits were first studied by Gleason (1910) in his extensive review of the inland sand deposits of Illinois. Rodgers and Anderson (1979) examined presettlement vegetation of the area, while Anderson and Brown (1983, 1986) determined effects of fire on sand savannas and adjacent forest. Later Jenkins et al. (1991) and Coates et al. (1992) studied the composition and structure of sand forest communities. Recently a study of the flora of a dry sand prairie was completed (McClain et al. 2004). As the Matanzas Prairie Nature Preserve contains the only preserved remnant of sedge meadow and wet-mesic sand prairie in the Illinois River sand deposits, it was decided to examine the vascular plant species composition, vegetation structure, and floristic quality of the native plant communities in this preserve.

DESCRIPTION OF THE STUDY SITE

The Matanzas Prairie Nature Preserve is located in southwestern Mason County, 2 km northeast of Bath, and 10 km south of Havana, Illinois (NE1/4 S4 T20N R9W). Officially dedicated in 1985, this site lies within the Illinois

River Section of the Mississippi River and Illinois River Sand Area Natural Division (Schwegman 1973). The preserve contains high quality wet-mesic sand prairie, shrub prairie, and sedge meadow communities that occur in only a few other places in Illinois. These communities have been assigned a rarity index of 5 (1 common, 5 very rare) by the Illinois Natural Areas Inventory (White 1978). The preserve is 23 ha in size, the northern part of about 11 ha contains high quality prairie and sedge meadow remnants, the southern part, 12 ha in size, is cultivated land that has been fallow since 1986.

In the Illinois River sand deposits, these wet area communities mostly occurred in the low-lying portions of out-wash and lake-plains. Most of these communities were drained and converted to agricultural lands. A few attempts were made to drain the preserve, and a few channels are still obvious. The large ditch that divides the preserve into two parts has been dammed by Beavers, and is usually filled with water. The northern part of the preserve was designated "grade B" wet sand prairie and sedge meadow by the Illinois Natural Areas Inventory (White 1978).

The soils of the preserve are wet Plainfield sands high in organic matter (Calsyn 1995). Climate is continental with warm summers and cold winters. Based on weather data from Havana, 10 km to the north, mean annual precipitation is 96.0 cm, with May having the highest rainfall (11.3 cm). Mean annual temperature is 10.8°C with the hottest month being July (average of 24.6°C), and the coldest January (average of -5.0°C). Frost free days range from 140 to 206, with the average being 173 day per year (Midwestern Regional Climate Center 2002).

METHODS

The Matanzas Prairie Nature Preserve was visited throughout the growing season of 1991 and at various other times between 1998 and 2002. During these visits the extent of the plant communities was determined and habitat data for each taxon noted. Voucher specimens of each plant species were collected, identified, and deposited in the herbarium of the Illinois Natural History Survey, Champaign, Illinois (ILLS). Criteria for designating non-native species followed Fernald (1950), Mohlenbrock (1986), and Gleason and Cronquist (1991). Nomenclature follows Mohlenbrock (1986).

Transects 25 m long were located randomly in cardinal compass directions throughout each of three community types: sedge meadow, wet-mesic sand prairie, and shrub prairie. The sedge meadow was studied in July 1999 and again in September 2001 using two transects each time (50 plots each year), the wet-mesic sand prairie was examined in September 1999 using four transects (100 plots), while the shrub prairie was surveyed in September 2001 using two transects (50 plots). Along each transect, $\frac{1}{4}$ m² quadrates were located at 1 m intervals, odd numbered quadrates on the right, even numbered quadrates on the left. Cover for each species was determined by using the Daubenmire cover class system (Daubenmire 1959) as modified by Bailey and Poulton (1968). Importance value (IV) for ground layer species was determined by summing relative cover and relative frequency.

The Floristic Quality Index (FQI) for each of the three communities was determined by using the coefficient of conservatism (CC) assigned to each species based on the species tolerance to disturbance and its fidelity to habitat integrity (Taft et al. 1997). As used here, the FQI is a weighted index of species richness (N), and is the arithmetic product of the mean CC, multiplied by the square-root of the species richness (\sqrt{N}) of a sites: $FQI = \text{mean CC}(\sqrt{N})$. For relatively small areas the FQI gives an indication of the

floristic integrity of the site. When used along with other floristic measures, such as quadrat-based sampling methods, it is useful in making site comparisons. Prairie sites with an FQI of 35 or higher are considered high quality natural areas (Taft et al. 1997).

RESULTS

A total of 342 species representing 194 genera and 71 families were documented for the site (Appendix I). Fern and fern-allies were represented by only five species. Of the remainder, 100 were monocots in 11 families and 47 genera, and 237 were dicots in 56 families and 143 genera. Of these taxa, 38 were not native to Illinois, and except for *Poa pratensis*, most were uncommon, only occasionally being encountered. One Illinois endangered species, *Platanthera flava* var. *herbiola*, and one Illinois threatened species, *Tomanthera auriculata* were found (Herkert and Ebinger 2002). The largest families were the Asteraceae (45 species), Poaceae (44), Cyperaceae (36), Polygonaceae (15), and Rosaceae (14).

Plant community types were determined according to the Illinois Natural Areas Inventory (White and Madany 1978). In addition to extensive disturbed and cultural communities, three communities with high to relatively high natural quality were designated: sedge meadow, wet-mesic sand prairie, and shrub prairie. All were located in the northern half of the preserve and show a gradual transition from one to the other depending on elevation and water availability.

Sedge Meadow Community: This community occupied the eastern half of the northern part of the preserve. Surface water was present during the winter and spring and the soil was nearly always saturated. In the 1999 survey the sedge meadow was dominated by *Carex stricta* (IV of 66.6), *Calamagrostis canadensis* and *Rosa palustris* (both with IV's of 28.3) (Table 1). Only 18 other taxa were present, all native species, and all with relatively low importance values. Numerous colonies of *Rosa palustris* existed throughout the sedge meadow, accounting for nearly one quarter of the entire area. In these colonies few other species occurred. The sedge meadow was burned in the fall of 2001. In the 2001 survey major changes in the flora of the sedge meadow were observed (Table 1). There was a dramatic decrease in *Rosa palustris* and a switch in dominance to *Calamagrostis canadensis* (IV of 69.1). This species was especially robust, covering over other species in some areas. In contrast, the *Rosa palustris* had been top-killed, but was resprouting. More species were encountered in the quadrates of the 2001 survey, some species previously present were more common, others much less common or not observed (Table 1). The FQI of the sedge meadow was _____ in 2001.

Wet-mesic sand prairie: In the wet-mesic sand prairie, located in the northwestern part of the preserve, surface water was present for short periods even during the growing season and the soil had a dark A horizon. This community was dominated by *Solidago canadensis* (IV of 34.2) followed by *Andropogon gerardii* (25.9) and *Carex* spp. (21.9) most of which was *C. stricta* (Table 2). Fourth in dominance, the exotic species *Poa pratensis*, was found throughout much of the wet-mesic prairie. Numerous species were encountered, 52 occurring in the quadrates with many other taxa being observed. Most were uncommon, however, 15 being encountered only once in the plots (Table 2). The FQI of the wet-mesic sand prairie was _____.

Shrub prairie: In some areas of the preserve the wet-mesic sand prairie gradually changes to shrub prairie. This shrub prairie exists since it rarely burns, mostly because of wet depressions that prevent the movement of fire.

Cornus drummondii, *Cornus obliqua*, *Rosa palustris*, *Rubus* spp., and *Salix discolor* were common woody species of the shrub prairie along with *Betula nigra*, which occasionally forms thickets. The ground layer species were similar to those associated with the wet-mesic sand prairie, although the rarer taxa were not encountered (Table 2). The FQI of the shrub prairie was _____.

DISCUSSION

The sedge meadow at the Matanzas Prairie Nature Preserve is similar to three sedge meadows studied in Lee County, Illinois (Handel et al. 2003). Species composition was similar with *Carex stricta* having very high importance values, and many of the subordinate species also being present. The Lee County sedge meadows, however, had higher species diversity, and *Onoclea sensibilis* ranked high in IV, while it was very uncommon at the Matanzas Prairie site. In contrast, *Calamagrostis canadensis* ranked high in IV on the Matanzas Prairie sedge meadow, being relative uncommon in the Lee County sedge meadows.

Similar results were obtained when comparing the prairie remnants studied in Lee County (Handel et al. 2003) with the wet-mesic sand prairie at Matanzas Prairie. Many of the species were common to both sites, as were some of the dominant species. Both sites had high species diversity and *Andropogon gerardii* and *Euthamia graminifolia* generally had high IV's. The Lee County prairie remnants, however, were drier and would be classified as mesic prairies.

The persistent presence of *Poa pratensis* in the ground layer of the wet-mesic sand prairie and the shrub prairie at the Matanzas Prairie Nature Preserve indicates that in the past this area was heavily grazed (Tables 3 and 4). More intensive management will be needed to reduce the importance of *Poa pratensis*. Management practices presently employed at the Matanzas Prairie Nature Preserve are only partially successful. A more aggressive fire regime is needed to decrease the extent of the exotic *Poa pratensis*, which would probably decrease in importance with the use of early spring burns.

In the past woody encroachment was extensive at the preserve and a 2-3 ha forest dominated by *Betula nigra* was present between the sedge meadow and the shrub prairie (Uhlarik et al. 1990). In this small forest, tree density averaged 579 stems/ha and basal area averaged 24.04 m²/ha. *Betula nigra*, which accounted for 90% of the importance value, had an average diameter of 22.6 cm dbd. Prescribed burns within the past 13 years have decreased the size of this small forest, and many of the trees are dead or dying. These prescribed burns should be continued, and probably increased in frequency, to stop woody encroachment in the shrub prairie. Also, the population of the state threatened *Tomanthera auriculata* on the preserve has decreased in the past 12 years. This rare species is associated with disturbances, and its seeds require light for germination (Midewin National Tallgrass Prairie 1999). In order to maintain this population, a management regime that includes disturbances such as fire and the removal of woody invaders must be undertaken.

ACKNOWLEDGMENTS

The authors would like to thank Dr. Gordon Tucker, Eastern Illinois University, for help with identification of the Cyperaceae, John Wilker, Natural Heritage Biologist, Illinois Department of Natural Resources, for his help and advice, and Daniel Busemeyer, Illinois Natural History Survey, for his field assistance and help in plant identification. The Illinois Department of Natural Resources, Wildlife Preservation Funds were used to support this project.

LITERATURE CITED

- Anderson, R.C. and L.E. Brown. 1983. Comparative effects of fire on trees in a midwestern savannah and an adjacent forest. *Bulletin of the Torrey Botanical Club* 110:87-90.
- Anderson, R.C. and L.E. Brown. 1986. Stability and instability in plant communities following fire. *American Journal of Botany* 73:364-368.
- Bailey, A.W., and C.E. Poulton. 1968. Plant communities and environmental relationships in a portion of the Tillamook burn, northwestern Oregon. *Ecology* 49:1-13.
- Calsyn, D.E. 1995. Soil survey of Mason County, Illinois. Soil Report 146, University of Illinois Agricultural Experiment Station, Urbana. ix+211 pp.
- Coates, D.T., S.E. Jenkins, and J.E. Ebinger. 1992. Woody vegetation survey of Barkhausen Woods, a closed canopy sand forest in Mason County, Illinois. *Erigenia* 12:1-6
- Daubenmire, R. 1959. A canopy coverage method of vegetation analysis. *Northwest Science* 33:43-64.
- Fernald, M.L. 1950. *Gray's Manual of Botany*, 8th edition, American Book Company, New York. lxiv+1632 pp.
- Gleason, H.A. 1910. The vegetation of the inland sand deposits of Illinois. *Bulletin of the Illinois State Laboratory of Natural History* 9:21-174.
- Gleason, H.A., and A.Cronquist. 1991. *Manual of the vascular flora of northeastern United States and adjacent Canada*. Second Edition. The New York Botanical Garden, Bronx, New York. lxxv+910 pp.
- Handel, W.C., L.R. Phillippe, and J.E. Ebinger. 2003. Floristic assessment of sand prairies and sedge meadows, Lee County, Illinois. *The Prairie Naturalist* 35:33-46.
- Herkert, J.R. and J.E. Ebinger, 2002. editors. *Endangered and threatened species of Illinois: Status and distribution*. Volume 1. Plants. Illinois Endangered Species Protection Board, Springfield, Illinois. 161 pp.
- Iverson, L.R., G.L. Rolfe, T.J. Jacob, A.S. Hodgins, and M.R. Jeffords. 1991. *Forests of Illinois*. Illinois Council on Forest development, Urbana, and Illinois Natural History Survey, Champaign, Illinois.
- Jenkins, S.E., J.E. Ebinger, and W.E. McClain. 1991. Woody vegetation survey of Bishop's Woods, a sand forest in Mason County, Illinois. *Transactions of the Illinois State Academy of Science* 84:20-27.
- King, J.E. 1981. Late Quaternary vegetational history of Illinois. *Ecological Monographs* 51:43-62.
- McClain, W.E., L.R. Phillippe, and J.E. Ebinger. 2004. Floristic assessment of the Henry Allan Gleason Nature Preserve, Mason County, Illinois. *Castanea* (submitted for publication)
- Midewin National Tallgrass Prairie. 1999. *Conservation Strategy: Earleaf floglove (Agalinus auriculata)*. Draft, Midewin National Tallgrass Prairie, Wilmington, Illinois.

Midwestern Regional Climate Center. 2002. <http://mcc.sws.uiuc.edu>

Mohlenbrock, R.H. 1986. Guide to the Vascular Flora of Illinois, revised and enlarged edition, Southern Illinois University Press, Carbondale, Illinois. viii+507 pp.

Rodgers, C.S. and R.C. Anderson. 1979. Presettlement vegetation of two prairie peninsula counties. *Botanical Gazette* 140:232-240.

Schwegman, J.E. 1973. Comprehensive plan for the Illinois nature preserves system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, Illinois. map+32 pp.

Taft, J.B., G.S. Wilhelm, D.M. Ladd, and L.A. Masters. 1997. Floristic quality assessment for vegetation in Illinois, a method for assessing vegetation integrity. *Erigenia* 15:1-95.

Transeau, E.N. 1935. The prairie peninsula. *Ecology* 16:423-437.

Uhlarik, C.A., J.E. Ebinger, and W.E. McClain. 1990. A study of four river birch stands in Mason and Cass counties, Illinois. *Transactions of the Illinois State Academy of Science* 83:149-155.

White J. 1978. Illinois Natural Areas inventory. Technical report. Volume I. Survey methods and results. Illinois Natural Areas Inventory, Urbana. xix+426 pp.

White, J., and M.H. Madany. 1978. Classification of natural communities in Illinois. Pp. 310-405 in Illinois Natural Areas Inventory. Technical report. (J. White, editor). Illinois Natural Areas Inventory, Urbana, Illinois.

Willman, H.B. 1973. Geology along the Illinois waterway - a basis for environmental planning. Illinois State Geological Survey Circular 478. Urbana. 48pp.

Willman, H.B. and J.C. Frye. 1970. Pleistocene stratigraphy of Illinois. Illinois State Geological Survey Bulletin 94:1-204.

Table 1. Average cover and importance value of the ground layer species encountered in the 1999 and 2001 study of the sedge meadow at the Matanzas Prairie Nature Preserve, Mason County, Illinois.

Species	1999		2001	
	Average Cover	I. V.	Average Cover	I.V.
<i>Carex stricta</i>	35.04	66.6	23.81	38.4
<i>Calamagrostis canadensis</i>	11.51	28.3	59.25	69.1
<i>Rosa palustris</i>	15.80	28.3	0.96	2.2
<i>Boehmeria cylindrica</i>	4.41	14.1	0.18	1.3
<i>Polygonum sagittatum</i>	2.20	14.0	0.79	4.2
<i>Thelypteris palustris</i>	2.78	10.5	10.17	20.3
<i>Aster umbellatus</i>	1.53	7.6	4.94	14.6
<i>Lycopus virginicus</i>	0.48	6.6	3.90	15.4
<i>Campanula aparinoides</i>	0.14	4.4	--	--
<i>Scutellaria galericulata</i>	0.79	3.8	0.91	3.6
<i>Mimulus ringens</i>	1.06	2.8	0.36	1.0
<i>Leersia oryzoides</i>	0.43	2.5	--	--
<i>Iris shrevei</i>	0.19	2.1	0.36	1.0
<i>Lycopus americanus</i>	0.18	1.6	--	--
<i>Lysimachia thrysiflora</i>	0.13	1.6	0.31	1.0
<i>Epilobium leptophyllum</i>	0.03	1.4	0.42	1.5
<i>Agrimonia parviflora</i>	0.02	0.9	--	--
<i>Eupatorium maculatum</i>	0.30	0.9	3.21	6.3
<i>Sagittaria latifolia</i>	0.30	0.9	--	--
<i>Fragaria virginiana</i>	0.06	0.6	--	--
<i>Galium trifidum</i>	0.01	0.5	0.11	2.2
<i>Triadenum fraseri</i>	--	--	0.48	5.0
<i>Bidens coronata</i>	--	--	0.63	3.8
<i>Eupatorium perfoliatum</i>	--	--	0.80	3.2
<i>Aster firmus</i>	--	--	0.61	1.6
<i>Onoclea sensibilis</i>	--	--	0.36	1.0
<i>Solidago canadensis</i>	--	--	0.12	0.8
<i>Liparis loeselii</i>	--	--	0.30	0.7
<i>Oxypolis rigidior</i>	--	--	0.06	0.5
<i>Rumex altissimus</i>	--	--	0.06	0.5
<i>Chelone glabra</i>	--	--	0.01	0.4
<i>Polygonum punctatum</i>	--	--	0.01	0.4
Totals	77.39	200.0	113.12	200.0
Average bare ground	13.11		3.15	

Table 2. Average cover and importance value of the ground layer species encountered in the wet-mesic sand prairie and the shrub prairie at the Matanzas Prairie Nature Preserve, Mason County, Illinois.

Species	SAND PRAIRIE		SHRUB PRAIRIE	
	Average Cover	I. V.	Average Cover	I. V.
<i>Solidago canadensis</i>	19.95	34.2	27.72	36.5
<i>Andropogon gerardii</i>	14.56	25.9	9.05	16.4
<i>Carex</i> (mostly <i>stricta</i>)	9.62	21.9	10.68	21.2
<i>Poa pratensis</i>	7.42	18.0	9.27	20.1
<i>Euthamia graminifolia</i>	7.20	17.7	9.21	17.3
<i>Fragaria virginiana</i>	4.52	11.0	6.02	11.5
<i>Rubus flagellaris</i>	4.86	10.9	8.73	14.9
<i>Vernonia missurica</i>	3.42	7.6	5.52	9.6
<i>Potentilla simplex</i>	2.25	5.3	1.20	2.1
<i>Sorghastrum nutans</i>	1.51	5.1	0.72	1.6
<i>Helianthus grosseserratus</i>	2.01	4.4	5.79	10.2
<i>Agrimonia parviflora</i>	1.85	4.1	3.15	4.9
<i>Rosa palustris</i>	2.13	4.1	3.41	5.2
<i>Cornus drummondii</i>	1.64	3.4	1.62	3.1
<i>Anemone canadensis</i>	1.20	3.1	1.80	4.0
<i>Ulmus americana</i>	1.05	2.8	1.62	3.1
<i>Dichanthelium acuminatum</i>	0.53	2.5	0.02	0.5
<i>Cassia fasciculata</i>	0.16	2.0	1.53	6.1
<i>Achillea millefolium</i>	0.21	1.8	0.01	0.2
<i>Pycnanthemum tenuifolium</i>	0.66	1.6	0.43	1.4
<i>Dichanthelium clandestinum</i>	0.37	1.1	0.60	1.0
<i>Pycnanthemum pilosum</i>	0.68	1.1	0.30	0.5
<i>Salix discolor</i>	0.75	1.1	--	--
<i>Cyperus strigosus</i>	0.36	1.0	0.06	0.3
<i>Muhlenbergia mexicana</i>	0.36	1.0	0.06	0.3
<i>Hypericum punctatum</i>	0.10	0.7	--	--
<i>Apocynum sibiricum</i>	0.07	0.5	0.37	1.0
<i>Lycopus americanus</i>	0.15	0.3	0.20	1.4
<i>Solidago gigantea</i>	0.16	0.5	0.66	1.3
<i>Equisetum arvense</i>	0.04	0.3	0.18	0.9
<i>Toxicodendron radicans</i>	--	--	0.36	0.8
<i>Lysimachia quadriflora</i>	0.03	0.1	0.12	0.6
Others (25 species/5 species)	1.59	4.9	0.96	2.0
Totals	91.41	200.0	111.37	200.0
Average bare ground	17.96		2.57	

APPENDIX 1

The vascular plant species encountered at Matanzas Prairie Nature Preserve are listed below by major groups, Pteridophytes (fern and fern-allies) and Spermatophytes (flowering plants), the latter divided into Monocots and Dicots. The families, genera, and species are alphabetically arranged within each group. An asterisk indicates species that have been introduced into Illinois (*). After the binomial and authority, the communities where the species were commonly encountered is given (1 = sand forest, 2 = wet sand prairie, 3 = shrub prairie, 4 = sedge meadow, 5 = cultural, 6 = open water of ditches). Collecting numbers are those of Morris (M), Feist (F), Ebinger (E), and Phillippe (P).

PTERIDOPHYTA

ASPLENIACEAE

Onoclea sensibilis L.;4; M429

EQUISETACEAE

Equisetum arvense L.;2,3; M180;

OPHIOGLOSSACEAE

Botrychium dissectum Spreng.;1,3; M606

Botrychium virginianum (L.) Sw.;1; M177

THELYPTERIDACEAE

Thelypteris palustris Schott;1,4,; M493

SPERMATOPHYTA

MONOCOTS

ALISMACEAE

Sagittaria latifolia Willd.;4;observed

COMMELINACEAE

Tradescantia ohiensis Raf.;2,3; M319

CYPERACEAE

Carex albolutescens Schw.;1; M289

Carex annectens Bickn.;1,2,5; M201

Carex blanda Dewey;1; M197

Carex brevior (Dewey) Mack.;1,2,5, M199

Carex bushii Mack.;1,2,3; M293

Carex buxbaumii Wahlenb.;1,4; M195

Carex conjuncta Boott;5; M200

Carex frankii Kunth;5; M377.1

Carex hirsutella Mack.;2,3; M414

Carex hyalinolepis Steud.;1; E28445

Carex hystericina Willd.;1; E28444

Carex jamesii Schwein.;1; M194

Carex lanuginosa Michx.;1,2,3; M198

Carex lurida Wahlenb.;4,5; M275

Carex molesta Mack.;2; M310

Carex normalis Mack.;2,3; E28448

Carex scoparia Willd.;1,2,5; M196

Carex stipata Muhl.;2,4,5; M200

Carex stricta Lam.;2,3,4; M195

Carex tenera Dewey;1; M199

Carex vulpinoidea Michx.;2,5; M270

Cyperus esculentus L.;5; M634

Cyperus filiculmis Vahl var. *macilentus* Fern.;5; M389

Cyperus schweintzii Torr.;5; M449

Cyperus strigosus L.;2,3,5; M523

Eleocharis acicularis (L.) Roem. & Schultes;5; M715

Eleocharis elliptica Kunth;4; E28630

- Eleocharis smallii* (L.) Britt.; 4; F205
Eleocharis verrucosa (Svens.) Harms; 2, 3, 5; M204
Scirpus atrovirens Willd.; 2, 5; M370
Scirpus cyperinus (L.) Kunth; 5; M460
Scirpus georgianus Harper; 5; E28847
Scirpus micranthus Vahl; 5; M630
Scirpus pendulus Muhl.; 2; M298
Scleria triglomerata Michx.; 2, 3; M357
 IRIDACEAE
Iris shrevei Small; 1, 4, 5; M222
 JUNCACEAE
Juncus acuminatus Michx.; 4; M426
Juncus canadensis J. Gay; 4; M492
Juncus dudleyi Wieg.; 2, 3; M279
Juncus interior Wieg.; 2, 3; M318
Juncus marginatus Rostk.; 2, 5; M450
Juncus tenuis Willd.; 2; M360
Juncus torreyi Coville; 2, 5; F212
 LEMNACEAE
Lemna minor L.; 6; M546
 LILIACEAE
 **Asparagus officinalis* L.; 5; M248
Polygonatum commutatum (Schult.) A. Dietr.; 1; M225
Smilacina racemosa (L.) Desf.; 1; M226
 ORCHIDACEAE
Liparis liliifolia (L.) Rich; 1; M372
Liparis loeselii (L.) Rich.; 2, 4; observed
Platanthera flava (L.) Lindl. var. *herbiola* (R. Br.) Luer; 5; M247
Spiranthes cernua (L.) Rich; 2, 4; M648
 POACEAE
Agrostis alba L.; 2, 3; M346
Agrostis hyemalis (Walt.) BSP.; 2, 3; M224
Alopecurus carolinianus Walt.; 5; M210
Andropogon gerardii Vitman; 2, 3; M554
Aristida purpurascens Poir.; 2, 5; M640
 **Bromus tectorum* L.; 5; M212
Calamagrostis canadensis (Michx.) Beauv.; 2, 3, 4; M280
Cenchrus longispinus (Hack.) Fern.; 5; M537
Dichanthelium acuminatum (Sw.) Gould & Clark; 2, 3; M385
Dichanthelium acuminatum (Sw.) Gould & Clark var. *lindheimeri* (Nash) Gould & Clark; 2, 5; M399
Dichanthelium clandestinum (L.) Gould; 2, 3; M368
Dichanthelium oligosanthos (Schult.) Gould; 2, 3, 5; M257
Dichanthelium villosissimum (Nash) Freckm.; 2, 3, 5; M302
 **Digitaria sanguinalis* (L.) Scop.; 5; M528
Echinochloa muricata (Beauv.) Fern.; 5; M614
Elymus canadensis L.; 2; E28851
Eragrostis spectabilis (Pursh) Steud.; 5; M572
Eragrostis trichodes (Nutt.) Wood; 5; M527
Festuca obtusa Biehler; 1; M283
Glyceria striata (Lam.) Hitchc.; 1; M286
Hordeum jubatum L.; 5; M261
Hordeum pusillum Nutt.; 5; M185
Leersia oryzoides (L.) Swartz; 1, 4; M622
Leersia virginica Willd.; 1; M657
Muhlenbergia mexicana (L.) Trin.; 2, 3; F216A
Muhlenbergia schreberi J. F. Gmel.; 5; M598

Panicum dichotomiflorum Michx.;5; M702
Panicum rigidulum Bosc;2,3; M530
Panicum virgatum L.;2; M416
Paspalum bushii Nash;5; M456
Paspalum ciliatifolium Michx. var. *stramineum* (Nash) Fern.;5; M456
Paspalum laeve Michx.;2; M555
**Phalaris arundinaceae* L.;4,5; M221
**Phleum pratense* L.;2; M353
**Poa compressa* L.;2,3; M306
**Poa pratensis* L.;2,3,5; M223
**Secale cereale* L.;5; M214
**Setaria faberi* Herrm.;5; M629
Sorghastrum nutans (L.) Nash;2,3; M553
Spartina pectinata Link;5; M538
Sphenopholis obtusata (Michx.) Scribn. var. *major* (Torr.) Erdman;1; M288
Sphenopholis obtusata (Michx.) Scribn. var. *obtusata*;2; M304
Tridens flavus (L.) Hitchc.;5; M575
Vulpia octoflora (Walt.) Rydb.;5; M186
POTAMOGETONACEAE
Potamogeton pusillus L.;6; M545
SMILACACEAE
Smilax hispida Muhl.;1; M596
TYPHACEAE
Typha latifolia L.;6; M582

DICOTS

ACANTHACEAE

Ruellia humilis Nutt.;2,3; M407

ACERACEAE

Acer negundo L.;1,3,5; M619

Acer saccharinum L.;1,5; M649

ANACARDIACEAE

Rhus aromatica Ait.;3; M164

Rhus glabra L.;2,3; M363

Toxicodendron radicans (L.) Kuntze;1,5; M441

APIACEAE

Cicuta maculata L.;1,5; M405

Eryngium yuccifolium Michx.;2; M410

Oxypolis rigidior (L.) Raf.;4; M490

Sanicula canadensis L.;1,2,3; M375

APOCYNACEAE

Apocynum cannabinum L.;2; M367

Apocynum sibiricum Jacq.;2,3; E28628

ASCLEPIADACEAE

Asclepias amplexicaulis Sm.;5; M259

Asclepias incarnata L.;4,5; M503

Asclepias hirtella (Pennell) Woodson;2; M472

Asclepias syriaca L.;2; M403

Cynanchum laeve (Michx.) Pers.;5; M459

ASTERACEAE

**Achillea millefolium* L.;2,3; M320

Ambrosia artemisiifolia L.;2,3; M557

Ambrosia trifida L.;1,5; M618

Antennaria neglecta Greene;2,3; M191

Aster dumosus L.;2; M214

Aster ericoides L.;2,3; M645A

Aster firmus Nees;4; M662

Lycopus rubellus Moench; 4; M504
Lycopus virginicus L.; 4; M518
Mentha arvensis L.; 1, 4; M604
Monarda punctata L.; 5; M522
 **Prunella vulgaris* L.; 1, 5; M448
Pycnanthemum tenuifolium Schrad.; 2, 3; M393
Pycnanthemum pilosum Nutt.; 2, 3, 5; M479
Pycnanthemum virginianum (L.) Dur. & Jacks.; 1; M438
Scutellaria galericulata L.; 2, 4; M364
Scutellaria lateriflora L.; 1, 2, 4; M364
Stachys palustris L. var. *homotricha* Fern.; 2, 3; M383
Teucrium canadense L. var. *virginicum* (L.) Eat.; 5; M455

LAURACEAE

Sassafras albidum (Nutt.) Nees; 1, 3; M165

LYTHRACEAE

Lythrum alatum Pursh; 2, 5; M366
Rotala ramosior (L.) Koehne; 5; M628

MENISPERMACEAE

Menispermum canadense L.; 1; M605

MOLLUGINACEAE

**Mollugo verticillatus* L.; 5; M526

MORACEAE

**Morus alba* L.; 1, 5; M172

NYCTAGINACEAE

**Mirabilis nyctaginea* (Michx.) MacM.; 2, 5; M461

OLEACEAE

Fraxinus pennsylvanica Marsh.; 1, 3; M665

ONAGRACEAE

Circaea lutetiana Aschers. & Magnus ssp. *canadensis* (L.); 1; M597
Gaura biennis L.; 2; M470
Epilobium coloratum Biehler; 2, 3; M516
Epilobium leptophyllum Raf.; 4; M488
Ludwigia alternifolia L.; 2, 3; M442
Ludwigia palustris (L.) Ell.; 4, 6; M512
Oenothera biennis L.; 2, 3; M552
Oenothera rhombipetala Nutt.; 5; M457
Oenothera laciniata Hill; 5; M215

OXALIDACEAE

Oxalis dillenii Jacq.; 2; M211
Oxalis stricta L.; 5; M380

PHYTOLACCACEAE

Phytolacca americana L.; 1; P26397

PLANTAGINACEAE

Plantago virginica L.; 5; M178

PLATANACEAE

Platanus occidentalis L.; 2, 5; M636

POLEMONIACEAE

Phlox glaberrima L.; 4; M430

POLYGALACEAE

Polygala cruciata L.; 2; M487A
Polygala sanguinea L.; 2; M423

POLYGONACEAE

Polygonum amphibium L.; 2, 4; M 513
Polygonum cristatum Engelm. & Gray; 1; M666
 **Polygonum hydropiper* L.; 5; M624
Polygonum hydropiperoides Michx.; 6; E28878
Polygonum lapathifolium L.; 6; M615

Polygonum pennsylvanicum L.; 2, 5; M451
 **Polygonum persicaria* L.; 5, 6; M508
Polygonum punctatum Ell.; 1, 2, 4, 5, 6; M507
Polygonum sagittatum L.; 4; M515
Polygonum scandens L.; 1; M602
Polygonum virginianum L.; 1; M496
 **Rumex acetosella* L.; 5; M187
Rumex altissimus Wood; 4; E28634
Rumex orbiculatus Gray; 5; M511
 **Rumex crispus* L.; 2, 5; M263
 PRIMULACEAE
Lysimachia lanceolata Walt.; 2; M365
Lysimachia quadriflora Sims; 2, 3, 4; F215
Lysimachia thysiflora L.; 4; F202
 RANUNCULACEAE
Anemone canadensis L.; 2, 3; M175
Anemone virginiana L.; 2; M347
Ranunculus abortivus L.; 1, 5; M189
Ranunculus longirostris Godr.; 6; M547
 RHAMNACEAE
 **Rhamnus cathartica* L.; 1, 3; M190
 **Rhamnus frangula* L.; 1, 2, 3; M651
 ROSACEAE
Agrimonia parviflora Ait.; 1, 2, 3, 4; M495
Fragaria virginiana Duchesne; 2, 3, 4; M174
Geum canadense Jacq.; 1; M381
 **Potentilla norvegica* L.; 5; M395
 **Potentilla recta* L.; 2, 5; E28642
Potentilla simplex Michx.; 2, 3; M188
Prunus serotina Ehrh.; 1, 5; M167
 **Rosa multiflora* Thunb.; 2, 5; M326
Rosa palustris Marsh.; 2, 3, 4; M299
Rosa setigera Michx.; 3, 5; M406
Rubus allegheniensis Porter; 2, 3, 5; E28436
Rubus flagellaris Willd.; 2, 3; M220
Rubus occidentalis L.; 5; M404
Rubus pennsylvanicus Poir.; 2; M327
 RUBIACEAE
Cephalanthus occidentalis L.; 3, 4; M421
Diodia teres Walt.; 5; M529
Galium aparine L.; 1; M192
Galium circaezans Michx.; 1; M290
Galium concinnum Torr. & Gray; 4; F201
Galium obtusum Bigel.; 2, 3; M313
Galium trifidum L.; 2, 4; E28436
Galium triflorum Michx.; 1; M373
 RUTACEAE
Zanthoxylum americanum Mill.; 1, 3; M166
 SALICACEAE
Populus deltoides Marsh.; 1, 2; M550
Salix discolor Muhl.; 1, 2, 3, 4, 5; M274
Salix exigua Nutt.; 1, 3, 5; M168
Salix nigra Marsh.; 1; M169
 SAXIFRAGACEAE
Penthorum sedoides L.; 6; M580
 SCROPHULARIACEAE
Agalinis aspera (Dougl.) Britt.; 2, 3; M483

Agalinus purpurea (L.) Pennell; 2, 3; F207
Chelone glabra L.; 4; M577
Gratiola neglecta Torr.; 1, 2, 4; M311
Lindernia dubia (L.) Pennell var. *anagallidea* (Michx.) Cooperrider; 5; M531
Mimulus ringens L.; 4; M501
Pedicularis lanceolata Michx.; 2, 4; M571
Penstemon pallidus Small; 2, 3; M173
Tomanthera auriculata (Michx.) Raf.; 2; M642
 **Verbascum thapsus* L.; 5; E28884
 **Veronica arvensis* L.; 5; M208
Veronica peregrina L.; 5; M209

SOLANACEAE

Physalis heterophylla Nees; 2, 3; M249
Solanum carolinense L.; 5; M401
Solanum ptycanthum Dunal; 1; M676

ULMACEAE

Ulmus americana L.; 2, 3; observed

URTICACEAE

Boehmeria cylindrica (L.) Sw.; 1, 4; M428
Laportea canadensis (L.) Wedd.; 1; M608
Parietaria pensylvanica Muhl.; 4; M277
Pilea pumila (L.) Gray; 1; M595

VERBENACEAE

Phyla lanceolata (Michx.) Greene; 4, 5; M422
Verbena hastata L.; 2, 3, 4, 5; M391
Verbena stricta Vent.; 5; M392

VIOLACEAE

Viola lanceolata L.; 2; M179
Viola pratensis Greene; 1; M219
 **Viola rafinesquii* Greene; 2; M182
Viola sagittata Ait.; 2; observed
Viola sororia Willd.; 2, 3; M218

VITACEAE

Ampelopsis cordata Michx.; 5; E28887
Parthenocissus quinquefolia (L.) Planch.; 1; M710
Vitis riparia Michx.; 1, 5; M254
Vitis aestivalis Michx.; 1, 5; M705

WHITE OAK CREEK WOODS NATURAL AREA, MASON COUNTY, ILLINOIS

William E. McClain, Loy R. Phillippe, Daniel T. Busemeyer, and John E. Ebinger, Illinois Natural History Survey, Champaign, Illinois 61820

ABSTRACT: *Quercus alba* (white oak), a dominant tree of mesic and wet-mesic forests in Illinois, is uncommon in the extensive sand deposits of the state. In the Kankakee sand deposits of northeastern Illinois white oak sometimes occurs on lower dune slopes and swales. This species, however, is rarely found in the Mississippi and Illinois River sand deposits. An exception is at White Oak Creek Woods Natural Area near the Illinois River in Mason County. Located on a flat upland terrace, white oak dominates the overstory where it accounts for nearly 70% of the importance value. *Quercus velutina* (black oak), *Prunus serotina* (black cherry), and *Sassafras albidum* (sassafras) account for most of the remaining stems. No oak saplings were found in the natural area, probably the result of a closed canopy due to past fire suppression.

INTRODUCTION

Quercus alba L. (white oak), an important forest tree throughout eastern United States, is common from Maine to Michigan and Minnesota, and south to eastern Texas and northern Florida (Gleason and Cronquist 1991). Throughout the northern half of its range, this species is a leading dominant of mesic and wet-mesic oak forests, woodlands, and savannas. In these communities white oak is commonly associated with *Quercus velutina* Lam. (black oak), *Quercus rubra* L. (red oak), and various species of *Carya* spp. (hickories) (Braun 1950, Ebinger 1997, Tyrrell et al. 1998).

White oak is well adapted to mesic habitats in Illinois, occurring in upland forests, open woodlands, prairie groves, and scattered trees associated with prairies. Known from all Natural Divisions of Illinois, it is commonly listed as a component of many different plant communities (Schwegman 1973, Mohlenbrock and Ladd 1978, White and Madany 1978). It is, however, not a common species in the sand deposits of the state.

Sand deposits account for nearly 5% of the land surface of Illinois and generally occur on glacial outwash plains associated with erosional events of Wisconsin glaciation in the northern half of the state (Schwegman 1973, Willman and Frye 1970, King 1981). Of the large sand deposits within Illinois, only the Kankakee Sand Area Section of the Grand Prairie Division consistently has communities in which white oak is a common component (Schwegman 1973).

In the Illinois River Section of the Mississippi River and Illinois Rivers Sand Areas Natural Division white oak is rarely present. The authors know of only one forest community of this natural division in which white oak is a common overstory component. The purpose of this study was to determine the composition and structure of this white oak

dominated forest and compares the results with other sand forests in Illinois.

DESCRIPTION OF THE STUDY SITE

White Oak Creek Woods Natural Area is located in Mason County, about 6 km south of Havana, Illinois (SE1/4 NW1/4 S23 T21N R9W). The site, about 2 ha in size, is located on a sandy upland terrace about 500 m west of the Illinois River immediately south of White Oak Creek. The soils are excessively drained Plainfield sand (Calsyn 1995), part of the dune and swale topography known as the Parkland Formation (Willman and Frye 1970).

Little is known about the history of this small tract of timber. This small tract, however, as well as the land north of White Oak Creek, was designated "grade B" dry sand forest by the Illinois Natural Areas Inventory (White 1978). Selective logging during the 1980 modified the area north of White Oak Creek and it is now designated "grade C" dry sand forest (Lerczak 2000). Registered as an Illinois Natural Heritage Landmark since 1983, it is presently designated the Speckman-Stelter Woods Land and Water Reserve by the Illinois Nature Preserves Commission (Lerczak 2000).

Based on weather data from Havana, mean annual precipitation is 96.0 cm, with May having the highest rainfall (11.3 cm). Mean annual temperature is 10.8°C with the hottest month being July (average of 24.6°C), and the coldest January (average of -5.0°C). Frost-free days range from 140 to 206, with the average being 173 day per year (Midwestern Regional Climate Center 2004).

METHODS

During late summer of 2004, a 75 m by 125 m section of the 2 ha natural area was surveyed by dividing the areas into 15 contiguous quadrates 25 m on a side. All living and dead-standing woody individuals ≥ 10.0 cm dbh were identified and their diameters recorded. From this data, living-stem density (stems/ha), basal area (m^2/ha), relative density, relative dominance, importance value (IV), and average diameter (cm) were calculated for each species. Determination of the IV follows the procedure used by McIntosh (1957), and is the sum of the relative density and relative dominance (basal area). Dead-standing density (stem/ha) and basal area (m^2/ha) was also determined.

Woody understory composition and density (stems/ha) was determined using nested circular plots 0.0001, 0.001, and 0.01 ha in size located at about 15 meter intervals along line transects within the study area (20 plots). Four additional 0.0001 ha circular plots were located 7 m from the center points of each of the 20 plot centers along cardinal compass directions (100 plots). In the 0.0001 ha plots, woody seedlings (≤ 50 cm tall) were counted; in the 0.001 ha circular plots small saplings (>50 cm tall and <2.5 cm dbh) were recorded; and in the 0.01 ha circular plots large saplings (2.5-9.9 cm dbh) were tallied.

RESULTS

Only 10 species were present in the overstory (Table 1). White oak dominated the larger diameter classes with an IV of 144.2 (200 possible), and an average diameter of 53.4 cm. Most of the larger white oaks had an open-grown appearance with large branches or branch scars

within 4 m of the ground. Black oak, also restricted to the higher diameter classes, was second in IV (12.7), and had an average diameter of 71.3 cm. The remaining trees were mostly in the 10-29 cm diameter classes. Dead-standing individuals averaged 11.7 stems/ha with an average basal area of 2.641 m²/ha, all being white and black oaks.

The woody understory was relatively sparse, being very open in many areas (Table 2). Throughout the woods *Sassafras albidum* (Nutt.) Nees (sassafras) dominated the seedling and sapling categories with 4300 seedlings/ha, 1600 small saplings/ha, and 545 large saplings/ha. White oak seedlings were also common, but no oak saplings were encountered.

DISCUSSION

Although the forests of the Speckman-Stelter Woods Land and Water Reserve and surrounding land were never clear-cut, they undoubtedly are now very different today compared to the early 1800s. According to Lerczak (2000), Ms. Stelter, the present owner of the property, recalled her great grandfather stating that it was possible to drive a wagon through the woods in the 1840s, an indication of the openness of the woods were at that time. Also mentioned was that many of the oaks were present as grubs which occurs when oaks are continually top-killed by frequent fires (Taft 1997).

The present appearance of White Oak Creek Woods compared to 150 years ago is due to a reduced fire frequency followed by a total absence of fire in recent decades (Taft 1997). In presettlement times frequent fires maintained much of this mesic to wet-mesic oak cover type, particularly along the western edge of its range (Ebinger and McClain 1991, McClain and Elzinga 1994). In general oaks are well adapted to fire due to their thick bark and ability to reproduce by sprouts, giving them a competitive advantage in areas of high fire frequencies. Oak densities in this presettlement landscape was dictated by fire frequency and intensity, ranging from low densities in savannas and woodlands that burned hot and frequently, to higher densities in closed forests where surface fires burned cooler and were less frequent (Anderson 1991, Abrams 1992).

The vegetation of White Oak Creek Woods Natural Area was surveyed in 1976 by the Illinois Natural Areas Inventory (INAI) (White 1978). At that time parts of the woods were designated "grade B" old-growth dry sand forest and tree density averaged 292 stems/ha with a basal area of 25.4 m²/ha. Black oak was the dominant overstory species with 100 stems/ha and a basal area of 17.0 m²/ha. *Ulmus americana* L. (American elm), white oak, and sassafras followed in importance. The area surveyed by the INAI, however, was to the north of White Oak Creek, not the small section to the south of White Oak Creek examined during the present survey.

No other forested areas examined in the Illinois River sand deposits contained white oak. Most were closed canopy dry sand forests on dune deposits where black oak and *Quercus marilandica* Muench. (blackjack oak) were the leading dominants along with a few hickory species in low numbers (Jenkins et al. 1991, Coates et al. 1992, McClain et al. 2002). One closed forest, Barton Woods, located on a terrace of Salt Creek in Mason County, is dominated by *Celtis occidentalis* L. (hackberry) and *Quercus macrocarpa* Michx. (bur oak) (McClain et al. 1993). In this wet-mesic forest white oak was not encountered though many mesic species

were present: *Gleditsia triacanthos* L. (honey locust), American elm, *Platanus occidentalis* L. (sycamore), *Ulmus rubra* Muhl. (slippery elm), *Juglans nigra* L. (black walnut), and *Quercus bicolor* Willd. (swamp white oak).

Rogers and Anderson (1979) used General Land Office survey records to determine the presettlement vegetation of Mason County in 1821-1824. In all community classes (prairie, savanna, open forest, and closed forest) black oak was the dominant woody species and usually accounted for more than half of the IV. Blackjack oak was second in IV in the open canopy communities (prairie and savanna) while in open and closed forests hickories and *Acer* spp. (maples) were second and third in importance. White oak was not common, but was occasionally recorded in savannas, and open and closed forests. These forests were mostly on the western edge of the county adjacent to the Illinois River and its backwater lakes. Decreased fire frequency in this region may have permitted the establishment of mesic to dry-mesic forest communities.

The only other sand area of Illinois where white oak is an occasional component of the vegetation is the Kankakee Sand Area Section of the Grand Prairie Natural Division. White oak is a component of dry-mesic sand savanna communities that exist on the lower slopes of dunes and swales (McDowell et al. 1983, Johnson and Ebinger 1992). On these sites the vegetation is an open savanna community with a herbaceous understory of native prairie species (Johnson and Ebinger 1995). In these communities periodic fire was commonly used to maintain an open overstory.

LITERATURE CITED

- Abrams, M. D. 1992. Fire and the development of oak forests. *BioScience* 42:346-353.
- Anderson R. C. 1991. Presettlement forest of Illinois. pages 9-19. in G. V. Burger, J. E. Ebinger, and G. S. Wilhelm (editors). Proceedings of the Oak Woods Management Workshop. Eastern Illinois University, Charleston, Illinois.
- Braun, E. L. 1950. Deciduous forests of Eastern North America. Blakiston, Philadelphia, Pennsylvania.
- Calsyn, D.E. 1995. Soil survey of Mason County, Illinois. Soil Report 146, University of Illinois Agricultural Experiment Station, Urbana. ix+211 pp.
- Coates, D. T., S. E. Jenkins, J. E. Ebinger, and W. E. McClain. 1992. Woody vegetation survey of Barkhausen Woods, a closed canopy sand forest in Mason County, Illinois. *Erigenia* 12:1-6.
- Ebinger, J. E. 1997. Forest Communities of the Midwestern United States. Pages 3-23. in M. W. Schwartz (editor). Conservation in highly fragmented landscapes. Chapman and Hall, New York.
- Ebinger, J. E. and W. E. McClain. 1991. Forest succession in the prairie peninsula of Illinois. *Illinois Natural History Survey Bulletin* 34:375-381.

Gleason, H. A. and A. Cronquist. 1991. Manual of vascular plants of northeastern United States and adjacent Canada. 2nd edition. The New York Botanical Garden, Bronx, New York.

Jenkins, S. E., J. E. Ebinger, and W. E. McClain. 1991. Woody vegetation survey of Bishop's Woods, a sand forest in Mason County, Illinois. Transactions of the Illinois State Academy of Science 84:20-27.

Johnson, K. C. and J. E. Ebinger. 1992. Effects of prescribed burns on the woody vegetation of a dry sand savanna, Hooper Branch Nature Preserve, Iroquois County, Illinois. Transactions of the Illinois States Academy of Science 85:105-111.

Johnson, K. C. and J. E. Ebinger. 1995. Effects of different fire regimes on the ground layer vegetation of a dry sand savanna, Hooper Branch Nature Preserve, Iroquois County, Illinois. *Erigenia* 14:37-40.

King, J. E. 1981. Late Quaternary vegetational history of Illinois. *Ecological Monographs* 51:43-62.

Lerczak, T. V. 2000. Proposal to register Speckman-Stelter Woods in Mason County as an Illinois Land and Water Reserve. Illinois Nature Preserves Commission, Springfield, Illinois. ii+14 pp.

McClain, W. E. and S. L. Elzinga. 1994. The occurrence of prairie and forest fires in Illinois and other Midwestern states, 1679 to 1854. *Erigenia* 13:79-90.

McClain, W. E., M. A. Jenkins, S. E. Jenkins, and J. E. Ebinger. 1993. Changes in the woody vegetation of a bur oak savanna remnant in central Illinois. *Natural Areas Journal* 13:108-114.

McClain, W. E., S. D. Turner, and J. E. Ebinger. 2002. Vegetation of forest communities at the Sand Prairie-Scrub Oak Nature Preserve, Mason County, Illinois. Transactions of the Illinois State Academy of Science. 95:37-46.

McDowell, B., J. Newman, and J. Ebinger. 1983. Survey of the woody vegetation of the Kankakee Sand Area Section of Indiana and Illinois. *Proceedings of the Indiana Academy of Science* 93:187-193.

McIntosh, R.P. 1957. The York Woods. A case history of forest succession in southern Wisconsin. *Ecology* 38:29-37.

Midwestern Regional Climate Center. 2004. <http://mcc.sws.uiuc.edu>

Mohlenbrock, R. H. and D. M. Ladd. 1978. Distribution of Illinois Vascular Plants. Southern Illinois University Press, Carbondale and Edwardsville, Illinois.

Rodgers, C. S. and R. C. Anderson. 1979. Presettlement vegetation of two prairie peninsula counties. *Botanical Gazette* 140:232-240.

Schwegman, J. E. 1973. Comprehensive plant for the Illinois nature preserves system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, Illinois.

Taft, J. B. 1997. Savanna and open woodland communities. Pages 24-54. in M. W. Schwartz (editor). Conservation in highly fragmented landscapes. Chapman and Hall, New York.

Tyrrell, L. E., G. J. Nowacki, T. R. Crow, D. S. Buckley, E. A. Nauertz, J. N. Niese, J. L. Rollinger, and J. C. Zasada. 1998. Information about old growth for selected forest type groups in the Eastern United States. USDA Forest Service. General Technical Report NC-197.

White, J. 1978. Illinois Natural Areas Inventory. Technical report. Volume I. Survey methods and results. Illinois Natural Areas Inventory, Urbana. xix+426 pp.

White, J., and M. N. Madany. 1978. Classification of natural communities in Illinois. pages 309-405. in J. White (editor). Illinois Natural Areas Inventory, Technical Report. Illinois Natural Areas Inventory, Urbana, Illinois.

Willman, H. B. and J. C. Frye. 1970. Pleistocene stratigraphy of Illinois. Illinois State Geological Survey Bulletin 94:1-204

Table 1. Density by diameter classes (stems/ha), basal area (m²/ha), relative density, relative dominance (basal area), importance value and average diameter (cm) of woody overstory species at White Oak Creek Natural Area, Mason County, Illinois.

Species	Diameter Class (cm)							Den. #/ha	Basal Area m ² /ha	Rel. Den.	Rel. Dom.	I. V.	Av. Diam. (cm)
	10-19	20-29	30-39	40-49	50-59	60-69	70+						
<i>Quercus alba</i>	--	--	11.7	29.9	35.2	24.5	4.3	105.6	24.577	58.6	85.6	144.2	53.4
<i>Quercus velutina</i>	--	--	--	--	1.1	1.1	4.3	6.5	2.611	3.6	9.1	12.7	71.3
<i>Prunus serotina</i>	18.1	2.1	--	--	--	--	--	20.2	0.284	11.2	1.0	12.2	12.9
<i>Sassafras albidum</i>	17.1	--	--	--	1.1	--	--	18.2	0.466	10.0	1.6	11.6	14.1
<i>Robinia pseudoacacia</i>	9.6	2.1	--	--	--	--	--	11.7	0.240	6.5	0.8	7.3	15.6
<i>Morus alba</i>	8.5	--	--	--	--	--	--	8.5	0.094	4.7	0.4	5.1	11.8
<i>Machura pomifera</i>	2.1	--	1.1	1.1	--	--	--	4.3	0.322	2.4	1.1	3.5	28.0
<i>Ulmus americana</i>	2.1	1.1	--	--	--	--	--	3.2	0.098	1.8	0.4	2.2	18.2
<i>Celtis occidentalis</i>	1.1	--	--	--	--	--	--	1.1	0.010	0.6	--	0.6	10.9
<i>Juglans nigra</i>	1.1	--	--	--	--	--	--	1.1	0.013	0.6	--	0.6	12.3
Totals	59.7	5.3	12.8	31.0	37.4	25.6	8.6	180.4	28.715	100.0	100.0	200.0	

Table 2. Density (individuals/ha) of woody understory species in a woodland community at White Oak Creek Natural Area, Mason County, Illinois.

Species	Seedlings	Small Saplings	Large Saplings
<i>Sassafras albidum</i>	4300	1600	545
<i>Celtis occidentalis</i>	1700	100	25
<i>Quercus alba</i>	1100	--	--
<i>Quercus velutina</i>	800	--	--
<i>Carya texana</i>	300	200	95
<i>Ulmus americana</i>	300	--	15
<i>Prunus serotina</i>	200	100	375
<i>Morus alba</i>	100	--	45
<i>Cercis canadensis</i>	100	--	20
<i>Asimina triloba</i>	100	--	--
<i>Carya tomentosa</i>	--	50	--
<i>Ulmus rubra</i>	--	--	145
<i>Robinia pseudoacacia</i>	--	--	75
<i>Crataegus mollis</i>	--	--	5
<i>Tilia americana</i>	--	--	5
<i>Fraxinus lanceolata</i>	--	--	5
<i>Juglans nigra</i>	--	--	5
<i>Cornus drummondii</i>	200	200	10
<i>Toxicodendron radicans</i>	7700	--	--
<i>Rubus allegheniensis</i>	1200	--	--
<i>Ribes missouriense</i>	900	--	--
Totals	19000	2250	1370

Herbaceous plant succession at Sand Prairie-Scrub Oak Nature Preserve,
Mason County, Illinois

William E. McClain, Adjunct Research Associate in Botany, Illinois
State Museum, Springfield, Illinois 62706
and

John E. Ebinger, Professional Scientist, Botanist, Illinois Natural
History Survey, Champaign, Illinois 61820

ABSTRACT: The herbaceous vegetation was studied at one-, three-, and twelve-year intervals following clearing of a pine plantation at the Sand Prairie-Scrub Oak Nature Preserve, Mason County, Illinois. The annual species *Setaria faberii* (giant foxtail), *Bidens bipinnata* (Spanish needles), and *Digitaria sanguinalis* (crabgrass) dominated the first year. The perennial *Dichanthelium villosissimum* (hairy panic grass) along with the non-native *Mollugo verticillata* (carpetweed) was prominent the third year while *Diodia teres* (buttonweed) and *Eragrostis trichodes* (sand love grass) dominated the twelfth year. Two dominants of mature sand prairie, *Schizachyrium scoparium* (little bluestem) and *Tephrosia virginiana* (goat's rue), were not present within the study area.

INTRODUCTION

Sand prairie communities are present in Illinois along the Green, Illinois, Kankakee, and Mississippi rivers, and Lake Michigan (Gleason 1910, Schwegman et al. 1973, Vestal 1913). Due to their arid nature they were the lasts of the extensive Illinois prairies to be converted to agriculture. In the early 1900s, thousands of acres of sand prairie still remained (Hart and Gleason 1907, Sampson 1921). With the advent of center pivot irrigation, much of the remaining sand prairie was converted to agriculture by 1960. The areas remaining were unsuitable for agriculture due to extensive sand dunes, large areas of actively moving sand, or depressions too difficult to drain.

The most extensive of the states' sand deposits is associated with the Illinois River in central Illinois. In these deposits the Illinois Department of Natural Resources, in cooperation with the Illinois Nature Preserves Commission, purchased and established several large nature preserves and state forests. One of the largest is Sand Prairie-Scrub Oak Nature Preserve (SP-SONP). At the time of purchase in 1969, this site was a mixture of remnant prairie, savanna, forest, and abandoned agricultural fields.

The opportunity to study the early stages of secondary plant succession at the SP-SONP occurred in the fall of 1992 when a large *Pinus banksiana* Lamb. (jack pine) plantation was cleared. This dense plantation was devoid of native woody and herbaceous vegetation, and pine removal exposed several hectares of bare mineral soil. The purpose of this study was to document plant succession changes over a period of twelve years.

DESCRIPTION OF THE STUDY AREA

SP-SONP is a 590 ha site located between the villages of Bath and Kilbourne (T20N, R9W, Sections 13, 14, 23, 26, of the 3rd PM), Mason County, Illinois. Part of the Illinois River Section of the Illinois and Mississippi River Sand Areas Natural Division (Schwegman et al.

1973), these sands were deposited by the Kankakee Torrent approximately 14,500 years ago (Willman and Frye 1970). Subsequent wind action created the dune and swale topography characteristic of the Parkland Formation (Willman and Frye 1970). The soils consist of about 95 % sand, 4 % silt, and 1 % clay. Soil pH ranges from 5.1 to 5.3 from prairie to open forest, and soils are low in organic matter and nitrogen (Benjamin et al. 1989).

The SP-SONP was a mixture of remnant prairie, sand forest, and savanna, tree plantations, and recently abandoned agricultural fields when purchased by the state. The only management activities have been prescribed burning for woody species control and invasive exotic species removal, mostly *Robinia pseudoacacia* L. (black locust). Abandoned agricultural fields have been allowed to re-vegetate naturally, and no plant propagules have been added.

The climate at SP-SONP is continental with warm summers and cold winters. Based on weather data from Havana, Illinois, about 12 km north of the site, mean annual precipitation is 96.0 cm, with May having the highest rainfall (11.3 cm). Mean annual temperature is 10.8° C, with July as the hottest month (average of 24.6° C), and the coldest being January (average of -5.0° C). Frost-free days range from 140 to 206, with the average being 173 days per year (Midwestern Regional Climate Center 2004).

METHODS

The study site was visited during September of 1993, 1995, and 2004. The ground layer vegetation was studied by aligning two randomly located transects, 25 m long, within the study area. Along each transect, 1/4 m² quadrates were located at 1 m intervals (n=25/transect), odd-numbered quadrates to the right even-numbered to the left. A random numbers table was used to determine the number of meters (0 to 9) a quadrate was located from the transect line. Cover was determined by using the Daubenmire cover class system (Daubenmire 1959) as modified by Bailey and Poulton (1968). Importance value (IV) for ground layer species was determined by summing relative cover and relative frequency. Only frequency of species was recorded the first year of the study.

Voucher specimens of each plant species were collected, identified, and deposited in the Stover-Ebinger Herbarium of Eastern Illinois University, Charleston, Illinois (EIU). Criteria for designating non-native species followed Mohlenbrock (2002), and Gleason and Cronquist (1991), while nomenclature follows Mohlenbrock (2002).

RESULTS

Eleven herbaceous species were present in sample plots during the first year of study, nine annuals and two perennials, four being non-native taxa (Table 1). The most frequently encountered species were *Bidens bipinnata* L. (Spanish needles), *Setaria faberii* F. Herrm. (giant foxtail), *Digitaria sanguinalis* (L.) Scop. (crabgrass), and *Diodia teres* Walt. (buttonweed) (Table 1).

The perennial *Dichanthelium villosissimum* (Nash.) Freckm. (hairy panic grass) was well established throughout the site by the third year. Annual species, such as Spanish needles and giant foxtail, had declined significantly, but the exotic annual *Mollugo verticillata* L.

(carpetweed) was second in importance. Seedlings of *Quercus velutina* Lam. (black oak), along with 23 other species were present in the plots. Of this total, 16 were perennial sand prairie species (Table 1).

Twenty-seven species were present in the plots during the twelfth year, including 16 native perennial sand prairie species (Table 1). Though buttonweed was the dominant species, the native perennial *Eragrostis trichodes* (Nutt.) Wood (sand love grass) was second followed by *Opuntia humifusa* (Raf.) Raf. (eastern prickly pear cactus), *Aristida desmantha* Trin. & Rupr. (three-awn grass), and hairy panic grass.

DISCUSSION

Abandoned agricultural fields located on sand deposits are colonized by native vegetation within a relatively short amount of time (McClain et al. 2005). During this twelve-year study, the herbaceous vegetation changed from non-native annuals to long-lived perennial sand prairie species (Table 1). One perennial, sand love grass, is an aggressive species that has dominated abandoned agricultural fields at SP-SONP for more than 60 years (McClain et al. 2005).

Though the vegetation of the study site currently consists of native sand prairie species, the composition is not similar to that of mature dry sand prairie. The most important grasses in mature dry sand prairie are *Schizachyrium scoparium* (Michx.) Nash. (little bluestem) and hairy panic grass, while *Tephrosia virginiana* (L.) Pers. (goat's rue), eastern prickly pear cactus, and *Ambrosia psilostachya* DC. (western ragweed) are usually among the top five in importance (Table 1).

Because no data are available on sand prairie restorations in Illinois, it is not known if planting seeds or plants of characteristic sand prairie species, such as little bluestem and goat's rue, will reduce the time required for their establishment. Prairie restoration guides contain little information on restoring sand prairies and most attempts to recreate prairie having taken place on loam soils (Packard and Mutel 1997, McClain 2003). Current management strategies allow for natural re-establishment of native flora within the old fields at SP-SONP.

The time required for prairie flora to invade abandoned fields at SP-SONP and become floristically similar in composition to native prairie is substantial. In a related study at SP-SONP, the flora of a 60-year old abandoned field still does not closely resemble the composition of mature remnant prairies (McClain et al. 2005). The reasons for this lengthy time requirement are not completely understood, but these results are consistent with other studies (Curtis 1959). The slow colonization of abandoned agricultural fields by dominant dry sand prairie flora may be due to heavy, gravity-dispersed seeds (Curtis 1959), disruption of the microrhizal community (Benjamin et al. 1989), or absence of a dispersal agent.

LITERATURE CITED

Bailey, A.W., and C.E. Poulton. 1968. Plant communities and environmental relationships in a portion of the Tillamook burn, northwestern Oregon. *Ecology* 49:1-13.

Benjamin, P., R.C. Anderson, and A.E. Liberta. 1989. Vesicular-arbuscular microrhizal ecology of little bluestem across a prairie-forest gradient. *Canadian Journal of Botany* 67:2678-2685.

- Curtis, J.T. 1959. The vegetation of Wisconsin. The University of Wisconsin Press. Madison. 657 pp.
- Daubenmire, R. 1959. A canopy coverage method of vegetation analysis. Northwest Science 33:43-64.
- Gleason, H.A. 1910. The vegetation of the inland sand deposits of Illinois. Bulletin of the Illinois State Laboratory of Natural History 9:21-174.
- Gleason, H.A., and A. Cronquist. 1991. Manual of the vascular flora of northeastern United States and adjacent Canada. Second Edition. The New York Botanical Garden, Bronx, New York. lxxv+910 pp.
- Hart, C.A., and H.A. Gleason. 1907. On the biology of the sand areas of Illinois. Bulletin of the Illinois State Laboratory of Natural History 7:137-272.
- McClain, W.E. 2003. Prairie Establishment and Landscaping, Second Edition. Illinois Department of Natural Resources. Springfield. 66 pp.
- McClain, W.E., J.E. Schwegman, T.A. Strole, L.R. Phillippe, and J.E. Ebinger. 2005. A floristic study of Sand Prairie-Scrub Oak Nature Preserve, Mason County, Illinois. (in review)
- Midwestern Regional Climate Center. 2004. <http://mcc.sws.uiuc.edu>
- Mohlenbrock, R.H. 2002. Vascular Flora of Illinois. Southern Illinois University Press. Carbondale. 490 pp.
- Packard, S. and C. Mutel (editors) 1997. The Tallgrass Prairie Restoration Handbook. Island Press, Washington D.C. 463 pp.
- Sampson, H.C. 1921. An ecological study of the prairie vegetation of Illinois. Bulletin of the Illinois State Laboratory of Natural History 13:523-577.
- Schwegman, J., G.B. Fell, M. Hutchison, W.M. Shepherd, and J. White. 1973. Comprehensive plan for the Illinois Nature Preserves System. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission. Rockford, Illinois. 32 p + map.
- Vestal, J.E. 1913. An association study of Illinois sand prairie. Bulletin of the Illinois State Laboratory of Natural History 10:1-96.
- Willman, H.B., and J.C. Frye. 1970. Pleistocene stratigraphy of Illinois. Illinois State Geological Survey Bulletin 94:1-204.

Table 1. Frequency (%), average cover, and importance value of the vascular plant species encountered in a mature sand prairie, a 12-year old successional field and a 3-year old successional field, and the frequency of species encountered in bare ground the year the area was bull-dozed to remove a pine plantation at Sand Prairie-Scrub Oak Nature Preserve, Mason County, Illinois

Species	MATURE PRAIRIE			12-YEAR OLD FIELD			3-YEAR OLD FIELD			BARE	
	Freq. %	Av. Cover	I.V.	Freq. %	Av. Cover	I.V.	Freq. %	Av. Cover	I.V.	Freq. %	
<i>Schizachyrium scoparium</i>	100	46.35	79.4	--	--	--	--	--	--	--	
<i>Dichanthelium villosissimum</i>	94	9.22	30.6	40	2.39	15.7	98	21.02	63.0	--	
<i>Tephrosia virginiana</i>	74	10.51	28.3	--	--	--	--	--	--	--	
<i>Ambrosia psilostachya</i>	68	6.74	22.2	8	0.48	3.1	2	0.30	1.0	--	
<i>Opuntia humifusa</i>	56	3.12	15.2	46	4.88	24.8	40	3.07	13.8	--	
<i>Eragrostis spectabilis</i>	24	0.91	6.0	6	0.13	1.6	--	--	--	--	
<i>Cyperus schweinitzii</i>	26	0.33	5.6	12	0.06	2.6	2	0.06	0.4	--	
<i>Calamagrostis longifolia</i>	18	0.09	3.7	--	--	--	--	--	--	--	
<i>Crotonopsis linearis</i>	10	0.05	2.1	4	0.02	0.9	98	2.33	22.3	--	
<i>Oenothera rhombipetala</i>	8	0.19	1.8	--	--	--	2	0.06	0.4	--	
<i>Cyperus grayoides</i>	8	0.09	1.7	2	0.01	0.4	38	0.54	7.9	58	
<i>Carex muhlenbergii</i>	4	0.12	1.0	8	0.19	2.2	18	1.21	5.8	--	
<i>Aristida tuberculosa</i>	4	0.07	0.9	--	--	--	--	--	--	--	
<i>Andropogon gerardii</i>	2	0.06	0.5	--	--	--	--	--	--	--	
<i>Bouteloua gracilis</i>	2	0.06	0.5	--	--	--	--	--	--	--	
<i>Pseudognaphalium obtusifolium</i>	2	0.06	0.5	--	--	--	--	--	--	--	
<i>Diodia teres</i>	--	--	--	84	7.97	42.1	14	0.90	4.5	38	
<i>Eragrostis trichodes</i>	--	--	--	54	6.21	30.5	6	0.90	3.1	--	
<i>Aristida desmantha</i>	--	--	--	78	1.72	21.2	--	--	--	--	
<i>Paspalum bushii</i>	--	--	--	24	2.35	12.3	2	0.01	0.3	--	
<i>Conyza canadensis</i>	--	--	--	26	1.25	9.2	2	0.06	0.4	--	
<i>Triplasis purpurea</i>	--	--	--	28	0.68	7.8	6	0.18	1.5	--	
<i>Rhus glabra</i>	--	--	--	10	1.47	6.6	16	2.40	8.0	--	
<i>Croton glandulosus</i>	--	--	--	14	0.07	3.0	92	2.20	21.0	12	
<i>Chamaecrista fasciculata</i>	--	--	--	4	0.60	2.7	--	--	--	--	
<i>Rubus allegheniensis</i>	--	--	--	4	0.60	2.7	6	0.42	2.1	--	
<i>Cyperus lupulinus</i>	--	--	--	10	0.05	2.2	--	--	--	--	

Floristic composition and structure of two dry sand prairies at Sand Ridge State Forest, Mason County, Illinois

Loy R. Phillippe, Paul B. Marcum, Daniel T. Busemeyer, Mary Ann Feist, William E. McClain, and John E. Ebinger, Illinois Natural History Survey, Champaign, Illinois 61820

ABSTRACT - Burns and Quiver prairies are dry sand prairies in small forest openings on ridges and swales of large stabilized dunes at Sand Ridge State Forest, Mason County, Illinois. Dominant species were nearly identical on both prairies. *Schizachyrium scoparium* (little bluestem) had an importance value of 40.1 on Quiver Prairie and 35.7 on Burns Prairie. *Tephrosia virginiana* (goat's-rue), *Opuntia humifusa* (common prickly pear), and *Ambrosia psilostachya* (western ragweed) were among the top five species on both prairies. Other common grasses were *Dichanthelium villosissimum* (hairy panic grass) on both prairies and *D. depauperatum* (panic grass) on Quiver Prairie.

INTRODUCTION

At the time of settlement by European man prairie vegetation covered about 60% of Illinois (Iverson et al. 1991). Most was tall-grass, black soil prairie common in the prairie peninsula of northeastern Illinois, though extensive tall grass prairies were also common in many parts of the state (Transeau 1935, Schwegman 1973). Depending upon soil and topography, other prairie types were common, including loess hill prairies, glacial till prairies, sand prairies, and gravel prairies (Schwegman 1973).

Sand prairies were relatively common in the northern half of Illinois, mostly on outwash plains that resulted from erosional events associated with Wisconsin glaciation (Willman and Frye 1970, King 1981). The most extensive of these sand deposits occurs in central Illinois in Mason and Cass counties (Gleason 1910, Schwegman 1973, Willman 1973). These deposits were formed about 14,500 years ago when glacial moraines and ice dams were breached, resulting in a flood known as the Kankakee Torrent. This flood removed extensive deposits of sand and gravel from glacial lakes in northeastern Illinois and adjacent Indiana. Most of this sand and gravel was deposited when the waters of the Kankakee Torrent slowed upon entering the broad lowlands of the Illinois River. These extensive sand deposits were then reworked by winds creating the present dune and swale topography.

Since early studies of Gleason (1910) very little has been published concerning the structure and composition of the ground layer vegetation of Illinois sand deposits. Though greatly modified by human activity, a few nature preserves and other high quality natural areas still occur in these sand deposits. The present study was undertaken to determine vascular plant species composition, vegetation structure, and floristic quality of two of the larger remaining dry sand prairie openings at Sand Ridge State Forest.

DESCRIPTION OF THE STUDY SITE

Sand Ridge State Forest is located in northwestern Mason County about 21 km northeast of Havana, and just west of Forest City, Illinois (parts of townships T22N R7W and T23N R7W). This 3,035 ha (11.7 sq. miles) state forest lies within the Illinois River Section of the Mississippi River and Illinois River Sand Area Natural Division (Schwegman 1973). Initial land purchases for the state forest began in 1939 for the purpose of stabilizing soil of abandoned farmlands,

developing a wood product industry, and setting land aside for public recreation. From the 1940s into the 1950's, pine plantations were established mostly on old pastureland and abandoned cultivated fields, but also in dry sand prairies that were scattered throughout the forest. Presently 1,012 ha of marketable pine plantations are present in the state forest while most of the remainder is oak-hickory dry sand forest and savanna (Andrews 2004).

A few small prairie openings containing various quality dry sand prairie remnants are scattered throughout the forest. These dry sand prairie communities are mostly less than 5 ha in size. The two prairies studied are:

Burns Sand Prairie (S1/2 NW1/4 SW1/4 S4 T22N R7W), about 4 ha in size, is located on a broad dune ridge and is surrounded by black oak dominated forest on three sides. To the east is an abandoned cultivated field with many prairie species. Small oaks, mostly less than 2 m tall, are scattered throughout the prairie, their concentration increasing near the forest edge.

Quiver Sand Prairie (E1/2 NW1/4 SE1/4 S28 T23N R7W), about 2.4 ha in size, is located in a wide, shallow depression on the sides of surrounding dunes that are covered with oak forest and closed savanna. A few small oaks, usually less than 1 m tall, occur throughout the prairie. Numerous small oaks are present at the forest/prairie boundary.

Sand Ridge State Forest has a continental climate with warm summers and cold winters. Based on weather data from Havana mean annual precipitation is 96.0 cm, with May having the highest rainfall (11.3 cm). Mean annual temperature is 10.8°C with the hottest month being July (average of 24.6°C), and the coldest January (average of -5.0°C). Frost-free days range from 140 to 206, with the average being 173 days per year (Midwestern Regional Climate Center 2004). The soils are mostly excessively drained Plainfield sands and Bloomfield sands (Calsyn 1995) that form the dune and swale topography known as the Parkland Formation (Willman and Frye 1970).

METHODS

Sand Ridge State Forest was visited more than 15 times in 2003 and 2004 to study the floristic composition of sand prairie and sand forest communities. Voucher specimens were collected, identified, and deposited in the herbarium of the Illinois Natural History Survey, Champaign, Illinois (ILLS). Determination of non-native species followed Mohlenbrock (2002) and Gleason and Cronquist (1991). Nomenclature follows Mohlenbrock (2002) while community classification follows White and Madany (1978). All species are listed in Appendix I along with the site, collector, and collecting number.

In late summer of 2004 four 25 m transects were located randomly along cardinal compass directions in Burns and Quiver prairies. Along each transect, 1 m² quadrates were spaced at 1 m intervals (n=25/transect), odd-numbered quadrates to the right, even-numbered quadrates to the left. A random numbers table was used to determine the number of meters (0 to 9) the quadrate was located from the transect line. Species cover was determined using the Daubenmire cover class system (Daubenmire 1959) as modified by Bailey and Foulton (1968). Importance value (IV) for ground layer species was determined by summing relative cover and relative frequency.

RESULTS

In the two sand prairies a total of 124 vascular plant species within 101 genera and 45 families were documented. Of these, none were fern and fern-allies, four were a gymnosperms, 83 dicots in 72 genera and 39 families, and 37 monocots in

27 genera and four families. One threatened species, *Cyperus grayioides* (sand prairie flatsedge) was encountered (Herkert and Ebinger 2002). The predominant plant families were the Poaceae with 24 species and the Asteraceae with 14 species (Appendix I).

Schizachyrium scoparium (little bluestem) had the highest importance value (IV) in both prairies. This species was present in nearly every plot on both prairies and had an IV of 35.7 on Burns Prairie and 40.1 on Quiver Prairie (Table 1 and 2). *Tephrosia virginiana* (goat's-rue), *Opuntia humifusa* (common prickly pear), and *Ambrosia psilostachya* (western ragweed) were among the top five species on both prairies. Other common grasses were *Dichanthelium villosissimum* (hairy panic grass) on both prairies and *D. depauperatum* (panic grass) on Quiver Prairie. Overall, the five native prairie species that are typical components of dry sand prairies had IV's greater than 10 (Table 1 and 2). All would be expected in high quality dry sand prairie communities in Illinois.

While two exotic species were encountered in the plots, only 11 were collected within the prairies (Appendix I). None were encountered in the plots on Quiver Prairie (Table 2), while *Mollugo verticillata* (carpetweed) were present in low numbers on Burns Prairie (Table 1). The remaining exotics were predominantly at the edges of the prairies, generally in areas of disturbance. The naturalized *Rumex acetosella* (sour dock), a pervasive species of many Illinois sand prairies, was not encountered in the plots or observed in either prairie.

DISCUSSION

Burns and Quiver dry sand prairies are very similar to the mature dry sand prairies at Henry Allen Gleason Nature Preserve (McClain et al. 2005), Long Branch Nature Preserve (Phillippe et al. 2005), and Sand Prairie-Scrub Oak Nature Preserve (McClain et al. 2005), all in Mason County, Illinois. Four of the top five dominants were identical in all preserves with little bluestem dominant and western ragweed, common prickly pear, and goat's-rue important subdominants. Of these species, goat's-rue was not found in the plots at Long Branch Nature Preserve. This species is relatively common on that prairie, but has a clumped distribution and was absent in the general area where the line transects were randomly placed (Phillippe et al. 2005).

Gleason (1910) referred to dry sand prairie communities as the bunch-grass association and listed many of species and dominants recorded during the present study. In all dry sand prairies little bluestem generally formed dense, various-sized clumps, usually circular in outline. These clumps were generally between 15 and 35 cm wide, some being dead in the center, an indication of their age. Open sand, common between the clumps, was occupied by other species. Bare ground and litter averaged 15.06 for Quiver Prairie and 27.31 for Burns Prairie (Table 1 and 2).

The lack of exotic species, high species diversity, and number of conservative prairie species, indicate that the mature dry sand prairie openings at Sand Ridge State Forest are of high natural quality. Many prairie openings were destroyed when pines were planted in them in the 1940s and early 1950s. Also, many small remnants have been lost due to woody encroachment and fire suppression. The few remaining sand prairie openings should be managed by periodic fires and by the removal of encroaching trees, particularly at the prairie/forest boundaries.

ACKNOWLEDGMENTS

The authors thank John Wilker, Natural Heritage Biologist, Illinois Department of Natural Resources, for his help and advice. The Illinois Department of Natural Resources, Wildlife Preservation Fund supported this project.

LITERATURE CITED

- Andrews, K. 2004. Forest Treasures. Outdoor Illinois. December 2004:2-5.
- Bailey, A.W. and C.E. Poulton. 1968. Plant communities and environmental relationships in a portion of the Tillamook burn, northwestern Oregon. *Ecology* 49:1-13.
- Calsyn, D.E. 1995. Soil survey of Mason County, Illinois. Soil Report 146, University of Illinois Agricultural Experiment Station, Urbana. ix+211 pp.
- Daubenmire, R. 1959. A canopy coverage method of vegetation analysis. *Northwest Science* 33:43-64.
- Gleason, H.A. 1910. The vegetation of the inland sand deposits of Illinois. *Bulletin of the Illinois State Laboratory of Natural History* 9:21-174.
- Gleason, H.A. and A. Cronquist. 1991. Manual of the vascular flora of northeastern United States and adjacent Canada. Second Edition. The New York Botanical Garden, Bronx, New York. lxxv+910 pp.
- Herkert, J.R. and J.E. Ebinger. 2002. editors. Endangered and Threatened Species of Illinois: Status and Distribution. Volume 1: Plants. Endangered Species Protection Board, Springfield, Illinois. 161 pp.
- Iverson, L.R., G.L. Rolfe, T.J. Jacob, A.S. Hodgins, and M.R. Jeffords. 1991. Forests of Illinois. Illinois Council on Forest Development, Urbana, and Illinois Natural History Survey, Champaign, Illinois.
- King, J.E. 1981. Late Quaternary vegetational history of Illinois. *Ecological Monographs* 51:43-62.
- McClain, W.E., L.R. Phillippe, and J.E. Ebinger. 2005. Floristic assessment of the Henry Allan Gleason Nature Preserve, Mason County, Illinois. *Castanea* (in press)
- McClain, W.E., J.E. Schwegman, T.A. Strole, L.R. Phillippe, and J.E. Ebinger. 2005. A floristic study of Sand Prairie-Scrub Oak Nature Preserve. (submitted).
- Midwestern Regional Climate Center. 2002. <http://mcc.sws.uiuc.edu>
- Mohlenbrock, R.H. 2002. Vascular Flora of Illinois. Southern Illinois University Press, Carbondale, Illinois. xi+490 pp.
- Phillippe, L.R., M.A. Feist, J.E. Ebinger, and W.E. McClain, 2005. Vascular Flora of Long Branch Nature Preserve, Mason County, Illinois. *Transactions of the Illinois State Academy of Science* (in press).
- Schwegman, J.E. 1973. Comprehensive plan for the Illinois nature preserves system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, Illinois. map+32 pp.
- Transeau, E.N. 1935. The prairie peninsula. *Ecology* 16:423-437.

White, J. and M.H. Madany. 1978. Classification of natural communities in Illinois. Pp. 310-405 in Illinois natural areas inventory. Technical report. (J. White, Editor). Illinois Natural Areas Inventory, Urbana, Illinois.

Willman, H.B. 1973. Geology along the Illinois waterway - a basis for environmental planning. Illinois State Geological Survey Circular 478. Urbana. 48 pp.

Willman, H.B. and J.C. Frye. 1970. Pleistocene stratigraphy of Illinois. Illinois State Geological Survey Bulletin 94:1-204.

Table 1. Frequency (%), average cover, relative frequency, relative cover, and importance value of the ground layer species encountered at Quiver sand prairie, Sand Ridge State Forest, Mason County, Illinois. (* non-native species)

Species	Freq. %	Average Cover	Rel. Freq.	Rel. Cover	I. V.
<i>Schizachyrium scoparium</i>	100	25.90	10.6	29.5	40.1
<i>Ambrosia psilostachya</i>	98	19.62	10.4	22.3	32.7
<i>Opuntia humifusa</i>	100	9.46	10.6	10.8	21.4
<i>Tephrosia virginiana</i>	43	11.80	4.6	13.4	18.0
<i>Dichanthelium depauperatum</i>	63	6.18	6.7	7.0	13.7
<i>Dichanthelium villosissimum</i>	56	2.12	5.9	2.4	8.3
<i>Cyperus lupulinus</i>	68	0.42	7.2	0.5	7.7
<i>Sporobolus clandestinus</i>	36	1.50	3.8	1.7	5.5
<i>Monarda punctata</i>	36	1.40	3.8	1.6	5.4
<i>Oenothera clelandii</i>	41	0.63	4.3	0.7	5.0
<i>Rhus aromatica</i>	7	2.56	0.7	2.9	3.6
<i>Carex muhlenbergii</i>	22	0.83	2.3	1.0	3.3
<i>Crotonopsis linearis</i>	29	0.15	3.1	0.2	3.3
<i>Conyza canadensis</i>	26	0.21	2.8	0.2	3.0
<i>Dichanthelium oligosanthes</i>	22	0.46	2.3	0.5	2.8
<i>Physalis heterophylla</i>	15	0.74	1.6	0.9	2.5
<i>Fallopia cristata</i>	19	0.25	2.0	0.3	2.3
<i>Solidago nemoralis</i>	20	0.15	2.1	0.2	2.3
<i>Pseudognaphalium obtusifolium</i>	14	0.63	1.5	0.7	2.2
<i>Bouteloua hirsuta</i>	13	0.44	1.4	0.5	1.9
<i>Chamaecrista fasciculata</i>	11	0.35	1.2	0.4	1.6
<i>Brickellia eupatorioides</i>	9	0.44	1.0	0.5	1.5
<i>Eragrostis spectabilis</i>	10	0.37	1.1	0.4	1.5
<i>Lespedeza capitata</i>	11	0.13	1.2	0.2	1.4
<i>Polygonum tenue</i>	10	0.05	1.1	0.1	1.2
<i>Carex tonsa</i>	6	0.28	0.6	0.3	0.9
<i>Euphorbia corollata</i>	8	0.09	0.8	0.1	0.9
<i>Asclepias verticillata</i>	7	0.04	0.7	--	0.7
<i>Commelina erecta</i>	5	0.05	0.5	0.1	0.6
<i>Cyperus schweinitzii</i>	6	0.03	0.6	--	0.6
<i>Leptoloma cognatum</i>	5	0.08	0.5	0.1	0.6
<i>Fragaria virginiana</i>	3	0.21	0.3	0.2	0.5
<i>Bouteloua curtipendula</i>	3	0.07	0.3	0.1	0.4
<i>Chrysopsis camporum</i>	2	0.06	0.2	0.1	0.3
<i>Eragrostis trichodes</i>	3	0.19	0.3	--	0.3
<i>Strophostyles helvula</i>	2	0.06	0.2	0.1	0.3
<i>Carex meadii</i>	2	0.01	0.2	--	0.2
<i>Chenopodium standleyanum</i>	2	0.01	0.2	--	0.2
<i>Froelichia floridana</i>	2	0.01	0.2	--	0.2
<i>Penstemon pallidus</i>	2	0.01	0.2	--	0.2
<i>Sorghastrum nutans</i>	2	0.04	0.2	--	0.2
<i>Croton glandulosus</i>	1	0.01	0.1	--	0.1
<i>Lactuca canadensis</i>	1	0.03	0.1	--	0.1
<i>Rhus glabra</i>	1	0.03	0.1	--	0.1
<i>Talinum rugospermum</i>	1	0.01	0.1	--	0.1
<i>Tridens flavus</i>	1	0.01	0.1	--	0.1
<i>Triplasis purpurea</i>	1	0.01	0.1	--	0.1
<i>Viola pedata</i>	1	0.01	0.1	--	0.1

Totals	88.14	100.0	100.0	200.0
Average bare ground and litter	15.06			

Table 2. Frequency (%), average cover, relative frequency, relative cover, and importance value of the ground layer species encountered at Burns sand prairie, Sand Ridge State Forest, Mason County, Illinois. (* non-native species)

Species	Freq. %	Average Cover	Rel. Freq.	Rel. Cover	I. V.
<i>Schizachyrium scoparium</i>	99	18.26	9.2	26.5	35.7
<i>Tephrosia virginiana</i>	60	13.49	5.6	19.6	25.2
<i>Dichanthelium villosissimum</i>	96	9.44	8.9	13.7	22.6
<i>Ambrosia psilostachya</i>	96	8.21	8.9	11.9	20.8
<i>Opuntia humifusa</i>	84	6.51	7.8	9.4	17.2
<i>Crotonopsis linearis</i>	94	0.50	8.7	0.7	9.4
<i>Pseudognaphalium obtusifolium</i>	45	2.56	4.2	3.7	7.9
<i>Cyperus lupulinus</i>	73	0.37	6.8	0.5	7.3
<i>Leptoloma cognatum</i>	33	2.76	3.0	4.0	7.0
<i>Carex muhlenbergii</i>	64	0.55	6.0	0.8	6.8
<i>Commelina erecta</i>	58	0.34	5.4	0.5	5.9
<i>Carex tonsa</i>	43	0.74	4.0	1.1	5.1
<i>Conyza canadensis</i>	50	0.33	4.6	0.5	5.1
<i>Croton glandulosus</i>	34	0.20	3.2	0.3	3.5
<i>Rhus aromatica</i>	10	1.77	0.9	2.6	3.5
<i>Oenothera clelandii</i>	21	0.13	2.0	0.2	2.2
<i>Eragrostis spectabilis</i>	14	0.49	1.3	0.7	2.0
<i>Lespedeza capitata</i>	10	0.54	0.9	0.8	1.7
<i>Dichanthelium depauperatum</i>	7	0.52	0.6	0.8	1.4
<i>Aristida tuberculosa</i>	12	0.11	1.1	0.2	1.3
<i>Cyperus schweinitzii</i>	10	0.05	0.9	0.1	1.0
<i>Eragrostis trichodes</i>	7	0.23	0.6	0.3	0.9
<i>Dichanthelium oligosanthes</i>	5	0.10	0.5	0.2	0.7
<i>Froelichia floridana</i>	9	0.07	0.8	0.1	0.9
<i>Liatris aspera</i>	3	0.19	0.3	0.3	0.6
<i>Quercus velutina</i>	3	0.21	0.3	0.3	0.6
<i>Polygonum tenue</i>	5	0.03	0.5	--	0.5
<i>Apocynum sibiricum</i>	3	0.07	0.3	0.1	0.4
<i>Bulbostylis capillaris</i>	4	0.02	0.4	--	0.4
<i>Talinum rugospermum</i>	4	0.02	0.4	--	0.4
<i>Fallopia cristata</i>	3	0.02	0.3	--	0.3
<i>Lactuca canadensis</i>	3	0.04	0.3	--	0.3
* <i>Mollugo verticillata</i>	3	0.02	0.3	--	0.3
<i>Paspalum bushii</i>	2	0.06	0.2	0.1	0.3
<i>Plantago patagonica</i>	3	0.02	0.3	--	0.3
<i>Cyperus grayoides</i>	2	0.01	0.2	--	0.2
<i>Diodia teres</i>	1	--	0.1	--	0.1
<i>Physalis virginiana</i>	1	0.03	0.1	--	0.1
<i>Solidago nemoralis</i>	1	0.03	0.1	--	0.1
<i>Chenopodium standleyanum</i>	1	0.05	--	--	--
<i>Euphorbia corollata</i>	1	0.05	--	--	--
Totals		69.14	100.0	100.0	200.0
Average bare ground and litter		27.31			

APPENDIX I. Vascular species encountered at two dry sand prairies at Sand Ridge State Forest, Mason County, Illinois, listed alphabetically by family under the major plant groups. An asterisk indicates non-native (exotic) species (*). Following the scientific name and in parenthesis is the name of the prairie where the taxon was found, (BP) for Burns dry sand prairie and (QP) for Quiver dry sand prairie. Following the name of the prairie, collecting numbers preceded by the initial of the collector's name are given (B) Daniel T. Busemeyer, (F) Mary Ann Feist, (G) Sophia Gehlhausen, (M) Paul B. Marcum, and (P) Loy R. Phillippe.

Sand Ridge State Forest Burns Prairie Species List

SPERMATOPHYTES: GYMNOSPERMS

CUPRESSACEAE

Juniperus virginiana L.: (BP) P 36479

PINACEAE

**Pinus resinosa* Ait.: (BP) P 37183

**Pinus strobus* L.: (BP) P 37175

**Pinus sylvestris* L.: (BP) P 36481

SPERMATOPHYTES: ANGIOSPERMS

MONOCOTS

COMMELINACEAE

Commelina erecta L. var. *erecta*: (BP) P 36950, (QP) F 2781, M 2846

Tradescantia ohiensis Raf.: (BP) P 36780, (QP) P 36752

CYPERACEAE

Bulbostylis capillaris (L.) C.B. Clarke: (BP) P 36952

Carex meadii Dewey: (BP) P 36782, (QP) P 36737

Carex muhlenbergii Schk.: (BP) P 36757, (QP) P 36736

Carex ionsa (Fern.) Bickn.: (BP) P 36478, (QP) F 2522

Cyperus grayioides Mohlenbr.: (BP) M 2684, P 36748

Cyperus lupulinus (Spreng.) Marcks var. *macilentus* (Fern.) Marcks: (BP) P 36949, (QP) F 2784

Cyperus schweinitzii Torr.: (BP) P 36945, (QP) F 2794

Eleocharis erythropoda Steud.: (BP) P 36955

Eleocharis ovata (Roth) Roem. & Schultes: (BP) P 36953

JUNCACEAE

Juncus acuminatus Michx.: (BP) P 36951

Juncus interior Wieg.: (BP) P 36763

POACEAE

Agrostis hyemalis (Walt.) BSP.: (BP) P 36759, (QP) F 2785

Andropogon gerardii Vitman: (BP) P 37173

Aristida tuberculosa Nutt.: (BP) P 37169, (QP) M 2848, M 2850

Bouteloua curtipendula (Michx.) Torr.: (QP) M 2826

Bouteloua hirsuta Lag.: (QP) M 2660

**Bromus tectorum* L.: (BP) P 36775

Calamovilfa longifolia (Hook.) Scribn.: (BP) M 2685

Dichanthelium depauperatum (Muhl.) Gould: (BP) P 36760, (QP) M 2844, P 36735

Dichanthelium oligosanthes (Schult.) Gould: (BP) P 36772, (QP) P 36738

Dichanthelium villosissimum (Nash) Freckm.: (BP) P 36756, (QP) P 36739

**Echinochloa crus-galli* (L.) P. Beauv.: (BP) P 36954

Eragrostis spectabilis (Pursh) Steud.: (BP) P 37170, (QP) M 2839

Eragrostis trichodes (Nutt.) Wood: (BP) Site Record, (QP) M 2845

Heterostipa spartea (Trin.) Barkworth: (BP) P 36781, (QP) F 2805

Hordeum pusillum Nutt.: (BP) P 36774

Leptoloma cognatum (Schult.) Chase: (BP) M 2683, (QP) M 2836

Paspalum bushii Nash: (BP) M 2682

**Poa pratensis* L.: (QP) F 2799

Schizachyrium scoparium (Michx.) Nash: (BP) Site Record, (QP) M 2829

Sorghastrum nutans (L.) Nash: (BP) P 37180, (QP) M 2834
Sporobolus clandestinus (Biehler) Hitchc.: (QP) M 2838
Tridens flavus (L.) Hitchc.: (QP) Site Record
Triplasis purpurea (Walt.) Chapm.: (QP) M 2847, M 2849
Vulpia octoflora (Walt.) Rydb. var. *tenella* (Willd.) Fern.: (BP) P 36773, (QP) P 36751

DICOTS

ACANTHACEAE

Ruellia humilis Nutt.: (QP) F 2791

AMARANTHACEAE

Froelichia floridana (Nutt.) Moq.: (BP) M 2679, (QP) Site Record

ANACARDIACEAE

Rhus aromatica Ait.: (BP) B 1675, P 36767, (QP) F 2801
Rhus glabra L.: (QP) F 2803
Rhus hirta L.: (QP) F 2802

APOCYNACEAE

Apocynum sibiricum Jacq.: (BP) Site Record

ASCLEPIADACEAE

Asclepias amplexicaulis Small: (BP) P 36766, (QP) F 2796
Asclepias hirtella (Pennell) Woodson: (BP) P 36956
Asclepias tuberosa L.: (QP) F 2790
Asclepias verticillata L.: (QP) M 2664

ASTERACEAE

**Achillea millefolium* L.: (QP) F 2783
Ambrosia psilostachya DC.: (BP) P 37172, (QP) M 2840
Brickellia eupatorioides (L.) Shinn.: (QP) M 2835
Chrysopsis camporum Greene: (BP) P 36795, P 37254, (QP) F 2780, M 2828
Conyza canadensis (L.) Cronq.: (BP) P 37178, (QP) M 2832
Erigeron strigosus Muhl.: (BP) P 36784, (QP) F 2778
Helianthus occidentalis Riddell: (QP) M 2852
Krigia virginica (L.) Willd.: (BP) B 1673, (QP) P 36741
Lactuca canadensis L.: (BP) P 37184, (QP) M 2842
Liatris aspera Michx. (BP) Site Record
Pseudognaphalium obtusifolium (L.) Hilliard & Burt.: (BP) P 37176, (QP) M 2837
Rudbeckia hirta L.: (QP) F 2787
Senecio plattensis Nutt.: (BP) P 36776, (QP) P 36749
Solidago nemoralis Ait.: (BP) P 37174, (QP) M 2833

BORAGINACEAE

Lithospermum croceum Fern.; (BP) P 36768, (QP) P 36740

BRASSICACEAE

**Alliaria petiolata* (Bieb.) Cavara & Grande: (B) B 1676
Arabis canadensis L.: (QP) P 36747
Draba reptans (Lam.) Fern.: (QP) F 2524
**Lepidium densiflorum* Schrad.: (BP) P 36770, (QP) P 36746
Lepidium virginicum L.: (BP) P 36769, (QP) F 2792

CACTACEAE

Opuntia humifusa (Raf.) Raf.: (BP) P 36755, (QP) F 2788

CAESALPINIACEAE

Chamaecrista fasciculata (Michx.) Greene: (BP) P 37253, (QP) M 2663

CAMPANULACEAE

Triodanis perfoliata (L.) Nieuwl.: (BP) P 36758, P 36946, (QP) F 2793

CHENOPODIACEAE

Chenopodium standleyanum Aellen: (QP) M 2665
Cycloloma atriplicifolium (Spreng.) Coult.: (BP) P 36958

CISTACEAE

Helianthemum canadense (L.) Michx.: (BP) P 36785, (QP) P 36750

CORNACEAE

Cornus drummondii C.A. Mey.: (BP) P 36790

EUPHORBIACEAE

Croton glandulosus L.: (BP) M 2680, (QP) F 2800
Crotonopsis linearis Michx.: (BP) M 2677, (QP) M 2662, M 2831
Euphorbia corollata L.: (BP) Site Record, (QP) F 2786

FABACEAE

Amorpha canescens Pursh: (QP) F 2789
Desmodium sessilifolium (Torr.) Torr. & Gray: (QP) M 2851
Lespedeza capitata Michx.: (BP) P 37179, (QP) M 2830
Strophostyles leiosperma (Torr. & Gray) Piper: (QP) M 2843
Tephrosia virginiana (L.) Pers.: (BP) P 36947, (QP) M 2841

FAGACEAE

Quercus marilandica Muench.: (QP) M 2667
Quercus velutina Lam.: (BP) P 37171

FUMARIACEAE

Corydalis micrantha (Englem.) Gray var. *micrantha*: (BP) P 36793

GERANIACEAE

Geranium carolinianum L.: (BP) P 36792

GROSSULARIACEAE

Ribes missouriense Nutt.: (BP) P 36482

LAMIACEAE

Monarda punctata L.: (QP) 2797
Teucrium canadense L.: (BP) P 37177, (QP) F 2798

MALVACEAE

Callirhoe triangulata (Leavenw.) Gray: (BP) M 2686, (QP) M 2661

MOLLUGINACEAE

**Mollugo verticillata* L.: (BP) P 36765

MORACEAE

**Morus tatarica* L.: (BP) P 36789

ONAGRACEAE

Oenothera clelandii W. Dietr., Raven, & W.L. Wagner: (BP) P 36957, (QP) F 2779

OXALIDACEAE

Oxalis violacea L.: (QP) P 36754

PLANTAGINACEAE

Plantago patagonica Jacq.: (BP) P 36761

POLEMONIACEAE

Phlox bifida Beck: (BP) P 36484, (QP) F 2525

POLYGONACEAE

Fallopia cristata (Engelm. & Gray) Holub.: (BP) P 37252, (QP) M 2666
Polygonum tenue Michx.: (BP) P 37251, (QP) M 2827

PORTULACACEAE

Talinum rugospermum Holz.: (BP) P 36764, (QP) G 36

PRIMULACEAE

Androsace occidentalis Pursh: (BP) P 36483, (QP) F 2523

ROSACEAE

Fragaria virginiana Duchesne: (BP) P 36777, (QP) P 36743
Geum canadense Jacq.: (BP) P 36778
Prunus serotina Ehrh.: (BP) P 36783, (QP) F 2804
Rosa carolina L. var. *carolina*: (BP) P 36786
Rubus allegheniensis Porter: (BP) P 36788
Rubus occidentalis L.: (BP) P 36779

RUBIACEAE

Diodia teres Walt. (BP) M 2681
Galium aparine L.: (BP) P 36771, (QP) P 36733

RUTACEAE

Ptelea trifoliata L.: (QP) P 36744

SCROPHULARIACEAE

Lindernia anagallidea (Michx.) Pennell: (BP) M 2678
Nuttallanthus canadensis (L.) D. Sutton: (BP) B 1674, (QP) P 36734
Penstemon pallidus Small: (QP) P 36748

SOLANACEAE

Physalis heterophylla Nees: (BP) P 37181, (QP) F 2795, M 2668

Physalis virginiana Mill.: (BP) P 36762

Solanum carolinense L.: (BP) P 36791

Solanum ptychanthum Dunal: (BP) P 36787

URTICACEAE

Parietaria pensylvanica Muhl.: (QP) P 36745


VIOLACEAE

Viola pedata L.: (QP) P 36753

**Viola rafinesquei* Greene: (BP) P 36480, (QP) P 36742

ZYGOPHYLLACEAE

**Tribulus terrestris* L.: (BP) P 36794



A Floristic Study of Sand Prairie-Scrub Oak Nature Preserve, Mason County, Illinois

William E. McClain¹, John E. Schwegman², Todd A. Strole³, Loy R. Phillippe⁴, and John E. Ebinger⁴

¹Adjunct Research Associate in Botany, Illinois State Museum, Springfield, Illinois 62706-5000, ²3626 Riverpoint Lane, Metropolis, Illinois 62960, ³Illinois Department of Natural Resources, Springfield, Illinois 72702, ⁴Illinois Natural History Survey, Champaign, Illinois 61820.

ABSTRACT

The 590 ha Sand Prairie-Scrub Oak Nature Preserve in Mason County, Illinois contains remnant sand prairies, sand savanna, sand forest, and successional fields. Purchased in 1969, these abandoned agricultural fields have been allowed to re-vegetate naturally. During the present study the vegetation of the mature dry sand prairie was compared with two successional fields, one 60- and one 30-years-old. The mature dry sand prairie remnants are dominated by *Schizachyrium scoparium* (Michx.) Nash, *Dichanthelium villosissimum* (Nash) Freckm., *Tephrosia virginiana* (L.) Pers, *Ambrosia psilostachya* DC., and *Opuntia humifusa* (Raf.) Raf. Though these five species are usually present, *Eragrostis trichodes* (Nutt.) Wood dominated the 60-year-old successional fields. The 30-year-old successional field was dominated by *Eragrostis trichodes*, *Strophostyles helvula* (L.) Ell., and *Monarda punctata* L. A total of 393 vascular plant species were documented for the preserve.

INTRODUCTION

Sand prairie communities are present in Illinois on windblown sands deposits located along the Illinois, Kankakee, Green, and Mississippi Rivers, and the shores of Lake Michigan (Gleason 1910, Vestal 1913). Due to their arid nature, they were the last of the extensive Illinois prairie to be converted to agriculture. In the early 1900's, thousands of acres of sand prairie still remained (Hart and Gleason 1907, Gleason 1910, Sampson 1921). With the advent of central pivot irrigation, much of the remaining sand prairie was converted to agriculture between 1940 and 1960. The majority of the natural areas that remained were unsuitable for cultivation due to extensive sand dunes, large areas of actively moving sand known as blowouts, or depressions difficult to drain.

The most extensive sand deposit in Illinois are those associated with the Illinois River in the central part of the state. Within these deposits the Illinois Nature Preserves Commission has preserved many of the best remaining examples of the sand communities that were once the dominant vegetation of the region. To help accomplish this goal the Illinois Department of Natural Resources purchased the Sand Prairie-Scrub Oak Nature Preserve (SP-SONP). When purchased, much of the preserve consisted of sand forest, open woodlands, and savanna though some high quality sand prairie was also present along with numerous abandoned agricultural fields.

Since being purchased in 1969 the SP-SONP has been used for prairie vegetation studies. The mycorrhizal dependence of two sand prairie grasses was examined (Anderson and Liberta 1987, Dhillion et al. 1988,

Benjamin et al. 1989, Anderson et al. 1994). Also, the effects of fire on savanna and adjacent forest vegetation were studied (Anderson and Brown 1986, Dhillon and Anderson 1994), while recently the composition and structure of the sand forest communities was examined (McClain et al. 2002). Except for the early works by Hart and Gleason (1907) and Gleason (1910), however, little information has been published on the composition of the sand prairie communities. High quality dry sand prairie remnants occur on the SP-SONP as well as various aged abandoned fields. The purpose of this study was to compare the plant species composition, vegetation structure, and floristic quality of a dry sand prairie remnant with various aged old-field communities, and to document the vascular flora of the SP-SONP.

DESCRIPTION OF THE STUDY AREA

The SP-SONP is a 590 ha site located between the villages of Bath and Kilbourne (T20N, R9W, S13, 14, 23, 26,) Mason County, in the Illinois River Section of the Illinois River and the Mississippi River Sand Areas Natural Division (Schwegman 1973). This sand deposit was created about 14,500 years ago when glacial moraines were breached, resulting in the Kankakee Torrent (Willman 1973, King 1981). These floodwaters carried huge amounts of sand and gravel that were deposited when the waters of the Kankakee Torrent slowed upon entering the broad lowlands of the Illinois River below present day Hennepin, Illinois. These sand deposits were subsequently shaped by winds, creating the dune and swale topography known as the Parkland Formation (Willman and Frye 1970). The soils of the SP-SONP are excessively drained Plainfield sands (Calsyn 1995). These soils consist of about 95% sand, 4% silt, and 1% clay. Soil pH ranges from 5.1 to 5.3 and is low in organic matter and nitrogen (Benjamin et al. 1989).

Within the SP-SONP many sand communities exist. Presently the most common are sand forests that were probably open woodlands and savannas during early settlement time (Rodgers and Anderson 1979). At the time of purchase a few remnant sand prairies were present based on a United States Department of Agriculture aerial photograph dated 9 July 1939. These remnants were probably grazed, but there is no indication that they were ever cultivated. One area, originally considered to be a prairie remnant, was determined to be in row crop agriculture according to the 1939 photograph. Subsequent aerial photographs indicate that farming of this site was abandoned in the early 1940's (60-year-old successional field). Other fields were taken out of agriculture at the time of purchase in 1969 (30-year-old successional fields).

The only management activities conducted on the SP-SONP have been prescribed burning to keep the forests open, and invasive exotic species control to eliminate *Robinia pseudoacacia* (black locust) from the prairies and old fields. The north side of the preserve, the location of the study sites, was prescribed burned in 1994. Prior to this date, prescribed fire was used in smaller areas in the interior of the preserve in 1984, 1987, 1990, and 1992 mostly in the sand forest vegetation.

Climate at the SP-SONP is continental with warm summers and cold winters. Based on weather data from Havana, Illinois about 12 km north of the SP-SONP, mean annual precipitation is 96.0 cm, with May having the highest rainfall (11.3 cm). Mean annual temperature is 10.8°C with the hottest month being July (average of 24.6°C), and the coldest

January (average of -5.0°C). Frost free days range from 140 to 206, with the average being 173 day per year (Midwestern Regional Climate Center 2002).

METHODS

The SP-SONP was visited throughout the growing season of 1976 by one of the authors (Schwegman 1977). The preserve has been visited many times since this initial study, with a concentrated effort during the 2000 to 2003 growing season. During these trips specimens were collected and deposited in the herbarium of the Illinois Natural History Survey, Champaign, Illinois (ILLS), and the Stover-Ebinger Herbarium, Eastern Illinois University, Charleston, Illinois (EIU). Gleason and Cronquist (1991) and Mohlenbrock (1986) were used to designating non-native species. Nomenclature follows Mohlenbrock (1986).

In late summer of 2000 two 25 m transects were located randomly along cardinal compass directions in a mature dry sand prairie remnant, a 60-year-old, and a 30-year-old successional field. Along each transect, 1 m² quadrats were located at 1 m intervals ($n=25/\text{transect}$). Odd-numbered quadrats were located on the right side; even-numbered to the left. A random numbers table was used to determine the number of meters (0 to 9) the quadrat was located from the transect line. Species cover was determined using the Daubenmire cover class system (Daubenmire 1959) as modified by Bailey and Poulton (1968). Importance value (IV) was determined by summing relative cover and relative frequency.

RESULTS

A total of 387 species within 235 genera and 76 families were documented on the preserve (Appendix I). Fern and fern-allies were represented by only one species, while gymnosperms accounted for two. Of the remainder, 108 were monocots in 11 families and 61 genera and 282 were dicots in 64 families and 182 genera. Non-native (exotic) species accounted for 71 taxa, about 19% of the species collected. The predominant plant families were the Asteraceae with 65 species and the Poaceae with 59. The state endangered *Echinodorus tenellus* and *Schoenoplectus hallii* were encountered at the edge of a sand pond, while the state threatened *Cyperus grayoides* was common in disturbed sand prairies and blowouts (Herkert and Ebinger 2002).

Some of the species associated with wet areas reported by Schwegman (1977) were found around a sand pond located in a large blowout near the middle of the SP-SONP. This was a sizeable pond prior to the lowering of the water table by drainage of the land surrounding the preserve. Presently water rarely accumulates in this blowout and these species many no longer occur on the preserve. It is possibly that they persist within the soil seed bank, however, and will appear in extremely wet years (McClain et al. 1997).

Mature Dry Sand Prairie: Of the species encountered *Schizachyrium scoparium* (little bluestem) was most important, having an average cover of 46.35, and an IV of 79.4 (Table 1). Also common, *Dichanthelium villosissimum* (hairy panic grass) was second with an IV of 30.6, *Tephrosia virginiana* (goat's-rue) was third with an IV of 28.3, followed by *Ambrosia psilostachya* (western ragweed), and *Opuntia humifusa* (common prickly pear). These five native prairie species, that are typical components of dry sand prairies in the Mason County sand deposits, had IV's greater than 10. All would be expected in high

quality dry sand prairie communities in Illinois. Of the 16 species encountered in the plots, 14 were perennials (Table 1).

60-year-old Successional Field: Last cultivated about 60 years ago this old-field contains many sand prairie species. *Eragrostis trichodes* (sand love grass), that was absent from the mature dry sand prairie was dominant with an average cover of 25.00, and an IV of 51.2. After 60 years this area contained many taxa commonly associated with mature sand prairies. Four of the top five species in IV (little bluestem, western ragweed, hairy panic grass, and common prickly pear) were among the top five encountered in the mature dry sand prairie (Table 1). Other important differences were the relatively low IV for little bluestem, and the absence of goat's-rue. Of the 20 herbaceous species encountered in the plots, 17 were perennials (Table 1).

30-year-old Successional Field: This field, taken out of cultivation when the preserve was purchased, contained many species not found in the mature sand prairie. Sand love grass was the dominant with an average cover of 61.70 and an IV of 97.4, while the herbaceous vine *Strophostyles helvula* (wild bean) was second with an IV of 43.7, and *Monarda punctata* (horsemint) was third with an IV of 16.9 (Table 1). Of the dominants associated with mature dry sand prairies, little blue stem, western ragweed, and goat's-rue were not encountered while hairy panic grass and common prickly pears were rare. Of the 22 species encountered in the plots, 18 were perennials (Table 1).

DISCUSSION

In the Illinois River sand deposits native sand prairie species rapidly colonize agricultural fields, especially in areas with a readily available native seed source. Most of these early successional species, however, are those associated with natural sand area disturbances such as blowouts or are native adventive species. After 30 years, however, few of the species associated with mature dry sand communities were found in the plots. Colonization by characteristic herbaceous species such as little bluestem, hairy panic grass, goat's-rue, and western ragweed require much longer, particularly when sites re-vegetate naturally. This slow colonization is not due to the lack of a seed source as native prairie is adjacent to the fields. This slow rate of colonization is not completely understood. It may be related to disruption of the mycorrhizal community in the soil, lack of certain soil nutrients, or competition from aggressive successional species such as sand love grass (Anderson and Liberta 1987, Dhillion et al. 1988, Anderson et al. 1994).

Though 71 non-native species were encountered at the SP-SONP, most were restricted to roadsides, fence rows, or in the remnants of pine plantations and abandoned home sites. Also, some were encountered around the "blowouts", others associated with past off-road vehicle disturbances. Most of the non-native species are not management problems. The major exceptions are black locust that is highly invasive in remnant prairies and old fields, and *Lonicera maackii* (Amur honeysuckle) and *Alliaria petiolata* (garlic mustard) which are serious invaders of the sand forests. The only exotic found in the study plots was *Mollugo verticillata* (carpetweed) which had an IV of 0.5 in the 30-year-old successional field.

ACKNOWLEDGMENTS

The authors thank Dr. Gordon Tucker, Eastern Illinois University for his help in the identification of the Cyperaceae, and John Wilker, Natural Heritage Biologist, Illinois Department of Natural Resources for his help and advice. We also thank the Illinois Wildlife Preservation Fund for their generous grant that helped support this project.

LITERATURE CITED

Anderson R.C. and L.E. Brown. 1986. Stability and instability in plant communities following fire. *American Journal of Botany* 73:364-368.

Anderson, R.C., B.A.D. Hetrick, and G.W.T. Wilson. 1994. Mycorrhizal dependence of *Andropogon gerardii* and *Schizachyrium scoparium* in two prairie soils. *American Midland Naturalist* 132:366-376.

Anderson, R.C. and A.E. Liberta. 1987. Variation in vesicular-arbuscular mycorrhizal relationships of two sand prairie species. *American Midland Naturalist* 127:56-63.

Benjamin, P., R.C. Anderson, and A.E. Liberta. 1989. Vesicular-arbuscular mycorrhizal ecology of little bluestem across a prairie-forest gradient. *Canadian Journal of Botany* 67:2678-2685.

Bailey, A.W., and C.E. Poulton. 1968. Plant communities and environmental relationships in a portion of the Tillamook burn, northwestern Oregon. *Ecology* 49:1-13.

Calsyn, D.E. 1995. Soil survey of Mason County, Illinois. Soil Report 146, University of Illinois Agricultural Experiment Station, Urbana. ix+211 pp.

Daubenmire, R. 1959. A canopy coverage method of vegetation analysis. *Northwest Science* 33:43-64.

Dhillion S.S., R.C. Anderson, and A.E. Liberta. 1988. Effects of fire on the mycorrhizal ecology of little blue stem (*Schizachyrium scoparium*). *Canadian Journal of Botany* 66:706-713.

Dhillion, S.S. and R.C. Anderson. 1994. Production on burned and unburned sand prairies during drought and non-drought years. *Vegetatio* 115:51-59.

Gleason, H.A. 1910. The vegetation of the inland sand deposits of Illinois. *Bulletin of the Illinois State Laboratory of Natural History* 9:21-174.

Gleason, H.A., and A. Cronquist. 1991. Manual of the vascular flora of northeastern United States and adjacent Canada. Second Edition. The New York Botanical Garden, Bronx, New York. lxxv+910 pp.

Hart, C.A. and H.A. Gleason. 1907. On the biology of the sand areas of Illinois. *Bulletin of the Illinois State Laboratory of Natural History* 7:137-272.

Herkert, J.R. and J.E. Ebinger. 2002. editors. *Endangered and Threatened Species of Illinois: Status and Distribution. Volume 1:*

Plants. Endangered Species Protection Board, Springfield, Illinois. 161 pp.

King, J.E. 1981. Late Quaternary vegetational history of Illinois. *Ecological Monographs* 51:43-62.

McClain, W.E., R.D. McClain, and J.E. Ebinger. 1997. Flora of temporary sand ponds in Cass and Mason counties, Illinois. *Castanea* 62:65-73.

McClain, W.E., S.D. Turner, and J.E. Ebinger. 2002. Vegetation of forest communities at the Sand Prairie-Scrub Oak Nature Preserve, Mason County, Illinois. *Transactions of the Illinois State Academy of Science* 95:37-46.

Midwestern Regional Climate Center. 2002. <http://mcc.sws.uiuc.edu>

Mohlenbrock, R.H. 1986. Guide to the Vascular Flora of Illinois, revised and enlarged edition, Southern Illinois University Press, Carbondale, Illinois. vii+507 pp.

Rodgers, C.S. and R.C. Anderson. 1979. Presettlement vegetation of two prairie peninsula counties. *Botanical Gazette* 140:232-240.

Sampson, H.C. 1921. An ecological survey of the prairie vegetation of Illinois. *Bulletin of the Illinois State Laboratory of Natural History* 13:523-577.

Schwegman, J.E. 1973. Comprehensive plan for the Illinois nature preserves system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, Illinois. map+32 pp.

Schwegman, J.E. 1977. An annotated list of the flowering plants of Sand Prairie-Scrub Oak Nature Preserve. Unpublished report submitted to the Illinois Department of Conservation, Springfield, Illinois map+26pp.

Vestel, A.G. 1913. An association study of Illinois sand prairie. *Bulletin of the Illinois State Laboratory of Natural History* 10:1-96.

Willman, H.B. 1973. Geology along the Illinois waterway - a basis for environmental planning. Illinois State Geological Survey Circular 478. Urbana. 48 pp.

Willman, H.B. and J.C. Frye. 1970. Pleistocene stratigraphy of Illinois. *Illinois State Geological Survey Bulletin* 94:1-204.

Table 1. Importance values (I.V.) and average cover of the ground layer species encountered in a mature sand prairie, and a 60, and a 30 year successional field at the Sand Prairie Scrub Oak Nature Preserve, Mason County, Illinois. Species with I.V.'s of 1.0 or less and found in only one of the areas are included as others.

Species	Mature Prairie		60 Year Field		30 Year Field	
	I.V.	Average Cover	I.V.	Average Cover	I.V.	Average Cover
<i>Schizachyrium scoparium</i>	79.4	46.35	31.7	14.83	--	--
<i>Dichanthelium villosissimum</i>	30.6	9.22	21.2	7.20	1.2	0.07
<i>Tephrosia virginiana</i>	28.3	10.51	--	--	--	--
<i>Ambrosia psilostachya</i>	22.2	6.74	23.2	7.23	--	--
<i>Opuntia humifusa</i>	15.2	3.12	18.1	6.22	0.6	0.06
<i>Eragrostis spectabilis</i>	6.0	0.91	9.5	2.70	10.6	2.29
<i>Cyperus schweinitzii</i>	5.6	0.33	4.6	0.23	--	--
<i>Calamovilfa longifolia</i>	3.7	0.09	--	--	--	--
<i>Crotonopsis linearis</i>	2.1	0.05	--	--	--	--
<i>Oenothera rhombipetala</i>	1.8	0.19	4.0	0.31	1.1	0.02
<i>Cyperus grayoides</i>	1.7	0.09	0.3	0.01	--	--
<i>Carex muhlenbergii</i>	1.0	0.12	2.3	0.90	--	--
<i>Pseudognaphalium obtusifolium</i>	0.5	0.06	3.5	1.08	0.6	0.06
Others (3 species)	1.9	0.19	--	--	--	--
<i>Eragrostis trichodes</i>	--	--	51.2	25.00	97.4	61.70
<i>Froelichia floridana</i>	--	--	11.1	2.24	--	--
<i>Conyza canadensis</i>	--	--	4.1	0.36	1.8	0.18
<i>Paspalum bushii</i>	--	--	3.9	0.91	9.8	1.00
<i>Croton glandulosus</i>	--	--	3.8	0.16	--	--
<i>Chenopodium simplex</i>	--	--	3.2	0.14	1.1	0.02
<i>Physalis heterophylla</i>	--	--	2.0	0.25	4.8	0.90
<i>Monarda punctata</i>	--	--	0.7	0.02	16.9	4.06
Others (2 species)	--	--	1.6	0.14	--	--
<i>Strophostyles helvula</i>	--	--	--	--	43.7	15.36
<i>Tridens flavus</i>	--	--	--	--	2.5	0.38
<i>Asclepias syriaca</i>	--	--	--	--	2.4	0.24
<i>Ampelamus albidus</i>	--	--	--	--	1.1	0.02
Others (7 species)	--	--	--	--	4.4	0.75
Totals	200.0	77.97	200.0	69.93	200.0	87.11

Asclepias amplexicaulis Small 29810
Asclepias syriaca L. 29934
Asclepias tuberosa L. 29809
Asclepias verticillata L. 29935

Asteraceae

**Achillea millefolium* L. 29811
Ageratina altissima (L.) R.M. King & H. Robins. 29937
Ambrosia artemisiifolia L. 25964
Ambrosia psilotachya DC. 30384
Ambrosia trifida L. 30385
Antennaria plantaginifolia (L.) Hook. 29936
Arnoglossum atripicifolium (L.) H. Robins. 30207
Artemisia campestris L. 30203
Aster cordifolius L. 31203
Aster dumosus L. (JES)
Aster ericoides L. 30483
Aster lanceolatus Willd. 30481
Aster lateriflorus (L.) Britt. 31202
Aster ontarionis Wieg. (JES)
Aster oolentangiensis Riddell 30484
Aster pilosus L. 30386
Aster racemosus L. 30482
Aster sagittifolius Willd. (JES)
Aster urophyllus Lindl. 25989
Bidens bipinnata L. 25950
Brickellia eupatorioides (L.) Shinnery 30387
**Carduus nutans* L. 29812
Chrysopsis camporum Greene 29528
Cirsium discolor (Muhl.) Spreng. 30388
Conoclinium coelestinum (L.) DC. 30390
Conyza canadensis (L.) Cronq. 30202
Coreopsis lanceolata L. 29678
Coreopsis palmata Nutt. 30208
Erechtites hieracifolia (L.) Raf. 30389
Erigeron annuus (L.) Pers. 29814
Erigeron strigosus Muhl. 29813
Eupatoriadelphus purpureus (L.) R.M. King & H. Robins. 30485
Eupatorium altissimum L. 30478
Eupatorium serotinum Michx. 30391
Euthamia graminifolia (L.) Nutt. 30392
Euthamia gymnospermoides Greene P36125
Helianthus divaricatus L. 25990
Helianthus grosseserratus Martens. (JES)
Helianthus hirsutus Raf. 30394
Helianthus occidentalis Riddell

25987
Helianthus pauciflorus Nutt. 30206
**Helianthus petiolaris* Nutt. 30205
Heliopsis helianthoides (L.) Sweet 29938
Heterotheca subaxillaris (Lam.) Britt. & Rusby 31000
Hieracium scabrum Michx. 25992
Ionactis linariifolius (L.) Greene P36123
Krigia virginica (L.) Willd. 29524
Lactuca canadensis L. 26006
Lactuca floridana (L.) Gaertn. (JES)
**Lactuca serriola* L. 29939
Liatris aspera Michx. 30395
Pseudognaphalium obtusifolium (L.) Hilliard & Burt. 30393
Pyrrhopappus carolinianus (Walt.) DC. P36119
Ratibida pinnata (Vent.) Barnh. 29940
Rudbeckia hirta L. 29815
Senecio plattensis Nutt. 29529
Solidago canadensis L. 30396
Solidago gigantea Ait. (JES)
Solidago missouriensis Nutt. 30995
Solidago nemoralis Ait. 30204
Solidago speciosa Nutt. (JES)
Solidago ulmifolia Muhl. P36100
Taraxacum officinale Weber 30201
**Tragopogon dubius* Scop. 29703
Xanthium strumarium L. 30480

Bignoniaceae

Campsis radicans (L.) Seem. 30209
**Catalpa speciosa* Warder 29941

Boraginaceae

**Buglossoides arvense* (L.) I.M. Johnston 29439
Hackelia virginiana (L.) I.M. Johnston 29942
Lithospermum canescens (Michx.) Lehm. (JES)
Lithospermum croceum Fern. 29527
Lithospermum incisum Lehm. 29526

Brassicaceae

**Alliaria petiolata* (Bieb.) Cavara & Grande 29523
Arabis canadensis L. 29687
Descurainia pinnata (Walt.) Britt. 29438
Draba reptans (Lam.) Fern. 29441
Erysimum capitatum (Dougl.) Greene 29531
**Lepidium densiflorum* Schrad. 29698

Rorippa sessiliflora (Nutt.) A.
Hitchc. 30855

Cactaceae

Opuntia humifusa (Raf.) Raf. 30376

Caesalpinaceae

Chamaechrista fasciculata (Michx.)
Greene 30210

Chamaechrista nictitans (L.) Moench.
25275

Senna marilandica (L.) Link 30211

Campanulaceae

Lobelia spicata Lam. (JES)

Triodanis perfoliata (L.) Nieuwl.
29816

Cannabaceae

**Cannabis sativa* L. 29817

Caprifoliaceae

**Lonicera maackii* (Rupr.) Maxim.
30212

Sambucus canadensis L. 30856

Caryophyllaceae

**Arenaria serpyllifolia* L. 29536

**Cerastium glomeratum* Thuill. 29535

**Dianthus armeria* L. 30999

**Holosteum umbellatum* L. 29444

Paronychia canadensis (L.) Wood
25276

Paronychia fastigiata (Raf.) Fern.
30397

**Saponaria officinalis* L. 29818

Silene antirrhina L. 29943

**Silene pratensis* (Spreng.) Godron &
Gren. 29671

Silene stellata (L.) Ait. f. 29944

**Stellaria media* (L.) Cyrillo 29537

Celastraceae

Celastrus scandens L. 31199

Chenopodiaceae

**Chenopodium album* L. 30398

Chenopodium desiccatum A. Nels.
25969

Chenopodium simplex (Torr.) Raf.
25945

Cycloloma atriplicifolium (Spreng.)
Coults. 29946

Cistaceae

Helianthemum bicknellii Fern. 30998

Helianthemum canadense (L.) Michx.

29673

Lechea tenuifolia Michx. (JES)

Convolvulaceae

**Ipomoea hederacea* (L.) Jacq. 25953

Cornaceae

Cornus drummondii C.A. Mey. 29819

Cornus racemosa Lam. 30213

Cuscutaceae

Cuscuta gronovii Willd. 30399

Euphorbiaceae

Acalypha gracilens Gray 29947

Acalypha rhomboidea Raf. P36129

Chamaesyce geyeri (Engelm.) Small
25959

Chamaesyce maculata (L.) Small 30486

Croton glandulosus L. 25961

Crotonopsis linearis Michx. 25962

Euphorbia corollata L. 29948

Poinsettia dentata (Michx.) Kl. &
Garcke 25960

Fabaceae

Amorpha canescens Pursh 29949

Amphicarpaea bracteata (L.) Fern.
30400

Astragalus distortus Torr. & Gray
29540

Baptisia bracteata Ell. P36116.2

Dalea candida (Michx.) Willd. (JES)

Desmodium glabellum (Michx.) DC.
30401

Desmodium illinoense Gray 30215

Desmodium sessilifolium (Torr.) Torr.
& Gray 26005

**Kummerowia stipulacea* (Maxim.)
Makino 29821

Lespedeza capitata Michx. 30402

**Medicago lupulina* L. 29823

**Melilotus alba* Medic. 29820

**Melilotus officinalis* (L.) Pallas
225971

**Robinia pseudoacacia* L. 29950

Strophostyles helvula (L.) Ell.
25966

Strophostyles leiosperma (Torr. &
Gray) Piper 30214

Tephrosia virginiana (L.) Pers.
25278

**Trifolium campestre* Schreb. 29822

**Trifolium pratense* L. 29824

**Trifolium repens* L. 29825

**Vicia villosa* Roth. 29672

Fagaceae

- Quercus x bushii* Sarg. 29951
Quercus marilandica Muenchh. 29952
Quercus velutina Lam. 29953

Fumariaceae

- Corydalis micrantha* (Engelm.) Gray
(JES)

Gentianaceae

- Gentiana alba* Muhl. (JES)

Geraniaceae

- Geranium carolinianum* L. 29681

Grossulariaceae

- Ribes missouriense* Nutt. 29685

Hypericaceae

- Hypericum majus* (Gray) Britt. (JES)
Hypericum mutilum L. (JES)
Hypericum sphaerocarpum Michx.
P36091

Juglandaceae

- Carya texana* Buckl. 29954
Carya tomentosa (Poir.) Nutt. 29955
Juglans nigra L. 29956

Lamiaceae

- Agastache nepetoides* (L.) Ktze.
30216
**Lamium amplexicaule* L. 29446
**Leonurus cardiaca* L. 29826
Lycopus americanus Muhl. 30403
Monarda fistulosa L. 29827
Monarda punctata L. 30217
Physostegia virginiana (L.) Benth.
P36122
Pycnanthemum pilosum Nutt. 25988
Scutellaria parvula Michx. P36077
Stachys palustris L. P36120
Teucrium canadense L. 25277

Lauraceae

- Sassafras albidum* (Nutt.) Nees 29828

Lythraceae

- Rotala ramosior* (L.) Koehne 30997

Malvaceae

- Callirhoe triangulata* (Leavenw.) Gray
29957

Molluginaceae

- **Mollugo verticillata* L. 25972

Moraceae

- **Maclura pomifera* (Raf.) Schneider
29958
**Morus alba* L. 29829

Nyctaginaceae

- **Mirabilis nyctaginea* (Michx.) MacM.
29702

Onagraceae

- Gaura biennis* L. 30404
Ludwigia alternifolia L. 29959
Oenothera biennis L. 30405
Oenothera laciniata Hill 29680
Oenothera rhombipetala Nutt. 25967

Oxalidaceae

- Oxalis stricta* L. 29686
Oxalis violacea L. (JES)

Phytolaccaceae

- Phytolacca americana* L. 29960

Plantaginaceae

- Plantago aristata* Michx. (JES)
**Plantago lanceolata* L. 29831
Plantago patagonica Jacq. 29692
Plantago rugelii Dcne. 29832
Plantago virginica L. 29676

Polemoniaceae

- Phlox bifida* Beck 29440

Polygalaceae

- Polygala polygama* Walt. 25958
Polygala sanguinea L. P36096
Polygala verticillata L. (JES)

Polygonaceae

- Antenoron virginianum* (L.) Roberty &
Vautier 30410
**Fallopia convolvulus* (L.) A. Love
29961
Fallopia cristata (Engelm. & Gray)
Holub. 25808
Fallopia scandens (L.) Holub. 30409
Persicaria coccinea (Muhl.) Greene
29962
Persicaria lapathifolia (L.) S.F.
Gray 29963
Persicaria pensylvanica (L.) Small
30407
**Persicaria vulgaris* Webb & Moq.
30408
**Polygonum aviculare* L. 30406
Polygonum tenue Michx. 26003
**Rumex acetosella* L. 29675

**Rumex crispus* L. 29830

Portulacaceae

Claytonia virginica L. (JES)
Talinum rugospermum Holz. 29833

Primulaceae

Androsace occidentalis Pursh 29445
Lysimachia ciliata L. (JES)

Ranunculaceae

Anemone canadensis L. P36130
Anemone caroliniana Walt. 29442
Anemone cylindrica Gray (JES)
Anemone virginiana L. 29964
Ranunculus abortivus L. 29522

Rhamnaceae

Ceanothus americanus L. 25279

Rosaceae

Agrimonia gryposepala Wallr. P36131
Fragaria virginiana Duchesne 29534
Geum canadense Jacq. 25812
Malus ioensis (Wood) Britt. (JES)
**Potentilla norvegica* L. 30857
**Potentilla recta* L. 29834
Potentilla simplex Michx. 29697
Prunus angustifolia Marsh. 29516
Prunus serotina Ehrh. 29517
Rosa carolina L. 29835
**Rosa multiflora* Thunb. 29965
Rubus flagellaris Willd. 29684
Rubus occidentalis L. 29691
Rubus pensylvanicus Poir. 26601

Rubiaceae

Diodia teres Walt. 25963
Galium aparine L. 29533
Galium circaezans Michx. 29836
Galium pilosum Ait. 25273

Rutaceae

Zanthoxylum americanum Mill. 29966

Salicaceae

**Populus alba* L. P36118
Populus deltoides Marsh. 29967
Salix humilis Marsh. 25810
Salix nigra Marsh. 30487

Santalaceae

Comandra umbellata (L.) Nutt. 29695

Saxifragaceae

Penthorum sedoides L. (JES)

Scrophulariaceae

Lindernia dubia (L.) Pennell (JES)
Nuttallanthus canadensis (L.) Sutton
29532
Penstemon pallidus Small 29696
Scrophularia lanceolata Pursh 29693
**Verbascum thapsus* L. 29837
**Veronica arvensis* L. 29538
Veronica peregrina L. 29677
Veronicastrum virginicum (L.) Farw.
30219

Solanaceae

Physalis heterophylla Nees 25970
Physalis virginiana Mill. 30488
Solanum carolinense L. 29968
Solanum ptychanthum Dunal. 25949

Ulmaceae

Celtis occidentalis L. 29969
Ulmus americana L. 29970
**Ulmus pumila* L. 29971
Ulmus rubra Muhl. P36127

Urticaceae

Parietaria pensylvanica Muhl. 25272
Urtica gracilis Ait. 30411

Verbenaceae

Verbena hastata L. 30996
Verbena stricta Vent. 29972
Verbena urticifolia L. 30218

Violaceae


Viola pedata L. 29682
Viola pedatifida G. Don (JES)
**Viola rafinesquii* Greene 29443
Viola sagittata Ait. P36098
Viola sororia Willd. (JES)

Vitaceae

Parthenocissus quinquefolia (L.)
Planch. 29973
Vitis aestivalis Michx. 29974
Vitis vulpina L. 29838

Zygophyllaceae

**Tribulus terrestris* L. 29975



VEGETATION SURVEY OF TOMLIN TIMBER NATURE PRESERVE, MASON COUNTY,
ILLINOIS

Loy R. Phillippe, James Ellis, Daniel T. Busemeyer, William E. McClain,
and John E. Ebinger
Illinois Natural History Survey, Champaign, Illinois 61820

ABSTRACT: Tomlin Timber Nature Preserve is a dry to dry-mesic sand forest. When surveyed by the Illinois Natural Areas Inventory in 1976, it was dominated by *Quercus velutina* (black oak), *Sassafras albidum* (sassafras) was second in importance and dominated the woody seedling and sapling layer and was common in the small tree diameter classes. In 2004 sassafras dominated the overstory with an importance value (IV) of 54.9 (possible 200), an average dbh of 24.4 cm, and with most individuals in the 10-40 cm diameter classes. Black oak, in contrast, was second with an IV of 38.5, an average dbh of 62.7, and dominated the larger diameter classes. Dead-standing and dead-downed black oaks were common and averaged 51.7 stems/ha. The change in dominance appears to be due to a combination of natural mortality and oak wilt, which was common in the preserve in the early 1980s. Over 157 vascular plant taxa were found in the preserve.

INTRODUCTION

Wind-blown sand deposits, which account for nearly 5% of the land surface of Illinois, occurs on glacial outwash plains associated with erosional events of Wisconsin glaciation in the northern half of the state (Schwegman 1973, King 1981). The Illinois River sand deposits of Cass and Mason counties in the central part of the state, were formed about 14,500 years ago when glacial moraines were breached, resulting in the Kankakee Torrent (Willman and Frye 1970). These flood waters carried huge amounts of sand and gravel that were deposited when the waters slowed upon entering the broad lowlands of the Illinois River valley. Many high quality natural areas that are now Illinois Nature Preserves occur on these sand deposits. We are studying these preserves to determine the composition and structure of the plant communities.

Early works by Hart and Gleason (1907) and Gleason (1910) were the first extensive studies of the sand area vegetation of the state. A few other studies were completed prior to the early 1990s, those being plant species lists for a few natural areas (Maier 1976, Schwegman 1977). More recently the structure and composition of some upland dry sand forest communities and dry sand savanna communities were examined (Jenkins et al. 1991, Coates et al. 1992, McClain et al. 2002). The purpose of the present study was to examine the composition and structure of Tomlin Timber Nature Preserve, a sand forest remnant on the Illinois River sand deposits in Mason County.

DESCRIPTION OF THE STUDY SITE

Dedicated as a nature preserve in 1987, Tomlin Timber is located about 3 km south of Easton, Mason County, Illinois (NE1/4 SW1/4 S11 T20N R7W). The preserve is on a series of low dunes that lie between 158 and 165 m above sea level in the Illinois River Section of the Mississippi River and Illinois Rivers Sand Areas Natural Division (Schwegman 1973). The soils are excessively drained Bloomfield sand (Calsyn 1995); part of the dune and swale topography known as the Parkland Formation (Willman and Frye 1970).

Little is known about the past history of this 8 ha tract of timber. A comment by Onstot (1902) mentions that Walker's Grove (of which Tomlin Timber is a remnant) "embraces 40 acres of as fine a body of timber as can be found anywhere; a fine growth of oaks, black walnut, soft and sugar maple, hickory, butternut, mulberry, sassafras, redbud, pawpaw, and dogwood". The size of the black oaks, most are in the 50-70 cm diameter classes, suggests that the woods was never clear-cut. When purchased by the Tomlin family in 1912 a one-room schoolhouse was located on the east edge of the property. When surveyed by the Illinois Natural Areas Inventory (White 1978), the owner indicated that the woods had been selectively logged 50 to 60 years ago, but never grazed. Don McFall (Illinois Department of Natural Resources, personal communication) mentions that he first walked through the woods in the early 1980s, and there was a fairly dense woody understory and a number of large dead black oaks.

Based on weather data from Havana about 21 km to the northwest, mean annual precipitation is 96.0 cm, with May having the highest rainfall (11.3 cm). Mean annual temperature is 10.8°C with the hottest month being July (average of 24.6°C), and the coldest January (average of -5.0°C). Frost-free days range from 140 to 206, with the average being 173 days per year (Midwestern Regional Climate Center 2004).

METHODS

During the growing seasons of 2003 and 2004 the study site was visited 12 times. All vascular plant species were collected, their habitat recorded, and voucher specimens deposited in the herbarium of the Illinois Natural History Survey, Champaign, Illinois (ILLS). Criteria for designating non-native species follows Mohlenbrock (2002) and Gleason and Cronquist (1991). Nomenclature follows Mohlenbrock (2002). All species are listed in Appendix I along with the collector and collecting number.

During late summer of 2004 a 150 m by 300 m section of the natural area was surveyed by dividing the area into 72 contiguous quadrates 25 m on a side (4 ha area). The GPS coordinates for the corners of the plot are listed in Appendix II. All living and dead-standing woody individuals >10.0 cm dbh were identified and diameters recorded. From this data, living-stem density (stems/ha), basal area (m²/ha), relative density, relative dominance, importance value (IV), and average diameter (cm) were calculated for each species. Determination of the IV follows the procedure used by McIntosh (1957), and is the sum of the relative density and relative dominance (basal area). Dead-standing density (stem/ha) and basal area (m²/ha) was also determined.

Woody understory composition and density (stems/ha) was determined using nested circular plots 0.0001, 0.001, and 0.01 ha in size located at 15 m intervals along randomly located east-west transects within the study area (72 plots). Four additional 0.0001 ha circular plots were located 7 m from the center points of each of the 72 plot centers along cardinal compass directions (360 plots). In the 0.0001 ha plots, woody seedlings (<50 cm tall) were counted; in the 0.001 ha circular plots small saplings (>50 cm tall and <2.5 cm dbh) were recorded; and in the 0.01 ha circular plots large saplings (2.5-9.9 cm dbh) were tallied.

RESULTS

At Tomlin Timber Nature Preserve a total of 157 vascular plant species within 125 genera and 61 families were documented. Of these, three were fern and fern-allies, one was a gymnosperms, 122 dicots in 98 genera and 50 families, and 31 monocots in 23 genera and seven families. No threatened or endangered species were found. The predominant plant families were the Poaceae and Asteraceae with 17 taxa each. Thirty one exotic plant taxa were found, about 19% of the flora of the preserve (Appendix I).

Nineteen tree species were present in the overstory (Table 1). *Sassafras albidum* (sassafras) dominated the smaller diameter classes with most individuals being less than 40 cm dbh. This species had an IV of 54.9, averaged 132.5 stems/ha, and averaged 24.4 cm dbh. *Quercus velutina* (black oak), restricted to the larger diameter classes, was second in IV (38.5), averaged 25.4 stems/ha, and averaged 62.7 cm dbh. The remaining trees were mostly in the 10-39 cm diameter classes, *Carya texana* (black hickory) averaged 90.1 stems/ha, *Celtis occidentalis* (hackberry) averaged 80.7 stems/ha, and *Prunus serotina* (black cherry) averaged 52.8 stems/ha, all with IVs below 31 and average diameters less than 20 cm dbh.

The woody understory was dense with 18,639 woody seedlings/ha, 4,862 small saplings/ha, and 1,222 large saplings/ha (Table 2). Few open areas existed in the woody understory, the more open areas being under the extensive colonies of *Asimina triloba* (pawpaw) (Table 2). Pawpaw averaged 4,028 seedlings/ha, 2,986 small saplings/ha, 854 large saplings/ha, along with 14.9 stems/ha that exceeded 10 cm dbh (Table 1 and 2). Few other taxa occurred under these dense pawpaw colonies. Hackberry, sassafras, *Carya cordiformis* (bitternut hickory), and *Ulmus americana* (American elm) were also extremely common. Many other tree species were present, but in relatively low numbers. Woody shrubs and vines were also important in the understory, *Toxicodendron radicans* (poison ivy), *Rubus pennsylvanicus* (blackberry), and *Ribes missouriense* (Missouri gooseberry) being the most common (Table 2).

Dead-standing individuals averaged 43.5 stems/ha with a basal area of 5.059 m²/ha, the most important being black oak, sassafras and American elm (Table 3). Sassafras exceeded black oak in the number of dead-standing stems/ha, but dead-standing black oak was responsible for most of the basal area (4.188 of 5.059 m²/ha). Dead-downed trees were common and averaged 46.7 stems/ha with a basal area of 6.115 m²/ha. Black oak was the most important taxon in this category accounting for 37.3 stems/ha and 5.781 m²/ha of basal area (Table 3).

DISCUSSION

Using Government Land Office survey records, Rodgers and Anderson (1979) described the presettlement vegetation of Mason County. They found that black oak was the dominant woody species in open forest communities, where it usually accounted for half of the IV. Species diversity was high in these open forests with *Carya* spp. (hickories), *Acer* spp. (maples), *Quercus alba* (white oak), *Ulmus* spp. (elms), and *Fraxinus* spp. (ash) following black oak in IV. The many small diameter witness trees reported in the GLO survey indicates that oaks and hickories were reproducing, and these relatively shade-intolerant species were replacing themselves (Rodgers and Anderson 1979).

Tomlin Timber is probably very different today compared to the early 1800s due to a reduced fire frequency followed by a total absence of fire in recent decades (Taft 1997). Frequent fires in presettlement times were probably responsible for maintaining a relatively open forest with a reduced understory (Ebinger and McClain 1991, McClain and Elzinga 1994). Their thick bark and ability to reproduce by sprouts gave oaks a competitive advantage in areas of high fire frequencies. Fire frequency and intensity dictated oak density in this presettlement landscape (Anderson 1991, Abrams 1992). The compositional stability of these open forest communities indicates that the open habitat necessary for the reproduction of these species was being maintained, probably by fire.

When surveyed by the Illinois Natural Areas Inventory in 1976, the preserve was considered an old-growth "grade A" dry upland sand forest due to the "excellent timber of good size, height, and form" (Wallace and Rowe 1976). Tree density averaged 244 stems/ha with a basal area of 27.6 m²/ha. Black oak was the dominant overstory species with 120 stems/ha and a basal area of 22.0 m²/ha, most of the individuals in the 50 to 70 cm diameter classes. Sassafras, *Carya tomentosa* (mockernut hickory), and black hickory followed in importance. The sapling layer averaged 3,900 stems/ha, sassafras being the dominant species with 1,700 stems/ha (Wallace and Rowe 1976).

Most of the forests previously studied in the Illinois River sand deposits were closed canopy dry sand forests located on dune deposits where black oak and *Quercus marilandica* (blackjack oak) were usually the leading dominants along with a few hickory species. Black hickory occasionally replaced blackjack oak as second in IV in those forests, while sassafras and elms were not present. All probably represented sand savannas that have since become closed forests due to fire suppression (Jenkins et al. 1991, Coates et al. 1992, McClain et al. 2002).

Tomlin Timber was probably an open, dry-mesic sand forest (woodland) community in presettlement times. Canopy closure and the increase in importance of mesic trees, resulting from fire suppression, has altered the structure of Tomlin Timber. With canopy closure shade-intolerant black oaks could not effectively reproduce. Sassafras, a fire-sensitive, but relatively shade-tolerant species, became the dominant understory species, eventually entered the canopy and now is the dominant species.

The high mortality of black oak observed in the woods indicates that this species was susceptible to oak wilt (Henry et al. 1944). Oak wilt disease was observed in the woods in the early 1980s, and several large diameter black oaks were killed (W.E. McClain, personal observation). The death of large oaks created canopy openings that were filled by sassafras. Though the growth of sassafras is not rapid, this species has a relatively high gap-phase-replacement potential and commonly reproduces by root suckers.

At the present time no black oaks were observed in the sapling or 10-20 cm diameter class, suggesting that this species will continue to decrease in importance at the veteran trees die. Also, few individuals of black oak were in the 20-29 cm diameter class. Tomlin timber is another example of oaks being replaced by more mesic tree species due

to a very reduced fire frequency (Ebinger and McClain 1991, Taft 1977). The loss of dominance by black oak since the 1976 inventory has been profound. This site no longer qualifies as an "old growth grade A" forest community.

LITERATURE CITED

- Abrams, M. D. 1992. Fire and the development of oak forests. *BioScience* 42:346-353.
- Anderson R. C. 1991. Presettlement forest of Illinois. pages 9-19. in G. V. Burger, J. E. Ebinger, and G. S. Wilhelm (editors). *Proceedings of the Oak Woods Management Workshop*. Eastern Illinois University, Charleston, Illinois.
- Calsyn, D.E. 1995. Soil survey of Mason County, Illinois. Soil Report 146, University of Illinois Agricultural Experiment Station, Urbana. ix+211 pp.
- Coates, D. T., S. E. Jenkins, J. E. Ebinger, and W. E. McClain. 1992. Woody vegetation survey of Barkhausen Woods, a closed canopy sand forest in Mason County, Illinois. *Erigenia* 12:1-6.
- Ebinger, J. E. and W. E. McClain. 1991. Forest succession in the prairie peninsula of Illinois. *Illinois Natural History Survey Bulletin* 34:375-381.
- Gleason, H. A. 1910. The vegetation of the inland sand deposits of Illinois. *Bulletin of the Illinois State Laboratory of Natural History* 9:21-174.
- Gleason, H. A. and A. Cronquist. 1991. *Manual of vascular plants of northeastern United States and adjacent Canada*. 2nd edition. The New York Botanical Garden, Bronx, New York.
- Hart, C. A. and H. A. Gleason. 1907. On the biology of the sand areas of Illinois. *Bulletin of the Illinois State Laboratory of Natural History* 7:137-272.
- Henry, B. W., C. S. Moses, C. A. Richards, and A. J. Riker. 1944. Oak wilt, its significance, symptoms, and cause. *Phytopathology* 34:636-645.
- Jenkins, S. E., J. E. Ebinger, and W. E. McClain. 1991. Woody vegetation survey of Bishop's Woods, a sand forest in Mason County, Illinois. *Transactions of the Illinois State Academy of Science* 84:20-27.
- King, J. E. 1981. Late Quaternary vegetational history of Illinois. *Ecological Monographs* 51:43-62.
- Maier, C. T. 1976. An annotated list of the vascular plants of Sand Ridge State Forest, Mason County, Illinois. *Transactions of the Illinois State Academy of Science* 69:153-175.
- McClain, W. E. and S. L. Elzinga. 1994. The occurrence of prairie and forest fires in Illinois and other Midwestern states, 1679 to 1854. *Erigenia* 13:79-90.

McClain, W. E., S. D. Turner, and J. E. Ebinger. 2002. Vegetation of forest communities at the Sand Prairie-Scrub Oak Nature Preserve, Mason County, Illinois. *Transactions of the Illinois State Academy of Science*. 95:37-46.

McIntosh, R.P. 1957. The York Woods. A case history of forest succession in southern Wisconsin. *Ecology* 38:29-37.

Midwestern Regional Climate Center. 2004. <http://mcc.sws.uiuc.edu>

Mohlenbrock, R. H. 2002. *Vascular Flora of Illinois*. Southern Illinois University Press, Carbondale and Edwardsville. X+490 pp.

Rodgers, C. S. and R. C. Anderson. 1979. Presettlement vegetation of two prairie peninsula counties. *Botanical Gazette* 140:232-240.

Schwegman, J.E. 1977. An annotated list of the flowering plants of Sand Prairie-Scrub Oak Nature Preserve. Unpublished report submitted to the Illinois Department of Conservation, Springfield, Illinois map+26pp.

Schwegman, J. E. 1973. Comprehensive plan for the Illinois nature preserves system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, Illinois.

Taft, J. B. 1997. Savanna and open woodland communities. Pages 24-54. in M. W. Schwartz (editor). *Conservation in highly fragmented landscapes*. Chapman and Hall, New York.

Wallace, D. and E. Rowe. 1976. Mattson Woods. Illinois Natural Areas Inventory final field survey form. Unpublished report on file at the Illinois Department of Natural Resources, Springfield, Illinois.

White, J. 1978. Illinois Natural Areas Inventory. Technical report. Volume I. Survey methods and results. Illinois Natural Areas Inventory, Urbana. xix+426 pp.

Willman, H. B. and J. C. Frye. 1970. Pleistocene stratigraphy of Illinois. *Illinois State Geological Survey Bulletin* 94:1-204.

Table 1. Densities (stems/ha), diameter classes (cm), basal areas (m²/ha), relative values, importance values and average diameters of the woody species at Tomlin Timber Nature Preserve, Mason County, Illinois.

Species	Diameter Classes (cm)							Total #/ha	Basal Area			Rel. Den.	Rel. Dom.	I. V.	AV. Diam. (cm)
	10-19	20-29	30-39	40-49	50-59	60-69	70+		m ² /ha	Den.	Dom.				
<i>Sassafras albidum</i>	41.1	59.8	26.7	4.2	0.7	--	--	132.5	6.898	26.5	28.4	54.9	24.4		
<i>Quercus velutina</i>	--	0.7	0.9	1.1	5.8	11.6	5.3	25.4	8.102	5.1	33.4	38.5	62.7		
<i>Carya texana</i>	51.6	34.0	3.8	0.7	--	--	--	90.1	2.905	18.1	12.0	30.1	19.3		
<i>Celtis occidentalis</i>	64.7	13.1	2.0	0.7	--	0.2	--	80.7	1.900	16.2	7.8	24.0	16.2		
<i>Prunus serotina</i>	42.0	8.0	2.2	0.4	--	--	0.2	52.8	1.380	10.6	5.7	16.3	16.7		
<i>Ulmus americana</i>	31.8	5.8	--	--	--	--	--	37.6	0.791	7.5	3.3	10.8	15.7		
<i>Carya tomentosa</i>	13.3	10.0	3.6	0.2	--	--	--	27.1	1.051	5.4	4.3	9.7	20.9		
<i>Ulmus rubra</i>	24.4	4.4	0.4	--	--	--	--	29.2	0.626	5.9	2.6	8.5	15.7		
<i>Asimina triloba</i>	14.9	--	--	--	--	--	--	14.9	0.166	3.0	0.7	3.7	11.8		
Others ¹ (10 species)	4.2	1.3	1.1	0.9	0.2	0.2	--	7.9	0.466	1.7	1.8	3.5			
Total	288.0	137.1	40.7	8.2	6.7	12.0	5.5	498.2	24.285	100.0	100.0	200.0			

1. Includes *Juglans nigra* (black walnut), *Carya cordiformis* (bitternut hickory), *Morus rubra* (red mulberry), *Morus alba* (white mulberry), *Acer saccharum* (sugar maple), *Robinia pseudoacacia* (black locust), *Catalpa bionoides* (catalpa), *Acer saccharinum* (silver maple), *Maclura pomifera* (Osage orange), and *Diospyros virginiana* (persimmon).

Table 2. Density (stems/ha) of woody understory species at Tomlin Timber Nature Preserve, Mason County, Illinois.

Species	Seedlings	Small Saplings	Large Saplings
<i>Asimina triloba</i>	4028	2986	854
<i>Celtis occidentalis</i>	2000	944	115
<i>Sassafras albidum</i>	1806	625	103
<i>Cary cordiformis</i>	1667	167	25
<i>Ulmus americana</i>	1222	56	18
<i>Ulmus rubra</i>	361	42	21
<i>Quercus velutina</i>	333	--	--
<i>Carya texana</i>	250	--	8
<i>Maclura pomifera</i>	139	--	11
<i>Morus alba</i>	111	28	--
<i>Prunus serotina</i>	83	14	61
<i>Gleditsia triacanthos</i>	28	--	--
<i>Jugland nigra</i>	--	--	4
<i>Acer negundo</i>	--	--	1
<i>Morus rubra</i>	--	--	1
<i>Toxicodendron radicans</i>	2194	--	--
<i>Rubus pensilvanica</i>	1944	--	--
<i>Ribes missouriense</i>	1306	--	--
<i>Sambus canadensis</i>	389	--	--
<i>Rubus occidentalis</i>	278	--	--
<i>Zanthoxylum americanum</i>	250	--	--
<i>Rubus flagellaris</i>	139	--	--
<i>Rosa multiflora</i>	83	--	--
<i>Lonicera maackii</i>	28	--	--
Totals	18639	4862	1222

Table 3. Density (stems/ha), basal area (m²/ha), and average diameter of the dead-standing and dead-downed tree species at Tomlin Timber Nature Preserve, Mason County, Illinois.

Tree Species	DEAD-STANDING			DEAD-DOWNED		
	Density (stems/ha)	Basal Area (m ² /ha)	Average Diameter (cm)	Density (stems/ha)	Basal Area (m ² /ha)	Average Diameter (cm)
<i>Quercus velutina</i>	14.4	4.188	59.2	37.3	5.781	43.1
<i>Sassafras albidum</i>	17.1	0.532	19.2	3.6	0.158	23.0
<i>Ulmus rubra</i>	4.2	0.083	15.5	2.9	0.079	18.3
<i>Ulmus americana</i>	2.7	0.085	19.3	1.8	0.059	20.0
<i>Prunus serotina</i>	2.0	0.057	17.7	0.2	0.010	23.8
<i>Carya tomentosa</i>	0.9	0.055	26.0	--	--	--
<i>Carya texana</i>	0.9	0.033	21.1	0.9	0.028	19.0
<i>Asimina triloba</i>	0.9	0.011	12.6	--	--	--
<i>Acer saccharinum</i>	0.4	0.015	20.8	--	--	--
Totals	43.5	5.059		46.7	6.115	

APPENDIX I. Vascular species encountered at Tomlin Timber Nature Preserve, Mason County, Illinois, listed alphabetically by family under the major plant groups. An asterisk indicates non-native (exotic) species (*). Following the scientific name, collecting numbers preceded by the initial of the collector's name are given (E) James Ellis and (P) Loy R. Phillippe.

PTERIDOPHYTES

DRYOPTERIDACEAE

Dryopteris cartusiana (Villars) H.P. Fuchs: P 36729

ONOCLEACEAE

Onoclea sensibilis L.: P 36064

OPHIOGLOSSACEAE

Botrychium virginianum (L.) Sw.: P 36680

SPERMATOPHYTES: GYMNOSPERMS

CUPRESSACEAE

Juniperus virginiana L.: P 36049

SPERMATOPHYTES: ANGIOSPERMS

MONOCOTS

ARACEAE

Arisaema triphyllum (L.) Schott: P 36685

COMMELINACEAE

**Commelina communis* L.: P 35988
Tradescantia ohiensis Raf.: P 35987

CYPERACEAE

Carex blanda Dewey: P 36678
Carex festucacea Schk.: P 36707, P 36728, P 36732
Carex grisea Wahl: P 36679, P 36690
Cyperus lupulinus (Spreng.) Marcks var. *macilentus* (Fern.) Marcks: P 35982

JUNCEAE

Juncus tenuis Willd.: P 36041

LILIACEAE

Lilium michiganense Farw.: P 36061
**Ornithogalum umbellatum* L.: P 36676
Smilacina racemosa (L.) Desf.: P 36014
Smilacina stellata (L.) Desf.: E 53, P 36013

POACEAE

**Bromus inermis* Leyss.: P 36068
**Bromus tectorum* L.: P 36688
**Dactylis glomerata* L.: P 36725
**Digitaria sanguinalis* (L.) Scop.: P 36040
Elymus villosus Muhl.: P 36043
Elymus virginicus L.: P 36025
Eragrostis spectabilis (Pursh) Steud.: P 35985

Festuca subverticillata (Pers.) E.B. Alexeev: P 36057
Hordeum pusillum Nutt.: P 36727
Leersia virginica Willd.: P 36061
Muhlenbergia frondosa (Poir.) Fern.: E 55
Muhlenbergia schreberi J.F. Gmel.: P 36032
Paspalum bushii Nash: P 35976
Poa pratensis L.: P 36699, P 36701
**Poa sylvestris* Gray: P 36677
**Setaria faberi* R.A.W. Herrm.: P 35998
Tridens flavus (L.) Hitchc.: P 35981

SMILICACEAE

Smilax tamnoides L. var. *hispida* (Muhl.) Fern.: P 36004
Smilax lasioneuron Hook.: P 36054

DICOTS

ACANTHACEAE

Ruellia humilis Nutt.: P 35978

ACERACEAE

Acer negundo L.: P 37298
Acer saccharinum L.: P 37360
Acer saccharum Marsh.: P 37356

ANACARDIACEAE

Toxicodendron radicans (L.) Kuntze: P 36003

ANNONACEAE

Asimina triloba (L.) Dunal: P 36016

APIACEAE

Chaerophyllum procumbens (L.) Crantz: P 36674
Osmorhiza longistylis (Torr.) DC.: P 36672.2
Sanicula canadensis L.: P 36011
Sanicula odorata (Raf.) Pryer & Phillippe: P 36021

ASTERACEAE

Ageratina altissima (L.) R.M. King & H. Rob.: P 36010
Ambrosia artemisiifolia L.: P 36036
Ambrosia trifida L.: P 36047
Arnoglossum atriplicifolia (L.) H. Rob.: P 35972
Aster ontarionis Wieg.: E 58, E 59
Aster pilosus Willd.: E 48
Bidens bipinnata L.: P 35979
Conyza canadensis (L.) Cronq.: P 36035
Eupatoriadelphus purpureus (L.) R.M. King & H. Rob.: P 36052
Eupatorium serotinum Michx.: P 36039
Helianthus divaricatus L.: P 36006
Heterotheca subaxillaris (Lam.) Britt. & Rusby: P 35986
Lactuca canadensis L.: P 36050
Lactuca floridana (L.) Gaertn.: P 36020
Rudbeckia hirta L.: P 35993
Solidago canadensis L.: P 35991
Solidago gigantea Ait.: E 51

BERBERIDACEAE

Podophyllum peltatum L.: P 36683

BIGNONIACEAE

**Catalpa bignonioides* Walt.: P 37357

BORAGINACEAE

**Buglossoides arvense* (L.) I.M. Johnston: P 36696
Hackelia virginiana (L.) I.M. Johnston: P 35994

BRASSICACEAE

**Alliaria petiolata* (Bieb.) Cavara & Grande: P 36673
Arabis glabra (L.) Bernh.: P 36730
**Capsella bursa-pastoris* (L.) Medik.: P 36693

CAESALPINIACEAE

Gleditsia triacanthos L.: P 36056

CANNABINACEAE

**Cannabis sativa* L.: P 36029
Humulus lupulus L.: P 36019

CAPRIFOLIACEAE

**Lonicera maackii* (Rupr.) Maxim.: P 36682
Sambucus canadensis L.: P 36051

CARYOPHYLLACEAE

**Arenaria serpyllifolia* L.: P 36691
**Cerastium pumilum* Curtis: P 36697
**Holosteum umbellatum* L.: P 36702
**Saponaria officinalis* L.: P 35983
Silene stellata (L.) Ait. f.: P 36009

CELASTRACEAE

Celastrus scandens L.: P 36048

CHENOPODIACEAE

**Chenopodium album* L.: E 49

CORNACEAE

Cornus drummondii C.A. Mey.: P 36007

CORYLACEAE

Corylus americana Walt.: P 36065

EBENACEAE

Diospyros virginiana L.: P 37361

ELAEAGNACEAE

**Elaeagnus umbellata* Thunb.: E 57, P 36723

EUPHORBIACEAE

Acalypha gracilens Gray: P 35974
Acalypha virginica L.: P 35975
Chamaesyce nutans (Lag.) Small: P 36053
Croton glandulosus L.: P 35973
Phyllanthus caroliniensis Walt.: P 36059
Poinsettia dentata Michx.: E 52P 35977

FABACEAE

**Medicago lupulina* L.: P 36726
**Melilotus albus* Medic.: P 35992
**Robinia pseudoacacia* L.: P 37297

FAGACEAE

Quercus velutina Lam.: P 36070

GROSSULARIACEAE

Ribes missouriense Nutt.: P 36012

HYDROPHYLLACEAE

Ellisia nyctelea L.: P 36681

JUGLANDACEAE

Carya cordiformis (Wangenh.) K. Koch: P 36026
Carya illinoensis (Wangenh.) K. Koch: P 36063
Carya texana Buckl.: P 36001
Carya tomentosa (Poir.) Nutt.: P 36018
Juglans nigra L.: P 36071

LAMIACEAE

Agastache nepetoides (L.) Ktze.: P 36008
**Leonurus cardiaca* L.: P 36722
**Nepeta cataria* L.: P 36005
Teucrium canadense L.: P 36055

LAURACEAE

Sassafras albidum (Nutt.) Nees.: P 36000

MALVACEAE

Callirhoë triangulata (Leavenw.) Gray: P 35970

MOLLUGINACEAE

**Mollugo verticillata* L.: P 36038

MORACEAE

**Maclura pomifera* (Raf.) Schneider: P 36066
**Morus alba* L.: P 36045
Morus rubra L.: P 36045

NYCTAGINACEAE

**Mirabilis nyctaginea* (Michx.) MacM.: P 35996

ONAGRACEAE

Circaea lutetiana L.: P 36024
Oenothera biennis L.: P 36042
Oenothera laciniata Hill: P 36704

OXALIDACEAE

Oxalis stricta L.: P 35989

PHRYMACEAE

Phryma leptostachya L.: E 56, P 36023

PHYTOLACCACEAE

Phytolacca americana L.: P 35984

PLANTAGINACEAE

Plantago rugelii Decne.: P 35997
Plantago virginica L.: P 36731

POLYGALACEAE

Polygala verticillata L.: P 35971

POLYGONACEAE

Antenoron virginianum (L.) Roberty & Vautier: P 36030
Fallopia scandens (L.) Holub.: E 50
Persicaria punctata (Eil.) Small: P 36028

PORTULACCACEAE

Claytonia virginica L.: P 36686

RANUNCULACEAE

Anemone virginiana L.: P 36037
Ranunculus abortivus L.: P 36687

ROSACEAE

Fragaria virginiana Duchesne: P 36689
Geum canadense Jacq.: P 35995
Prunus serotina Ehrh.: P 36067
Rosa carolina L.: P 36724
**Rosa multiflora* Thunb.: E 54
Rubus flagellaris Willd.: P 36698
Rubus occidentalis L.: P 36706
Rubus pensilvanicus Poir.: P 36705

RUBIACEAE

Galium aparine L.: P 36675

RUTACEAE

Zanthoxylum americanum Mill.: P 37359

SCROPHULARIACEAE

Penstemon pallidus Small: P 36703
Scrophularia lanceolata Pursh: P 36027

**Verbascum thapsus* L.: P 36033
Veronica arvensis L.: P 36694

SOLANACEAE

Physalis heterophylla Nees: P 35980
Physalis subglabrata Mack. & Bush: P 35999
Solanum caroliniense L.: P 35990
Solanum ptychanthum Dunal: P 36058

ULMACEAE

Celtis occidentalis L.: P 36002
Ulmus americana L.: P 37358
Ulmus rubra Muhl.: P 36044

URTICACEAE

Laportea canadensis (L.) Wedd.: P 36022

VERBENACEAE

Verbena stricta Vent.: P 36015
Verbena urticifolia L.: P 36069

VIOLACEAE

Viola pratincola Greene: P 36684
**Viola rafinesquei* Greene: P 36072
Viola sororia Willd.: P 36695

VITACEAE

Vitis aestivalis Michx.: P 36034
Vitis riparia Michx.: P 36060

Changes in a sand savanna due to disturbance and fire suppression, Sand Ridge State Forest, Mason County, Illinois.

Loy R. Phillippe, Paul B. Marcum, Daniel T. Busemeyer, William E. McClain, and John E. Ebinger, Illinois Natural History Survey, Champaign, Illinois 61820

ABSTRACT - Sand savannas, dominated by *Quercus velutina* Lam. (black oak) are still relatively common in the major sand deposits of Illinois. Most, however, are extensively degraded by fire suppression and invasion by woody and exotic species. Many are now closed forests that lack characteristic ground layer species. Degraded savannas, that are presently dry sand forests, are the dominant community of ridges and slopes on large stabilized dunes at Sand Ridge State Forest, Mason County, Illinois. In the community examined black oak, with an importance value of 143.5, averaged 321.1 stems/ha, and accounted for 78% of the total basal area. *Quercus marilandica* Muench. (blackjack oak) was second followed by *Pinus strobus* L. (white pine) and *Carya texana* Buckl. (black hickory). Based on early aerial photographs this closed forest had an open overstory in the early 1940s.

INTRODUCTION

Savanna communities are generally defined as having overstories consisting of scattered, open-grown trees and a herbaceous grass dominated ground layer (Curtis 1959, Bray 1960, Nuzzo 1986, White and Madany 1978). In the prairie forest interface of the prairie peninsula of Illinois the presence of prairie, savanna, woodland, and forest communities was determined largely by environmental factors, including the extent and intensity of fire; climate, and topography (Transeau 1935, Anderson 1991, Ebinger and McClain 1991, Abrams 1992). Other contributing factors important on a local level included coarseness of the soil, frequency of local droughts, and browsing by large herbivores (Nuzzo 1986).

Savanna communities were extremely common in the landscape of Illinois in the 1800s. Journals of many early travelers and settlers recount the open park-like landscape in many parts of the state (Bourne 1820, Engelmann 1863, Vestal 1936). Government Land Office (GLO) survey records also indicate that many "forests" were actually savanna and woodland communities based on the distance of witness trees to corner posts (Cottam and Curtis 1949, Clements 1958, Hutchison 1988). Furthermore, many present day old growth forests still retain a few open-grown "wolf trees" with low branches and branch-scars, indicating that they were formerly part of an open landscape (Curtis 1959, Ebinger and McClain 1991).

European man destroyed most "black soil" savannas of Illinois soon after settlement. The few trees and thinner, often drier sod, made savannas easier to plow with the early wooden plows than "black soil" prairies. The few savanna communities that remain are extensively degraded by a massive influx of exotic species and canopy closure due to fire suppression and subsequent woody invasion.

Savanna communities are still relatively common in the northern half of Illinois on major sand deposits. These deposits are mostly on outwash plains that resulted from erosional events associated with Wisconsin glaciation (Willman and Frye 1970, King 1981). Gleason (1910), and more recently Jenkins et al. (1991), Coates et al. (1992), McClain et al. (2002) studied the structure and composition of woodland communities of the Illinois River sand deposits. Also, Rodgers and Anderson (1979) studied the presettlement vegetation of Mason

County. Mostly modified by human activity, a few nature preserves and other good quality natural areas remain. The present study was undertaken to determine the woody overstory and understory species composition and structure of a degraded savanna that still retains some of its original structure at Sand Ridge State Forest.

DESCRIPTION OF THE STUDY SITE

Sand Ridge State Forest is located in northwestern Mason County about 21 km northwest of Havana, and just west of Forest City, Illinois (parts of townships T22N R7W and T23N R7W). This 3,035 ha (11.7 sq. miles) state forest, with initial land purchases starting in 1939, lies within the Illinois River Section of the Mississippi River and Illinois River Sand Area Natural Division in Mason and Cass counties (Schwegman 1973, Willman 1973). These deposits were formed about 14,500 years ago when glacial moraines and ice dams were breached. The resulting flood, the Kankakee Torrent, carried extensive deposits of sand and gravel from glacial lakes in northeastern Illinois and adjacent Indiana. Most of this sand and gravel was deposited when the waters of the Kankakee Torrent slowed upon entering the broad lowlands of the Illinois River. These deposits were then reworked by winds creating the present dune and swale topography.

The original reason for purchasing the land for what is now the Sand Ridge States Forest was to stabilize soil on abandoned farmlands, develop a wood product industry, and set land aside for recreation (Andrews 2004). During the early years, and into the 1950's, pine plantations were established within this state forest, mostly on old pastureland and abandoned cultivated fields, but also in dry sand prairies. Some pines were also planted in savannas. Presently 1,012 ha of marketable pine plantations are found in the state forest with most of the remainder in oak-hickory dry sand forest and savanna (Andrews 2004).

Sand Ridge State Forest has a continental climate with warm summers and cold winters. Based on weather data from Havana, mean annual precipitation is 96.0 cm, with May having the highest rainfall (11.3 cm). Mean annual temperature is 10.8°C with the hottest month being July (average of 24.6°C), and the coldest January (average of -5.0°C). Frost-free days range from 140 to 206, with the average being 173 days per year (Midwestern Regional Climate Center 2004). The soils are mostly excessively drained Plainfield and Bloomfield sands (Calsyn 1995) that form the dune and swale topography known as the Parkland Formation (Willman and Frye 1970).

METHODS

During late summer of 2004 a 100 m by 300 m section of the state forest was surveyed by dividing the area into 48 contiguous quadrates 25 m on a side. This 3 ha area was located on a large stabilized dune having an east/west orientation, the centerline of the transect running along the ridge of the dune (N1/2 NW1/4 NE1/4 S4 T22N R7W). The GPS readings for the transect line are reported in Appendix II. All living and dead-standing woody individuals ≥ 10.0 cm dbh were identified and their diameters recorded. From this data, living-stem density (stems/ha), basal area (m^2/ha), relative density, relative dominance, importance value (IV), and average diameter (cm) were calculated for each species. Determination of the IV follows the procedure used by McIntosh (1957), and is the sum of the relative density and relative dominance (basal area). Dead-standing density (stem/ha) and basal area (m^2/ha) was also determined. Nomenclature follows Mohlenbrock (2002).

Woody understory composition and density (stems/ha) were determined using nested circular plots 0.0001, 0.001, and 0.01 ha in size located at about 15 meter intervals along randomly located east-west line transects within the study area

(48 plots). Four additional 0.0001 ha circular plots were located 7 m from the center points of each of the 48 plot centers along cardinal compass directions (240 plots). In the 0.0001 ha plots, woody seedlings (≤ 50 cm tall) and shrubs and vines were counted; in the 0.001 ha circular plots small saplings (> 50 cm tall and < 2.5 cm dbh) were recorded; and in the 0.01 ha circular plots large saplings (2.5-9.9 cm dbh) were tallied.

To determine the change in total overstory cover within the state forest aerial photographs from 1939, 1967, and 1988 were digitized to determine the extent of woody encroachment (trees and large shrubs). The 1939 aerial photographs were taken in July, the 1967 in May, and the 1988 in April. These photographs were borrowed from the University of Illinois Map Library and scanned with a Microtek ScanMaker. A total of 20 stratified randomly located 5 ha circular plots (100 ha total area), representing approximately 20% of the study sites, were interpreted and then digitized using ARC/INFO.

RESULTS

Eleven tree species were encountered in the overstory (Table 1). Black oak dominated all diameter classes with the 10-29 cm diameter classes accounting for more than 75% of all individuals and only 3 stems/ha greater than 60 cm dbh. This species had an IV of 143.5, averaged 321.1 stems/ha, averaged 23.6 cm dbh, and accounted for 78.1% of the total basal area. *Quercus marilandica* Muench. (blackjack oak), second in IV, was mostly restricted to smaller diameter classes, averaged 111.6 stems/ha, and averaged 16.5 cm dbh. The remaining species were mostly in the 10-39 cm diameter classes, *Carya texana* Buckl. (black hickory) averaged 26.3 stems/ha, while *Pinus strobus* L. (white pine) averaged 26.1 stems/ha. Coppice stems accounted for about 16% of the stems encountered. Black oak accounted for the majority, averaging 27 coppice trees/ha with 57.7 stems/ha (Table 2).

Dead-standing individuals averaged 24.6 stems/ha with a basal area of 1.01 m²/ha, nearly all being oaks. Black oak averaged 15.6 dead-standing stems/ha while blackjack oak accounted for nearly all of the remainder. Most of the dead-standing individuals were in the lower diameter classes and most had basal fire-scars. A few dead-standing black oaks exceeded 40 cm dbh.

The woody understory averaged 15,200 seedlings/ha, 1,775 small saplings/ha, and 295 large saplings/ha (Table 3). Seedling density was relatively high but the majority was shrubby species. Black oak and black hickory accounted for nearly all of the tree seedlings. Because of the relatively few saplings, the woody understory was open. Again, black oak and black hickory accounted for the majority of the individuals (Table 3). Woody shrubs were also important in the understory, *Rubus allegheniensis* Porter (common blackberry), *Rhus aromatica* Ait. (fragrant sumac), *Toxicodendron radicans* (L.) Kuntze (poison ivy), and *Cornus drummondii* C. A. Mey. (rough-leaved dogwood) being the most common (Table 3).

Based on aerial photographs from 1939, 1967, and 1988, the density of the woody overstory has increased dramatically. Of the 100 ha (20 plots each 5 ha in size) analyzed in 1939 all sites were very open (00.000 ha of woody overstory), by 1967 woody cover had nearly doubled (00.000 ha), while by 1988 woody cover again increased significantly (00.000 ha). This represents an increase of 00.000 ha of woody overstory in about 50 years (Table 4).

DISCUSSION

The forests of Sand Ridge State Forest are very different today compared to the early 1800s, mostly due to a reduced fire frequency followed by the total absence of fire in recent decades (Taft 1997). There has been a dramatic

increase in canopy cover in the past 65 years: from _____ in 1939 to _____ in 1988 (Table 4). In presettlement times repeated fires were probably responsible for maintaining an open forest with a sparse understory (Ebinger and McClain 1991, McClain and Elzinga 1994). In those forests the larger trees maintained an open-grown appearance with low branches and branch-scars. A few large, open-grown trees remained in the study plots. Because of fire and droughty conditions, most of this forest was originally woodland and savanna communities with numerous small, scattered prairie openings. Canopy closure resulted from fire suppression and the subsequent woody invasion by native species.

Presently occasional fires and the droughty conditions have allowed for the perpetuation of oak species. Black oak is reproducing on the site with numerous seedlings and saplings in the understory (Table 3). Blackjack oak, in contrast, has a very low rate of reproduction. The large number of seedlings, saplings, and small diameter trees of black hickory suggests that this species will increase in importance. As canopy closure continues, the shade-intolerant oaks may not effectively reproduce. Black hickory, a fire-sensitive, but relatively shade-tolerant species, could become the dominant understory species and become more common in the lower diameter classes.

Woody exotic species are common in Sand Ridge State Forest. At least 10 species of pine were planted in the 1940s and early 1950s, and many pine plantations are present (Maier 1976, Andrews 2004). The most commonly planted species was white pine. Rows of individuals of this introduced exotic species were present in our study plots, indicating this species was also planted in the native hardwood forests and savannas. Smaller individuals, plus occasional seedlings indicate that this species is reproducing.

Using GLO survey records, Rodgers and Anderson (1979) described the presettlement vegetation of Mason County. They found that tree density averaged 7.44 trees/ha with an average basal area of 1.19 m²/ha in savanna communities. Black oak was, by far, the dominant woody species, accounting for more than half of the IV. Blackjack oak was second in IV followed by various species of hickory. The many small diameter witness trees reported in the GLO survey indicate that oaks and hickories were reproducing, and these relatively shade-tolerant species were replacing themselves in savanna, woodland, and closed forest communities (Rodgers and Anderson 1979).

Most forests studied within the Illinois River sand deposits were closed canopy dry sand forests located on dune deposits where black and blackjack oaks were usually the leading dominants along with a few hickory species. Black hickory occasionally replaced blackjack oak as second in IV in those forests (Jenkins et al. 1991, Coates et al. 1992, McClain et al. 2002). All probably represented sand savannas that have become closed forests due to fire suppression and woody species invasion. An increased fire frequency and some timber harvesting will be necessary to restore the woodland and savanna communities that once existed at Sand Ridge State Forest.

ACKNOWLEDGMENTS

The authors thank John Wilker, Natural Heritage Biologist, Illinois Department of Natural Resources, for his help and advice. The Illinois Department of Natural Resources, Wildlife Preservation Fund supported this project.

LITERATURE CITED

Abrams, M. D. 1992. Fire and the development of oak forests. *BioScience* 42:346-353.

- Anderson R. C. 1991. Presettlement forest of Illinois. pages 9-19. in G. V. Burger, J. E. Ebinger, and G. S. Wilhelm (editors). Proceedings of the Oak Woods Management Workshop. Eastern Illinois University, Charleston, Illinois.
- Andrews, K. 2004. Forest Treasures. Outdoor Illinois. December 2004:2-5.
- Bourne, A. 1820. On the prairies and barrens of the west. American Journal of Science and Arts 2:30-34.
- Bray, J.R. 1960. The composition of savanna vegetation in Wisconsin. Ecology 41:721-732.
- Calsyn, D.E. 1995. Soil survey of Mason County, Illinois. Soil Report 146, University of Illinois Agricultural Experiment Station, Urbana. ix+211 pp.
- Clements, D.B. 1958. Public land surveys-history and accomplishments. Survey and Mapping 18:213-219.
- Coates, D. T., S. E. Jenkins, J. E. Ebinger, and W. E. McClain. 1992. Woody vegetation survey of Barkhausen Woods, a closed canopy sand forest in Mason County, Illinois. *Erigenia* 12:1-6.
- Cottam, G. and J.T. Curtis. 1949. A method for making rapid surveys of woodlands by means of pairs of randomly selected trees. Ecology 30:101-104.
- Curtis, J.T. 1959. The vegetation of Wisconsin. The University of Wisconsin Press. Madison. 657 pp.
- Ebinger, J. E. and W. E. McClain. 1991. Forest succession in the prairie peninsula of Illinois. Illinois Natural History Survey Bulletin 34:375-381.
- Engelmann, H. 1863. Remarks upon the causes producing the different characters of vegetation known as prairie, flats, and barrens in southern Illinois, with special reference to observations made in Perry and Jackson counties. American Journal of Science 86:384-396.
- Gleason, H. A. 1910. The vegetation of the inland sand deposits of Illinois. Bulletin of the Illinois State Laboratory of Natural History 9:21-174.
- Hutchison, M. 1988. A guide to understanding, interpreting, and using the Public Land Survey field notes in Illinois. Natural Areas Journal 8:245-255.
- Jenkins, S.E., J.E. Ebinger, and W.E. McClain. 1991. Woody vegetation survey of Bishop's Woods, a sand forest in Mason County, Illinois. Transactions of the Illinois State Academy of Science 84:20-27.
- King, J.E. 1981. Late Quaternary vegetational history of Illinois. Ecological Monographs 51:43-62.
- Maier, C.T. 1976. An annotated list of the vascular plants of Sand Ridge State Forest, Mason County, Illinois. Transactions of the Illinois State Academy of Science 69:153-175.
- McClain, W. E. and S. L. Elzinga. 1994. The occurrence of prairie and forest fires in Illinois and other Midwestern states, 1679 to 1854. *Erigenia* 13:79-90.

- McClain, W.E., S.D. Turner, and J.E. Ebinger. 2002. Vegetation of forest communities at the Sand Prairie-Scrub Oak Nature Preserve, Mason County, Illinois. *Transactions of the Illinois State Academy of Science* 95:37-46.
- McIntosh, R.P. 1957. The York Woods. A case history of forest succession in southern Wisconsin. *Ecology* 38:29-37.
- Midwestern Regional Climate Center. 2002. <http://mcc.sws.uiuc.edu>
- Mohlenbrock, R.H. 2002. Vascular Flora of Illinois. Southern Illinois University Press, Carbondale, Illinois. xi+490 pp.
- Nuzzo, V.A. 1986. Extent and status of mid-west oak savanna: presettlement and 1985. *Natural Areas Journal* 6:6-36.
- Rodgers, C.S. and R.C. Anderson. 1979. Presettlement vegetation of two prairie peninsula counties. *Botanical Gazette* 232-240.
- Schwegman, J.E. 1973. Comprehensive plan for the Illinois nature preserves system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Rockford, Illinois. map+32 pp.
- Taft, J. B. 1997. Savanna and open woodland communities. Pages 24-54. in M. W. Schwartz (editor). *Conservation in highly fragmented landscapes*. Chapman and Hall, New York.
- Transeau, E.N. 1935. The prairie peninsula. *Ecology* 16:423-437.
- Vestal, A.G. 1936. Barrens vegetation in Illinois. *Transactions of the Illinois State Academy of Science* 29:29-30.
- Willman, H.B. 1973. Geology along the Illinois waterway - a basis for environmental planning. Illinois State Geological Survey Circular 478. Urbana. 48 pp.
- Willman, H.B. and J.C. Frye. 1970. Pleistocene stratigraphy of Illinois. *Illinois State Geological Survey Bulletin* 94:1-204.
- White, J. and M.H. Madany. 1978. Classification of natural communities in Illinois. Pp. 310-405 in Illinois natural areas inventory. Technical report. (J. White, Editor). Illinois Natural Areas Inventory, Urbana, Illinois.

Table 1. Densities (stems/ha), diameter classes, basal areas (m²/ha), relative values, importance values and average diameters of the woody species at Sand Ridge State Forest, Mason County, Illinois.

Species	Diameter Classes (cm)						Total #/ha	Basal Area m ² /ha	Rel. Den.	Rel. Dom.	I.V.	Av. Diam. (cm)
	10-19	20-29	30-39	40-49	50-59	60+						
<i>Quercus velutina</i>	145.3	107.7	37.7	20.7	6.7	3.0	321.1	16.995	65.4	78.1	143.5	23.6
<i>Quercus marilandica</i>	92.0	17.0	2.3	0.3	--	--	111.6	2.601	22.8	11.9	34.7	16.5
<i>Pinus strobus</i>	12.7	8.7	2.7	2.0	--	--	26.1	1.243	5.3	5.7	11.0	23.0
<i>Carya texana</i>	20.0	3.3	1.7	1.0	0.3	--	26.3	0.849	5.4	3.9	9.3	17.9
Others (7 species*)	5.7	--	--	--	--	--	5.7	0.080	1.1	0.4	1.5	
Total	275.7	136.7	44.4	24.0	7.0	3.0	490.8	21.768	100.0	100.0	200.0	

*Other species include: *Carya tomentosa* (Poir.) Nutt., *Diospyros virginiana* L., *Juniperus virginiana* L., *Pinus banksiana* Lamb., *Pinus sylvestris* L., *Prunus serotina* Ehrh., *Ulmus americana* L.

Table 2. Density (#/ha) of coppice trees and stems, coppice stems per tree, average basal area (m²/ha) of coppice stems, and the average diameter (cm) of coppice stems at Sand Ridge State Forest, Mason County, Illinois.

Species	Trees (#/ha)	Stems (#/ha)	Stems/trees	Basal Area (m ² /ha)	Average Diameter (cm)
<i>Quercus velutina</i>	27.0	57.7	2.1	2.721	23.4
<i>Quercus marilandica</i>	9.0	19.3	2.2	0.540	17.9
<i>Carya texana</i>	1.7	3.3	2.0	0.099	17.1
Totals	37.7	80.3		3.360	

Table 3. Density (individuals/ha) of woody understory species in a woodland community at Sand Ridge State Forest, Mason County, Illinois.

Species	Seedlings	Small Saplings	Large Saplings
<i>Quercus velutina</i>	3750	575	100.0
<i>Carya texana</i>	2850	600	85.0
<i>Prunus serotina</i>	250	250	20.0
<i>Quercus marilandica</i>	250	25	30.0
<i>Carya tomentosa</i>	150	125	17.5
<i>Pinus strobus</i>	150	25	17.5
<i>Juniperus virginiana</i>	--	100	15.0
<i>Pinus sylvestris</i>	--	--	5.0
<i>Ulmus americana</i>	--	--	2.5
<i>Celtis occidentalis</i>	--	--	2.5
<i>Rubus allegheniensis</i>	2250	--	--
<i>Rhus aromatica</i>	1850	--	--
<i>Toxicodendron radicans</i>	1650	--	--
<i>Cornus drummondii</i>	1600	50	--
<i>Rubus occidentalis</i>	300	--	--
<i>Ribes missouriense</i>	100	--	--
<i>Viburnum prunifolium</i>	50	--	--
<i>Lonicera maackii</i>	--	25	--
Totals	15200	1775	295.0