# horned lark Eremophila alpestris

Kingdom: Animalia Phylum: Chordata Class: Aves Order: Passeriformes Family: Alaudidae

### **ILLINOIS STATUS**

common, native

# ILLINOIS RANGE



### FEATURES

The horned lark averages seven to eight inches in length (tail tip to bill tip in preserved specimen). The male is a brown bird with black "sideburns," two, small black horns and a black breast splotch. The female and immature are similarly patterned but not as striking in coloration. This bird has an extended claw on its rear-facing toes.

#### **BEHAVIORS**

The horned lark is a common, permanent resident statewide. It lives in open areas, such as fields, pastures, golf courses, airports and along roads. This bird can fly but is often seen walking. In the spring, it flies high into the air and sings an irregular, highpitched song. The breeding season occurs from February through July. The nest is built on the ground, usually in a shallow depression. This nest of plant materials lined with grasses is built by the female in two to four days. One side of the nest may have small stones or dirt clods. Four, green or gray eggs with dark markings are deposited by the female, and she alone incubates them for the 11-day incubation period. Two broods are produced per year. The horned lark eats grains, grass seeds and insects.

© Illinois Department of Natural Resources. 2021. *Biodiversity of Illinois*. Unless otherwise noted, photos and images © Illinois Department of Natural Resources.





© Illinois Department of Natural Resources. 2021. *Biodiversity of Illinois*. Unless otherwise noted, photos and images © Illinois Department of Natural Resources.



adult

### **Aquatic Habitats**

wet prairies and fens

### **Woodland Habitats**

none

# Prairie and Edge Habitats

black soil prairie; dolomite prairie; edge; gravel prairie; hill prairie; sand prairie; shrub prairie