

# mourning dove

## *Zenaida macroura*

Kingdom: Animalia  
Phylum: Chordata  
Class: Aves  
Order: Columbiformes  
Family: Columbidae

### ILLINOIS STATUS

common, native



adult at nest

### FEATURES

The mourning dove averages about 12 inches in length. It has gray-brown body feathers with some dark spots at the base of the wings. The pointed tail feathers have white edging.

### BEHAVIORS

The mourning dove is a common migrant and summer resident statewide. It is also a common winter resident statewide. Spring migration into Illinois begins in late April. Nesting occurs from March through September. This bird nests in trees, especially evergreens, about 10 to 25 feet above the ground. The nest is a flimsy-looking platform of sticks. This dove may place its twig nest over an old nest of another bird. Two white eggs are laid by the female. The male (during the day) and the female (at night) incubate for the 13- to 14-day incubation period. Several broods are raised each year. Many doves seen in Illinois in late summer are migrants from farther north. Illinois doves begin migrating in September and winter in Georgia, Florida, Louisiana, Texas and Central America. Some doves do overwinter in Illinois. The mourning dove may be found in open habitats including shrubs, hedgerows, clover fields, grasslands, orchards, marshes, urban areas and open woods. It makes a “coah, cooo, cooo, coo” sound. This bird feeds on foxtail grass, corn, wheat, crabgrass, prairie grass and other plant materials.

# ILLINOIS RANGE



juvenile



© Illinois Department of Natural Resources. 2020. *Biodiversity of Illinois*.  
Unless otherwise noted, photos and images © Illinois Department of Natural Resources.



adult

**Aquatic Habitats**

bottomland forests

**Woodland Habitats**

bottomland forests; coniferous forests; upland deciduous forests

**Prairie and Edge Habitats**

none